



Lifestyles and their Influence on Sleep Disorders

Shashi K. Agarwal ^{a*}

^a 2227 US Highway 1, Suite 309, North Brunswick, NJ 08902, USA.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/INDJ/2021/v16i330182

Editor(s):

(1) Dr. Takashi Ikeno, National Institute of Mental Health & National Center of Neurology and Psychiatry, Japan.

Reviewers:

(1) Julia Nowowiejska, Medical University of Bialystok, Poland.

(2) Mohammad Maroof Siddiqui, Dhofar University, Oman.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/83544>

Received 15 October 2021

Accepted 19 December 2021

Published 20 December 2021

Mini-review Article

ABSTRACT

Sleep is a complex physiological event that occupies nearly one-third of human life. Poor sleep adversely affects several bodily systems. Prolonged sleep deprivation can lead to premature death. Adequate sleep duration and sleep quality are important for a healthy life. For most adults, sleep duration of 7 to 9 hours is healthy. Going to sleep rapidly, sleeping undisturbed through the night, and waking up refreshed are important qualities that should accompany an ideal sleep duration. The determinants of sleep duration and quality are diverse and include several physical and mental health factors. Lifestyle factors also play an important role. The five major lifestyles that adversely impact human health are smoking, obesity, excess alcohol consumption, lack of physical exercise, and an unhealthy diet. These can also disturb sleep duration and sleep quality, leading to sleep disorders. This manuscript provides a narrative review of the various interactions between these lifestyles and human sleep.

Keywords: Sleep disorders; smoking; lifestyles; exercise; alcohol; diet; obesity.

1. INTRODUCTION

Sleep is vital for life. It helps repair the body, allows tissue regrowth, rebuilds bone and muscle, strengthens the immune system, and allows memory consolidation [1-3]. Restorative

sleep is characterized by a sleep duration of between 7-9 hours per day, falling asleep easily, staying asleep without interruption, waking up feeling refreshed, and staying awake (without sleepiness, lack of alertness, fatigue, or irritability) during the day [4-6]. Sleep duration

*Corresponding author: E-mail: usacardiologist@gmail.com;

and quality, however, vary according to age – babies may sleep 16 to 18 hours per day, schoolchildren need about 9.5 hours of sleep per night, while at age 60, sleep is shorter, lighter, and may be interrupted by multiple awakenings [7,8]. Poor sleep (sleep disorder) is commonly due to insomnia, sleep apnea, restless leg syndrome, narcolepsy, REM sleep behavior disorder, and circadian-related sleep disorder [9]. These can cause several problems. Children may become overly active, misbehave, be less attentive and suffer in school performance. They may also appear angrier, exhibit impulsiveness, have mood swings, and show a lack of motivation [10]. In adults, improper sleep may result in decreased work efficiency, absenteeism, accidents, and low productivity [11]. Personal and social life is also often affected. The overall quality of life is diminished [12]. Sleep disorders are often present in many chronic diseases [13-20]. These include cardiovascular diseases (CVD) [13], type 2 diabetes mellitus (T2DM) [14], asthma [15], hepatitis [16], chronic kidney disease (CKD) [17], metabolic disorders [18], obesity [19], and even some cancers [20]. Patients with insomnia or obstructive sleep apnea (OSA) demonstrate an increased risk for several CVDs including, arrhythmias, atherosclerosis, coronary heart disease, heart failure, hypertension, and stroke. They also increase the risk of several CVD causing metabolic disorders such as obesity, T2DM, and dyslipidemia [13]. The relationship between T2DM and sleep disturbances has been well studied. Sleep disorders increase the risk of T2DM by altering insulin sensitivity resulting in lower glucose tolerance. Hypoxia associated with several sleep disorders may lead to a lower pancreatic β -cell function. Disordered sleep may also increase the risk for T2DM by modulating the sympathetic nervous system, cortisol levels, melatonin actions and by increasing inflammation [14]. OSA is more common in patients with asthma, and CPAP treatments help improve comorbid asthma in these patients. Insomnia is also known to result in poor control of asthma [15]. Sleep problems are common in patients with cirrhosis of the liver. They are encountered in almost 60% of patients with untreated hepatitis C. In patients on interferon- α therapy, the incidence improves to 30% [16]. Sleep problems are noted in almost 50% of patients with early CKD, and this percentage increases as high as 80% in patients with end-stage kidney disease (ESKD) [17]. Metabolic dysfunction has been noted in patients with insufficient sleep, insomnia, sleep apnea, narcolepsy, and many

other sleep disorders. This may lead to weight gain, obesity, and T2DM by food intake, disrupting energy balance, inflammation, impairing glucose tolerance, and insulin sensitivity. These abnormalities are not fully compensated by recovery sleep [18]. Several studies have established the causal relationship between short sleep duration and obesity. There is also a reverse causation as obesity leads to many co-morbidities (such as depression, gastrointestinal reflux disease, degenerative joint disease, asthma etc.) which deleteriously affect sleep [19]. Chen et al. reported that short sleep duration increased the risk of cancer in Asians while a long sleep duration increased the risk for colorectal cancer [20]. Sleep disorders have a mutual relationship with many skin disorders [21]. Several dermatosis result in pruritus, pain, burning sensations, pain, or inability to move, disturbing sleep. A bidirectional relationship with psoriasis and insomnia, obstructive sleep apnea syndrome, and restless leg syndrome has been noted. They have also been linked with infertility [22]. Sleep cycles normally parallel the reproductive cycle. Caetano et al. reviewed [33] studies published in the past 20 years and surmised that disturbed sleep results in an increased risk of menstrual irregularity, lower sperm count, and concentration, and a lower percentage of sperm with normal morphology when compared with normal sleep [22]. Mental health may also be affected 23-29. Sleep-related mental conditions include depression [22], anxiety [23], attention-deficit/hyperactivity disorder [24], Asperger syndrome [25,26], schizophrenia [27], and borderline personality/antisocial personality [28,29]. The relationship is usually bidirectional - most of these chronic non-communicable physical and mental diseases also interfere with normal sleep [30-33]. Poor sleepers face higher premature mortality [34]. Sleep duration-mortality risk has been studied in several meta-analytic studies – these show a 10% to 12% increased risk of mortality for short sleep duration and a 30% to 38% increased risk of mortality for long sleep duration [35]. Sleep is usually assessed by several techniques, including polysomnography, electrophysiological recordings of brain activity, muscle activity, and eye movements [36]. Sleep disorders impart a high economic burden on society [37].

2. DISCUSSION

Five lifestyles have been recognized as having a major impact on morbidity and mortality [38]. The

healthy behaviors are non-smoking, abstinence or low to moderate alcohol intake, a normal body mass index, regular exercise, and a prudent quality of diet. According to Li et al.[39], adherence to these lifestyles significantly improves survival.

3. SMOKING

Cigarette smoking is a major preventable cause of non-communicable diseases in the world [40]. Smoking is bi-directionally associated with sleep disorders. Smokers are more likely to develop poor sleep habits while poor sleepers are more likely to start smoking [41]. Cohrs et al. found that difficulty falling asleep and short sleep duration was related to smoking, and this increased as the number of pack-years smoked increased [42]. Smoking-related suboptimal sleep has also been confirmed in studies in the US, Germany, and Australia [43-45]. Smoking is independently associated with OSA [46]. Sleep apnea is a common medical condition that leads to disturbed sleep. Kashyap et al. found that current smoking was 35% in OSA patients compared to 18% in non-OSA patients [47]. Several subsequent studies have confirmed a higher prevalence of smoking in OSA patients [48,49]. Using polysomnography, Jaehne et al. found adult smokers, compared to nonsmokers, had more sleep apnea [50]. Ioannidou et al. reported that using univariate analysis, the prevalence of OSA was 1.2 times higher in current smokers compared with never and former smokers combined [51]. They also found that the prevalence in former smokers was 1.49 times higher when compared with never smokers [51]. Heavy smokers are also diagnosed at an earlier age with OSA [52]. They exhibit increased apnea-hypopnea index (commonly referred to as AHI) and reduced oxygen saturation, to a higher degree than non-smokers with OSA [53]. Smoking disturbs the sleep architecture via several mechanisms. Nicotine disturbs sleep by influencing sleep-regulating neurotransmitters [54]. In OSA patients, smoking causes several harmful changes. These include chronic mucosal inflammation and damaged cilia function in the nose, thickening, and swelling of the uvular mucosa, impairment of the neuromuscular protective reflexes of the upper airway, and an accelerated loss of lung function [55-57]. Although smoking is a risk factor and worsens OSA, untreated OSA is also related to smoking addiction [58]. Overall, smoking disrupts both the quantity and quality of sleep,

leading to a reduction in the quality of life in these individuals [59].

4. ALCOHOL

Alcohol is normally a relaxant and may even have euphoric effects [60]. It is commonly consumed during social occasions all over the world [61]. Because of its central nervous system depressant actions, it is often used as a non-prescription sleep-aid [62]. Evidence shows that a single alcoholic drink before sleep (in the evening) improves sleep patterns during the first half of sleep by reducing sleep onset latency and increasing non-rapid eye movement (NREM) sleep [63]. It however disrupts sleep during the second half. These effects may be related to the effect of alcohol on gamma-aminobutyric acid (GABA) and dopamine [64]. Alcohol, if taken in excess, has more detrimental effects on sleep [65]. Alcohol and sleep disturbances are usually bi-directionally related [66]. A prospective longitudinal study, involving 8,960 Finnish adults found that heavy drinking resulted in 1.5 higher odds of having problems with sleep [66]. In the same study, poor sleepers were 1.3 more likely to be heavy drinkers [66]. Alcohol intake has been implicated in sleep disturbances in children, adolescents, and adults [65,67-69]. Sleep disruptions are common in binge drinkers and chronic alcohol abusers - and there appears to be a dose-response relationship [70]. These individuals usually demonstrate an increase in sleep onset latency and a reduced sleep duration. This is associated with an increased tendency to sleep during the normal 'awake' day period. Alcohol-dependent patients have more sleep problems than healthy individuals, and this, in turn, can facilitate the development of alcoholism [71]. Alcoholics admitted for treatment have a higher rate of insomnia compared to healthy individuals [72]. In addition, alcoholism has a positive association with OSA [73]. Finally, the higher the sleep disturbance in alcoholics, the higher the relapse in those trying to quit [74].

5. OBESITY

Body mass index (BMI) is widely used as a measure to define body weight [75]. BMI is commonly measured by this equation: $BMI = \frac{\text{weight (lb)}}{\text{height (in)}^2} \times 703$. In adults (age over 18 years) BMI is categorized into several groups: < 19.9 (underweight), 20.0–24.9. (normal weight), and 25 to 29.9 (overweight). Obesity is defined as a BMI exceeding 30 [76] and is subclassified into class 1 (30–34.9), class 2 (35–

39.9), and class 3 or severe obesity (≥ 40). A BMI >50 is considered morbid obesity. The International Obesity Task Force recommends different BMI categories for Asians, due to their physical structure and increased percentage of abnormal fat accumulation. These are as follows: underweight (<18.5), normal (between 18.5 and 23), overweight (between 23 and 25), obese (between 25 and 30), severe obese (≥ 30). BMI is not used for children and adolescents aged 2 to 18 years; instead, a percentile scale based on the child's sex and age is used. Central (or visceral) obesity, as determined by a waist circumference >102 cm in males and >88 cm in females, a waist to hip ratio of more than 0.85 in women and >0.9 in men, and a waist-height ratio >0.5 , appears to be more harmful than obesity defined by BMI [77]. Obesity is strongly associated with several chronic diseases, including CVD, hypertension, T2DM, nonalcoholic fatty liver disease (NAFLD), gallbladder disease, pancreatitis, sleep apnea, and osteoarthritis [78]. It also increases the risk of several cancers [79]. Obesity also has a strong association with sleep disorders [80-90]. People who are obese have twice as many subjective sleep problems as those that are non-obese [80,81]. Being obese or overweight is associated with a decreased amount of sleep compared to those with normal body weight [82]. Obese individuals often complain of daytime sleepiness and fatigue, even if they sleep undisturbed during the night [83]. They may also suffer from ailments such as depression [84], gastroesophageal reflux disease (GERD) [85], osteoarthritis [86], and asthma [87], all of which can further disrupt sleep. Obesity is also associated with OSA, a major cause of sleep fragmentation [88]. Weight loss has a positive effect on sleep quality and duration [89,90]. Sleep disturbances also increase the risk of obesity [91]. This may result from increased food intake or physical inactivity, both often causing poor sleep [92,93]. The relationship is therefore bidirectional.

6. EXERCISE

One metabolic equivalent (MET) equals an energy expenditure of 1 kcal/kg/hour or an oxygen uptake of 3.5 ml/kg/min [94]. Sedentary behavior involves an energy expenditure <1.5 METs [95], while exercise expends more than 1.5 METs - light-intensity activities expend less than 3 METs, moderate-intensity activities expend 3 to 6 METs, and vigorous activities expend 6 or more METs [96]. Exercise may be

an effective non-pharmacological intervention to improve sleep. Several studies have validated this beneficial effect [97,98], irrespective of the form of exercise used [99]. Acute and chronic aerobic exercise improves sleep in midlife and later life [100]. A systemic review of thirteen studies suggested that resistance exercise also helps improve sleep quality [101]. In a study of older adults, both aerobic or resistance exercises significantly reduce sleep onset latency and medication use [102]. Even non-traditional exercises like Tai chi and yoga help improve sleep [103,104]. As often noted with other lifestyle factors, exercise also helps improve several chronic co-morbid conditions such as CVD [105], DM [106], obesity [107], OSA [108], and depression [109] – all of these are known to impair restful sleep.

7. DIET

Diet plays an important role in maintaining good health. Controlling portions and restricting calories are important dietary interventions that help prevent overweight/obesity [110]. In some studies, normal-weight adults exhibit higher sleep efficiency than overweight or obese individuals [111]. Weight loss thru dietary modification (combined with exercise) helps improve sleep [112]. On the other hand, lack of sleep has been shown to increase snacking, the number of meals consumed per day, and the preference for energy-rich foods, increasing the tendency to gain excess body weight [113-117]. It has been estimated that for every 1 hour later start of eating, there is a 1.25 unit increase in percent body fat [118]. Further, eating late may also increase the risk of gastroesophageal reflux disease (GERD), which in turn may promote sleep fragmentation [119]. The relationship between macronutrient intake and healthy sleep is complex. A high fat intake, especially from saturated fats, leads to less restorative sleep [120]. The 2007–2008 National Health and Nutrition Examination Survey results from 4,552 individuals, showed that a cholesterol-rich diet is associated with non-restorative sleep [121]. Intake of polyunsaturated fatty acids, docosahexaenoic acid (DHA), and arachidonic acid, on the other hand, appears to enhance sleep initiation and sleep duration [122,123]. In a study of 395 healthy children from schools in the United Kingdom, supplementation with DHA over 16-weeks led to an improvement in sleep quality [122]. Catalá reported that DHA may play a role in converting serotonin into melatonin, thereby, promoting sleep [123]. Alterations in both

quantity and quality of carbohydrate intake can also influence sleep [124]. High-carbohydrate meals with high glycemic indexes (especially high energy drinks and sugar-sweetened beverages) lead to poorer sleep quality [124]. Carbohydrates with a high glycemic index cause compensatory hyperinsulinemia, which along with counter-regulatory hormone responses results in disturbed sleep [125]. On the other hand, carbohydrates with high fiber, such as whole grains, fruits, and vegetables, are sleep friendly. As far as protein is concerned, a high protein intake diet helps sleep, especially in overweight and obese individuals [126]. Tryptophan, an essential amino acid, is needed to produce serotonin, which in turn creates melatonin [127]. Melatonin has a strong hypnotic and chronobiotic property [128] and helps sleep. Sleep quality can therefore be improved by eating foods rich in tryptophan or by taking a melatonin supplement. Deficiency of vitamins A, B, C, D, E, and K [129,130], and of minerals such as iron [131], zinc [132], calcium [133], and magnesium [134] is also associated with disturbed sleep. Some dietary supplements have been noted to improve sleep. These include Phlorotannins [135], GABA, Apocynum venetum leaf extract [136], Chlorophytum bovivilianum root, and velvet bean [137]. However, further studies are needed to confirm the therapeutic practicality of using these. Finally, stimulants like tea and coffee [138] and spicy foods [139] may also interfere with a night of restorative sleep. Overall, a balanced diet that is rich in non-starchy vegetables, fruits, whole grains, and legumes, moderate in consumption of nuts, seafood, lean meats, low-fat dairy products, and vegetable oil, and limited or void in trans-fats, saturated fats, sodium, red meat, refined carbohydrates, and sugar-sweetened beverages is associated with good sleep [140]. A prudent diet is also important in mitigating several chronic diseases, such as cardiovascular and respiratory disorders, several cancers, and T2DM [141]. Diets such as the Mediterranean [142] and the dietary approaches to stop hypertension (DASH) [143], are primarily plant-based and rich in healthy ingredients and help improve sleep duration and quality.

8. CONCLUSION

Proper sleep is vital for a healthy life. It is necessary for bodily repair, consolidation of memories, and good mental health. Proper sleep is characterized by falling asleep easily and staying asleep (uninterrupted) for 7-9 hours. It is

also associated with feeling refreshed after waking up and not feeling sleepy during the day. Several diseases, such as those involving the cardiovascular, respiratory, metabolic, and neurological systems, can alter sleep quantity and quality. Inadequate or fragmented sleep can in turn detrimentally affect these ailments. As this minireview shows, sleep can be adversely affected by several unhealthy lifestyles. These include smoking, excessive alcohol intake, excess body weight, sedentary behavior, and poor nutrition. Healthy lifestyles are therefore essential for regular salubrious sleep.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Benington JH, Craig Heller H. Restoration of brain energy metabolism as the function of sleep. *Prog. Neurobiol.* 1995;45:347–360.
DOI: 10.1016/0301-0082(94)00057-O
2. Gómez-González B, Domínguez-Salazar E, Hurtado-Alvarado G, et al. Role of sleep in the regulation of the immune system and the pituitary hormones. *Ann. N. Y. Acad. Sci.* 2012;1261:97–106.
DOI: 10.1111/j.1749-6632.2012.06616.x
3. Schönauer M. Sleep spindles: Timed for memory consolidation. *Curr. Biol.* 2018;28:R656–R658.
DOI: 10.1016/j.cub.2018.03.046
4. Ohayon M, Wickwire EM, Hirshkowitz M, et al. National Sleep Foundation's sleep quality recommendations: First report. *Sleep Health.* 2017;3:6–19.
DOI: 10.1016/j.sleh.2016.11.006.
5. Watson NF, Badr MS, Belenky G, et al. Joint consensus statement of the American academy of sleep medicine and sleep research society on the recommended amount of sleep for a healthy adult: Methodology and discussion. *J. Clin. Sleep Med.* 2015;11:931–952.
DOI: 10.5665/sleep.4716

6. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's sleep time duration recommendations: Methodology and results summary. *Sleep health*. 2015;1:40–43.
DOI: 10.1016/j.sleh.2014.12.010.
7. Available: https://www.cdc.gov/sleep/about_sleep/how_much_sleep.html
8. Available: <https://order.nia.nih.gov/sites/default/files/2020-12/sleep-and-older-adults-booklet.pdf>.
9. Sateia MJ. International classification of sleep disorders-third edition: Highlights and modifications. *Chest*. 2014;146(5):1387-1394.
DOI: 10.1378/chest.14-0970
10. Steptoe A, O'Donnell K, Marmot M, Wardle J. Positive affect, psychological well-being, and good sleep. *Journal of Psychosomatic Research*. 2008;64(4):409–15.
DOI: 10.1016/j.jpsychores.2007.11.008.
11. Rosekind MR, Gregory KB, Mallis MM, Brandt SL, Seal B, Lerner D. The cost of poor sleep: workplace productivity loss and associated costs. *J Occup Environ Med*. 2010;52:91–98.
DOI: 10.1097/JOM.0b013e3181c78c30
12. Simon GE, Vonkorff M. Prevalence, burden, and treatment of insomnia in primary care. *American Journal of Psychiatry*. 1997;154(10):1417-1423.
13. Grandner MA, Brown D, Conroy MB, Jean-Louis G, Coons M, Bhatt DL. Sleep duration and quality: Impact on lifestyle behaviors and cardiometabolic health: A scientific statement from the American Heart Association. *Circulation*. 2016;134:e367–e386.
DOI: 10.1161/CIR.0000000000000444.
14. Ogilvie RP, Patel SR. The Epidemiology of Sleep and Diabetes. *Curr Diab Rep*. 2018;18(10):82.
DOI: 10.1007/s11892-018-1055-8.
15. Kavanagh J, Jackson DJ, Kent BD. Sleep and asthma. *Curr Opin Pulm Med*. 2018;24(6):569-573.
DOI: 10.1097/MCP.0000000000000526.
16. Sockalingam S, Abbey SE, Alosaimi F, Novak M. A review of sleep disturbance in hepatitis C. *J Clin Gastroenterol*. 2010;44(1):38-45.
DOI: 10.1097/MCG.0b013e3181b314ea.
17. Natale P, Ruospo M, Saglimbene VM, Palmer SC, Strippoli GF. Interventions for improving sleep quality in people with chronic kidney disease. *Cochrane Database Syst Rev*. 2019;5(5):CD012625.
DOI: 10.1002/14651858.CD012625.pub2.
18. Depner CM, Melanson EL, Eckel RH, et al. Ad libitum Weekend Recovery Sleep Fails to Prevent Metabolic Dysregulation during a Repeating Pattern of Insufficient Sleep and Weekend Recovery Sleep. *Curr Biol*. 2019;29(6):957-967.e4.
DOI: 10.1016/j.cub.2019.01.069
19. Ogilvie RP, Patel SR. The epidemiology of sleep and obesity. *Sleep Health*. 2017;3(5):383-388.
DOI: 10.1016/j.sleh.2017.07.013
20. Chen Y, Tan F, Wei L, et al. Sleep duration and the risk of cancer: A systematic review and meta-analysis including dose-response relationship. *BMC Cancer*. 2018;18(1):1149.
DOI: 10.1186/s12885-018-5025-y
21. Nowowiejska J, Baran A, Flisiak I. Mutual Relationship Between Sleep Disorders, Quality of Life and Psychosocial Aspects in Patients With Psoriasis. *Front Psychiatry*. 2021;12:674460.
DOI: 10.3389/fpsy.2021.674460
22. Caetano G, Bozinovic I, Dupont C, Léger D, Lévy R, Sermondade N. Impact of sleep on female and male reproductive functions: A systematic review. *Fertil Steril*. 2021;115:715-731.
23. Benca RM, Obermeyer WH, Thisted RA, Gillin JC. Sleep and psychiatric disorders. A meta-analysis. *Arch Gen Psychiatry*. 1992;49(8):651-68; discussion 669-70.
DOI: 10.1001/archpsyc.1992.01820080059010
24. Willis TA, Gregory AM. Anxiety Disorders and Sleep in Children and Adolescents. *Sleep Med Clin*. 2015;10(2):125-31.
DOI: 10.1016/j.jsmc.2015.02.002
25. Becker SP. ADHD and sleep: Recent advances and future directions. *Curr Opin Psychol*. 2020;34:50-56.
DOI: 10.1016/j.copsyc.2019.09.006
26. Paavonen EJ, Vehkalahti K, Vanhala R, von Wendt L, Nieminen-von Wendt T, Aronen ET. Sleep in children with Asperger syndrome. *J Autism Dev Disord*. 2008;38(1):41-51.
DOI: 10.1007/s10803-007-0360-x
27. Johnson KP, Malow BA. Sleep in children with autism spectrum disorders. *Curr Neurol Neurosci Rep*. 2008;8(2):155-61.
DOI: 10.1007/s11910-008-0025-y
28. Ferrarelli F. Sleep disturbances in schizophrenia and psychosis. *Schizophr Res*. 2020;221:1-3.
DOI: 10.1016/j.schres.2020.05.022.

29. Hafizi S. Sleep and borderline personality disorder: A review. *Asian J Psychiatr.* 2013;6(6):452-9.
DOI: 10.1016/j.ajp.2013.06.016
30. Redeker NS. Sleep disturbance in people with heart failure: implications for self-care. *J Cardiovasc Nurs.* 2008;23(3):231-8.
DOI: 10.1097/01.JCN.0000305094.20012.76
31. Kavanagh J, Jackson DJ, Kent BD. Sleep and asthma. *Curr Opin Pulm Med.* 2018;24(6):569-573.
DOI: 10.1097/MCP.0000000000000526
32. Benca RM, Peterson MJ. Insomnia and depression. *Sleep Med.* 2008;9(1):S3-9.
DOI: 10.1016/S1389-9457(08)70010-8
33. Available:<https://www.sleepfoundation.org/>. Accessed January 26, 2022
34. Åkerstedt T, Ghilotti F, Grotta A, Zhao H, Adami HO, Trolle-Lagerros Y, Bellocco R. Sleep duration and mortality - Does weekend sleep matter? *J Sleep Res.* 2019;28(1):e12712.
DOI: 10.1111/jsr.12712
35. Mullington JM, Simpson NS, Meier-Ewert HK, Haack M. Sleep loss and inflammation. *Best Pract Res Clin Endocrinol Metab.* 2010;24(5):775-784.
DOI:10.1016/j.beem.2010.08.014
36. Baglioni C, Nanovska S, Regen W, et al. Sleep and mental disorders: A meta-analysis of polysomnographic research. *Psychol Bull.* 2016;142(9):969-990.
DOI:10.1037/bul0000053
37. Institute of Medicine (US) Committee on Sleep Medicine and Research; Colten HR, Altevogt BM, editors. *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem.* Washington (DC): National Academies Press (US); 2006. 4, Functional and Economic Impact of Sleep Loss and Sleep-Related Disorders. Available:<https://www.ncbi.nlm.nih.gov/books/NBK19958/>
38. Loef M, Walach H. The combined effects of healthy lifestyle behaviors on all cause mortality: A systematic review and meta-analysis. *Prev Med.* 2012;55:163-170.
39. Li Y, Pan A, Wang DD, Liu X, Dhana K, Franco OH, Kaptoge S, Di Angelantonio E, Stampfer M, Willett WC, Hu FB. Impact of Healthy Lifestyle Factors on Life Expectancies in the US Population. *Circulation.* 2018 Jul 24;138(4):345-355.
DOI:10.1161/CIRCULATIONAHA.117.032047
40. White WB. Smoking-related morbidity and mortality in the cardiovascular setting. *Prev Cardiol.* 2007;10(2 Suppl 1):1-4.
DOI: 10.1111/j.1520-037x.2007.06050.x
41. Bellatorre A, Choi K, Lewin D, Haynie D, Simons-Morton B. Relationships Between Smoking and Sleep Problems in Black and White Adolescents. *Sleep.* 2016;40
DOI: 10.1093/sleep/zsw031
42. Cohrs S, Rodenbeck A, Riemann D, Szagun B, Jaehne A, Brinkmeyer J, et al. Impaired sleep quality and sleep duration in smokers-results from the German Multicenter Study on Nicotine Dependence. *Addict Biol.* 2014;19(3):486-96.
Available:<https://doi.org/10.1111/j.1369-1600.2012.00487.x>
43. Bellatorre A, Choi K, Lewin D, Haynie D, Simons-Morton B. Relationships Between Smoking and Sleep Problems in Black and White Adolescents. *Sleep.* 2016;40
DOI: 10.1093/sleep/zsw031
44. Jaehne A, Unbehau T, Feige B, Lutz UC, Batra A, Riemann D. How smoking affects sleep: A polysomnographical analysis. *Sleep Med.* 2012;13:1286-1292.
DOI: 10.1016/j.sleep.2012.06.026
45. Metse AP, Clinton-McHarg T, Skinner E, Yogaraj Y, Colyvas K, Bowman J. Associations between Suboptimal Sleep and Smoking, Poor Nutrition, Harmful Alcohol Consumption and Inadequate Physical Activity ('SNAP Risks'): A Comparison of People with and without a Mental Health Condition in an Australian Community Survey. *Int J Environ Res Public Health.* 2021;18(11):5946.
DOI: 10.3390/ijerph18115946
46. Rundo JV. Obstructive sleep apnea basics. *Cleve Clin J Med.* 2019;86(9 Suppl 1):2-9.
DOI: 10.3949/ccjm.86.s1.02
47. Kashyap R, Hock LM, Bowman TJ. Higher prevalence of smoking in patients diagnosed as having obstructive sleep apnea. *Sleep Breath.* 2001;5:167-172.
DOI: 10.1055/s-2001-18805.8
48. Deleanu OC, Pocora D, Mihălcuță S, Ulmeanu R, Zaharie AM, Mihălțan FD. Influence of smoking on sleep and obstructive sleep apnea syndrome. *Pneumologia.* 2016;65:28-35.
49. Bielicki P, Trojnar A, Sobieraj P, Wąsik M. Smoking status in relation to obstructive sleep apnea severity (OSA) and cardiovascular comorbidity in patients with

- newly diagnosed OSA. *Adv. Respir. Med.* 2019;87:103–109.
DOI: 10.5603/ARM.a2019.0011
50. Jaehne A, Unbehau T, Feige B, Lutz UC, Batra A, Riemann D. How smoking affects sleep: A polysomnographical analysis. *Sleep Med.* 2012;13:1286–1292.
DOI: 10.1016/j.sleep.2012.06.026
 51. Ioannidou D, Kalamaras G, Kotoulas SC, Pataka A. Smoking and Obstructive Sleep Apnea: Is There An Association between These Cardiometabolic Risk Factors?-Gender Analysis. *Medicina (Kaunas).* 2021;57(11):1137.
DOI: 10.3390/medicina57111137
 52. Varol Y, Anar C, Tuzel OE, Guclu SZ, Ucar ZZ. The impact of active and former smoking on the severity of obstructive sleep apnea. *Sleep Breath.* 2015;19:1279–1284.
DOI: 10.1007/s11325-015-1159-1
 53. Yosunkaya S, Kutlu R, Vatansev H. Effects of smoking on patients with obstructive sleep apnea syndrome. *Clin. Respir. J.* 2021;15:147–153.
DOI: 10.1111/crj.13278
 54. Jaehne A, Unbehau T, Feige B, Cohrs S, Rodenbeck A, Schütz AL, Uhl V, Zober A, Riemann D. Sleep changes in smokers before, during and 3 months after nicotine withdrawal. *Addict Biol.* 2015;20(4):747-55.
DOI: 10.1111/adb.12151
 55. Yaniv E, Shvili Y, Koren R, Shvero J. Histopathological changes of the nasal mucosa induced by smoking. *Inhal. Toxicol.* 2009;21:1119–1122.
DOI: 10.3109/08958370902767070
 56. Dicipinigaitis PV. Cough reflex sensitivity in cigarette smokers. *Chest.* 2003;123:685–688.
DOI: 10.1378/chest.123.3.685
 57. Knoell DL, Smith D, Bao S, Sapkota M, Wyatt TA, Zweier JL, Flury J, Borchers MT, Knutson M. Imbalance in zinc homeostasis enhances lung Tissue Loss following cigarette smoke exposure. *J Trace Elem Med Biol.* 2020;60:126483.
DOI: 10.1016/j.jtemb.2020.126483
 58. Schrand JR. Is sleep apnea a predisposing factor for tobacco use? *Med. Hypotheses.* 1996;47:443–448.
DOI: 10.1016/S0306-9877(96)90155-3
 59. Hattatoğlu DG, Aydın Ş, Yıldız BP. Does smoking impair sleep hygiene? *Arq Neuropsiquiatr.* 2021;S0004-282X2021005024104.
DOI: 10.1590/0004-282X-ANP-2020-0578
 60. Freed EX. Alcohol and mood: an updated review. *Int J Addict.* 1978;13(2):173-200.
DOI: 10.3109/10826087809039273
 61. Sudhinaraset M, Wigglesworth C, Takeuchi DT. Social and Cultural Contexts of Alcohol Use: Influences in a Social-Ecological Framework. *Alcohol Res.* 2016;38(1):35-45.
Available:www.Sleepfoundation.org.
 63. Thakkar MM, Sharma R, Sahota P. Alcohol disrupts sleep homeostasis. *Alcohol.* 2015;49(4):299-310.
DOI: 10.1016/j.alcohol.2014.07.019.
 64. Koob GF, Colrain IM. Alcohol use disorder and sleep disturbances: a feed-forward allostatic framework. *Neuropsychopharmacology.* 2020;45(1):141-165.
DOI: 10.1038/s41386-019-0446-0
 65. Canham SL, Kaufmann CN, Mauro PM et al. Binge drinking and insomnia in middle-aged and older adults: the Health and Retirement Study. *Int J Geriatr Psychiatry* 2015;30:284–291.
 66. Haario P, Rahkonen O, Laaksonen M, Lahelma E., Lallukka TEA. Bidirectional associations between insomnia symptoms and unhealthy behaviours. *J. Sleep Res.* 2013;22:89–95.
DOI: 10.1111/j.1365-2869.2012.01043.x
 67. Zhabenko O, Austic E, Conroy DA et al. Substance use as a risk factor for sleep problems among adolescents presenting to the emergency department. *Journal of Addiction Medicine* 2016;10:331–338.
 68. Wong MM, Brower KJ, Nigg JT, Zucker RA. Childhood sleep problems, response inhibition, and alcohol and drug outcomes in adolescence and young adulthood. *Alcoholism, clinical and experimental research* 2010;34:1033–1044.
 69. Wang YM, Chen HG, Song M et al. Prevalence of insomnia and its risk factors in older individuals: A community-based study in four cities of Hebei Province, China. *Sleep Medicine* 2016;19:116–122.
 70. Popovici I, French MT. Binge drinking and sleep problems among young adults. *Drug and Alcohol Dependence.* 2013;132:207–215.
 71. Gann H, van Calker D, Feige B, Riemann D. Die Bedeutung des Schlafs für gesunde Alkoholkonsumenten und alkoholabhängige Patienten [The importance of sleep for healthy alcohol consumers and alcohol dependent

- patients]. *Nervenarzt*. 2004;75(5):431-41. German.
DOI: 10.1007/s00115-003-1637-8
72. Brower KJ, Hall JM. Effects of age and alcoholism on sleep: A controlled study. *Journal of Studies on Alcohol*. 2001;62:335–343.
 73. Taveira KVM, Kuntze MM, Berretta F, de Souza BDM, Godolfim LR, Demathe T, De Luca Canto G, Porporatti AL. Association between obstructive sleep apnea and alcohol, caffeine and tobacco: A meta-analysis. *J Oral Rehabil*. 2018;45(11):890-902.
DOI: 10.1111/joor.12686
 74. Chakravorty S, Chaudhary NS, Brower KJ. Alcohol Dependence and Its Relationship With Insomnia and Other Sleep Disorders. *Alcohol Clin Exp Res*; 2016.
 75. Available: https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmicalc.htm - accessed October 27;2021.
 76. Hirsch J, Salans LB, Aronne LJ. Obesity. Becker, KL eds. *Principles and Practice of Endocrinology and Metabolism*. 3rd ed. Lippincott, Williams, and Wilkins Philadelphia. 2001; 1239–1246.
 77. Ross R, Neeland IJ, Yamashita S, et al. Waist circumference as a vital sign in clinical practice: a Consensus Statement from the IAS and ICCR Working Group on Visceral Obesity. *Nat Rev Endocrinol*. 2020;16(3):177-189.
DOI:10.1038/s41574-019-0310-7
 78. Schelbert KB. Comorbidities of Obesity. *Prim Care*. 2009;36(2):271–85.
10.1016/j.pop.2009.01.009.
 79. Lauby-Secretan B, Scoccianti C, Loomis D, et al. Body fatness and cancer — viewpoint of the IARC Working Group. *N Engl J Med*. 2016;375:794-798.
 80. Vgontzas AN, Lin HM, Papaliaga M, Calhoun S, Vela-Bueno A, et al. Short sleep duration and obesity: The role of emotional stress and sleep disturbances. *Int J Obes (Lond)*. 2008;32(5):801–809.
 81. Ko GT, Chan JC, Chan AW, Wong PT, Hui SS, et al. Association between sleeping hours, working hours and obesity in Hong Kong Chinese: the ‘better health for better Hong Kong’ health promotion campaign. *Int J Obes (Lond)*. 2007;31(2):254–260.
 82. Vorona RD, Winn MP, Babineau TW, Eng BP, Feldman HR, et al. Overweight and obese patients in a primary care population report less sleep than patients with a normal body mass index. *Arch Intern Med*. 2005;165(1):25–30.
 83. Vgontzas AN, Bixler EO, Tan TL, Kantner D, Martin LF, Kales A. Obesity without sleep apnea is associated with daytime sleepiness. *Archives of internal medicine*. 1998;158(12):1333–1337.
Available: <https://doi.org/10.1001/archinte.158.12.1333>
 84. Faith MS, Butryn M, Wadden TA, Fabricatore A, Nguyen AM, Heymsfield SB. Evidence for prospective associations among depression and obesity in population-based studies. *Obesity reviews: An official journal of the International Association for the Study of Obesity*. 2011;12(5):e438–453.
 85. Chang P, Friedenberg F. Obesity and GERD. *Gastroenterol Clin North Am*. 2014;43(1):161–173.
 86. Lementowski PW, Zelicof SB. Obesity and osteoarthritis. *Am J Orthop (Belle Mead NJ)* 2008;37(3):148–151.
 87. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: A meta-analysis of prospective epidemiologic studies. *American journal of respiratory and critical care medicine*. 2007;175(7):661–666.
 88. Romero-Corral A, Caples SM, Lopez-Jimenez F, Somers VK. Interactions between obesity and obstructive sleep apnea: implications for treatment. *Chest*. 2010;137(3):711–719.
DOI: 10.1378/chest.09-0360
 89. Gangwisch JE, Malaspina D, Boden-Albala B, Heymsfield SB. Inadequate sleep as a risk factor for obesity: analyses of the NHANES I. *Sleep*. 2005;28(10):1289.
 90. Kohatsu ND, Tsai R, Young T, Vangilder R, Burmeister LF. Sleep duration and body mass index in a rural population. *Arch Intern Med*. 2006;166(16):1701–1705.
 91. Ogilvie RP, Patel SR. The epidemiology of sleep and obesity. *Sleep Health*. 2017 Oct;3(5):383-388.
DOI: 10.1016/j.sleh.2017.07.013
 92. Knutson KL, Spiegel K, Penev P, Van Cauter E. The metabolic consequences of sleep deprivation. *Sleep Med Rev*. 2007;11(3):163–178.
 93. Li L, Zhang S, Huang Y, Chen K. Sleep duration and obesity in children: A systematic review and meta-analysis of prospective cohort studies. *J Paediatr Child Health*. 2017 Apr;53(4):378-385.
DOI: 10.1111/jpc.13434

94. Jetté M, Sidney K, Blümchen G. Metabolic equivalents (METs) in exercise testing, exercise prescription, and evaluation of functional capacity. *Clin Cardiol.* 1990;13(8):555-65.
DOI: 10.1002/clc.4960130809
95. Pinto BM, Kindred MD, Dunsiger SI, Williams DM. Sedentary behavior among breast cancer survivors: A longitudinal study using ecological momentary assessments. *J Cancer Surviv.* 2021;15(4):546-553.
DOI: 10.1007/s11764-020-00948-x
96. Available: https://www.cdc.gov/nccdphp/dnpa/physical/pdf/PA_Intensity_table_2_1.pdf
97. Passos GS, Poyares D, Santana MG, D'Aurea CV, Youngstedt SD, Tufik S, de Mello MT. Effects of moderate aerobic exercise training on chronic primary insomnia. *Sleep Med.* 2011;12(10):1018-27.
DOI: 10.1016/j.sleep.2011.02.007
98. Abd El-Kader SM, Al-Jiffri OH. Aerobic exercise affects sleep, psychological wellbeing and immune system parameters among subjects with chronic primary insomnia. *Afr Health Sci.* 2020;20(4):1761-1769.
DOI: 10.4314/ahs.v20i4.29
99. Bonardi JMT, Lima LG, Campos GO, Bertani RF, Moriguti JC, Ferriolli E, Lima NKC. Effect of different types of exercise on sleep quality of elderly subjects. *Sleep Med.* 2016;25:122-129.
DOI: 10.1016/j.sleep.2016.06.025
100. King AC, Oman RF, Brassington GS, Bliwise DL, Haskell WL. Moderate-intensity exercise and self-rated quality of sleep in older adults: a randomized controlled trial. *JAMA.* 1997;277(1):32-37.
DOI: 10.1001/jama.1997.03540250040029
101. Kovacevic A, Mavros Y, Heisz JJ, Fiatarone Singh MA. The effect of resistance exercise on sleep: A systematic review of randomized controlled trials. *Sleep Med Rev.* 2018;39:52-68.
DOI: 10.1016/j.smrv.2017.07.002
102. Yang PY, Ho KH, Chen HC, Chien MY. Exercise training improves sleep quality in middle-aged and older adults with sleep problems: A systematic review. *J Physiother.* 2012;58(3):157-63.
DOI: 10.1016/S1836-9553(12)70106-6.
103. Si Y, Wang C, Yin H, Zheng J, Guo Y, Xu G, Ma Y. Tai Chi Chuan for Subjective Sleep Quality: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Evid Based Complement Alternat Med.* 2020;2020:4710527.
DOI: 10.1155/2020/4710527
104. Wang WL, Chen KH, Pan YC, Yang SN, Chan YY. The effect of yoga on sleep quality and insomnia in women with sleep problems: a systematic review and meta-analysis. *BMC Psychiatry.* 2020;20(1):195.
DOI: 10.1186/s12888-020-02566-4.
105. Fiuza-Luces C, Santos-Lozano A, Joyner M, Carrera-Bastos P, Picazo O, Zugaza JL, Izquierdo M, Ruilope LM, Lucia A. Exercise benefits in cardiovascular disease: beyond attenuation of traditional risk factors. *Nat Rev Cardiol.* 2018;15(12):731-743.
DOI: 10.1038/s41569-018-0065-1
106. Lumb A. Diabetes and exercise. *Clin Med (Lond).* 2014;14(6):673-6.
DOI: 10.7861/clinmedicine.14-6-673
107. Celik O, Yildiz BO. Obesity and physical exercise. *Minerva Endocrinol (Torino).* 2021;46(2):131-144.
DOI: 10.23736/S2724-6507.20.03361-1
108. Andrade FM, Pedrosa RP. The role of physical exercise in obstructive sleep apnea. *J Bras Pneumol.* 2016;42(6):457-464.
DOI: 10.1590/S1806-37562016000000156
109. Schuch FB, Stubbs B. The Role of Exercise in Preventing and Treating Depression. *Curr Sports Med Rep.* 2019;18(8):299-304.
DOI: 10.1249/JSR.0000000000000620
110. Baker B. Weight loss and diet plans. *Am J Nurs.* 2006;106(6):52-9; quiz 60.
DOI: 10.1097/00000446-200606000-00027
111. Kahlhöfer J, Karschin J, Breusing N, Bösych Westphal A. Relationship between actigraphy-assessed sleep quality and fat mass in college students. *Obesity (Silver Spring).* 2016;24(2): 335-341.
112. Available: <https://www.health.harvard.edu/blog/losing-weight-and-belly-fat-improves-sleep-201211145531>.
113. St-Onge MP, Pizinger T, Kovtun K, RoyChoudhury A. Sleep and meal timing influence food intake and its hormonal regulation in healthy adults with overweight/obesity. *Eur J Clin Nutr.* 2019;72(Suppl 1):76-82.
DOI: 10.1038/s41430-018-0312-x
114. Thomas EA, Zaman A, Cornier MA, Catenacci VA, Tussey EJ, Grau L, Arbet J, Broussard JL, Rynders CA. Later Meal and Sleep Timing Predicts Higher

- Percent Body Fat. Nutrients. 2020;13(1):73.
DOI: 10.3390/nu13010073
115. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Sleep symptoms associated with intake of specific dietary nutrients. *Journal of sleep research*. 2014;23(1):22–34.
Available:https://doi.org/10.1111/jsr.12084
 116. Spiege K, Tasali E, Penev P, Van Cauter E. Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Annals of internal medicine*. 2004;141(11):846–850.
Available:https://doi.org/10.7326/0003-4819-141-11-200412070-00008
 117. Greer SM, Goldstein AN, Walker MP. The impact of sleep deprivation on food desire in the human brain. *Nature communications*. 2013;4:2259.
Available:https://doi.org/10.1038/ncomms3259.
 118. Available:https://read.qxmd.com/read/33383648/late-meal-and-sleep-timing-predicts-higher-percent-body-fat.
 119. Fujiwara Y, Machida A, Watanabe Y, Shiba M, Tominaga K, Watanabe T, Oshitani N, Higuchi K, Arakawa T. Association between dinner-to-bed time and gastro-esophageal reflux disease. *Am J Gastroenterol*. 2005;100(12):2633-6.
DOI: 10.1111/j.1572-0241.2005.00354.x
 120. St-Onge M, Roberts A, Shechter A, Choudhury AR. Fiber and saturated fat are associated with sleep arousals and slow wave sleep. *J Clin Sleep Med*. 2016;12(1):19–24.
 121. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Sleep symptoms associated with intake of specific dietary nutrients. *Journal of Sleep Research*. 2014;23:22–34.
DOI: 10.1111/jsr.12084
 122. Lavialle M, Champeil-Potokar G, Alessandri JM, et al. An (n-3) polyunsaturated fatty acid-deficient diet disturbs daily locomotor activity, melatonin rhythm, and striatal dopamine in Syrian hamsters. *J. Nutr*. 2008;138:1719–1724.
 123. Catalá A. The function of very long chain polyunsaturated fatty acids in the pineal gland. *Biochimica et Biophysica Acta*. 2010;1801:95–99.
 124. Gangwisch JE, Hale L, St-Onge MP, et al. High glycemic index and glycemic load diets as risk factors for insomnia: analyses from the Women's Health Initiative. *Am J Clin Nutr*. 2020;111(2):429-439.
DOI:10.1093/ajcn/nqz275
 125. Banarer S, Cryer PE. Sleep-related hypoglycemia-associated autonomic failure in type I diabetes: reduced awakening from sleep during hypoglycemia. *Diabetes*. 2003;52:1195–203.
 126. Zhou J, Kim JE, Armstrong CL, Chen N, Campbell WW. Higher-protein diets improve indexes of sleep in energy-restricted overweight and obese adults: results from 2 randomized controlled trials. *American Journal of Clinical Nutrition*. 2016;103(3):766.
DOI: 10.3945/ajcn.115.124669
 127. Naseem M, Parvez S. Role of melatonin in traumatic brain injury and spinal cord injury. *ScientificWorldJournal*. 2014;2014:586270.
DOI:10.1155/2014/586270
 128. Kostoglou-Athanassiou I. Therapeutic applications of melatonin. *Ther Adv Endocrinol Metab*. 2013;4(1):13-24.
DOI: 10.1177/2042018813476084
 129. Okawa M, Mishima K, Nanami T, Shimizu T, Iijima S, Hishikawa Y, Takahashi K. Vitamin B12 treatment for sleep-wake rhythm disorders. *Sleep*. 1990;13:15–23.
 130. Lichstein KL, Payne KL, Soeffing JP, Heith Durrence H, Taylor DJ, Riedel BW, Bush AJ. Vitamins and sleep: an exploratory study. *Sleep Med*. 2007;9:27–32.
 131. O'Keefe ST, Gavin K, Lavan JN. Iron status and restless legs syndrome in the elderly. *Age Ageing*. 1994;23:200–203.
 132. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Dietary nutrients associated with short and long sleep duration. Data from a nationally representative sample. *Appetite*. 2013;64:71–80.
DOI: 10.1016/j.appet.2013.01.004
 133. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Sleep symptoms associated with intake of specific dietary nutrients. *J Sleep Res*. 2014;23(1):22-34.
DOI:10.1111/jsr.12084
 134. Cao Y, Zhen S, Taylor AW, Appleton S, Atlantis E, Shi Z. Magnesium Intake and Sleep Disorder Symptoms: Findings from the Jiangsu Nutrition Study of Chinese Adults at Five-Year Follow-Up. *Nutrients*. 2018;10(10):1354.
DOI: 10.3390/nu10101354
 135. Cho S, Yoon M, Pae A, et al. Marine polyphenol phlorotannins promote non-

- rapid eye movement sleep in mice via the benzodiazepine site of the GABA A receptor. *Psychopharmacology*. 2014;231:2825–2837.
DOI: 10.1007/s00213-014-3445-1
136. Yamatsu A, Yamashita Y, Maru I, Yang J, Tatsuzaki J, Kim M. The Improvement of Sleep by Oral Intake of GABA and Apocynum venetum Leaf Extract. *J. Nutr. Sci. Vitaminol*. 2015;61:182–187.
DOI: 10.3177/jnsv.61.182
137. Alleman RJ, Canale RE, McCarthy CG, Bloomer RJ. Chlorophytum bovivilianum (root) and velvet bean a blend of chlorophytum borivilianum and velvet bean increases serum growth hormone in exercise-trained men. *Nutr. Metab. Insights*. 2011;4.
138. Wells AS, Read NW, Idzikowski C, Jones J. Effects of meals on objective and subjective measures of daytime sleepiness. *J Appl Physiol*. 1998. 84:507–515.
139. Edwards SJ, Montgomery IM, Colquhoun EQ, Jordan JE, Clark MG. Spicy meal disturbs sleep: An effect of thermoregulation? *Int J Psychophysiol*. 1992 Sep;13(2):97-100.
DOI: 10.1016/0167-8760(92)90048-g
140. U.S. Department of Health and Human Services and the U.S. Department of Agriculture Dietary Guidelines for Americans 2015-2020. 8th ed. Available: <https://health.gov/dietaryguidelines/2015/guidelines/>
141. World Health Organization. World Health Statistics 2018; 2018. Available: www.who.int/gho/publications/world_health_statistics/2018/en
142. Muscogiuri G, Barrea L, Aprano S, et al. for Opera Prevention Project. Sleep Quality in Obesity: Does Adherence to the Mediterranean Diet Matter?. *Nutrients*. 2020;12(5).
143. Liang H, Beydoun HA, Hossain S, Maldonado A, Zonderman AB, et al. Dietary Approaches to Stop Hypertension (DASH) Score and Its Association with Sleep Quality in a National Survey of Middle-Aged and Older Men and Women. *Nutrients*. 2020;12(5): 1510. Available: <https://doi.org/10.3390/nu12051510>

© 2021 Agarwal; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83544>