



Multifaceted Health Benefit of *Ficus capensis* Fruits and Vegetables

N. N. Umerah ^{a*}, P. N. Ani ^b and N. M. Oly-Alawuba ^c

^a Department of Food Science and Technology, Enugu State University of Science and Technology, P.M.B. 01660, Enugu, Nigeria.

^b Department of Nutrition and Dietetics, University of Nigeria, Nsukka, Nigeria.

^c Department of Nutrition and Dietetics, Imo State University, Owerri, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author NNU designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author PNA managed the analyses of the study. Author NMOA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AFSJ/2022/v21i930468

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/62391>

Original Research Article

Received 09 September 2020

Accepted 12 November 2020

Published 10 August 2022

ABSTRACT

This study was carried out to evaluate the chemical properties of *Ficus capensis* fruits and vegetables. *Ficus capensis* vegetables and fruits were separately plucked, sorted, cleaned and milled using electric blender until the desired particle size was obtained (150 - 850 microns). Analyses of the proximate, mineral, vitamins, phytochemicals and anti-nutrient contents of the vegetables and fruits were carried out using standard methods. Proximate analysis showed that *Ficus capensis* had a protein mean value of 6.11 and 8.02 g/100 g, ash 7.63 and 2.92 g/100 g, fibre 9.82 and 6.33 g/100 g, moisture 43.28 and 45.20 g/100 g, carbohydrate 31.48 and 35.61 g/100 g and fat content of 1.68 and 1.92 g/100 g for the vegetables and fruits respectively. Mineral constituents were iron 14.24 and 11.68 mg/100 g, calcium 25.46 and 24.20 mg/100 g, magnesium 21.48 and 28.10 mg/100 g, zinc 2.42 and 5.22 mg/100 g, sodium 0.51 and 1.25 mg/100 g, potassium 126.80 and 5.01 mg/100 g and phosphorus 0.62 and 1.24 mg/100 g for the vegetables and fruits respectively. However, *Ficus capensis* had some anti-nutrients such as tannin 4.28 and 1.67 mg/100 g, cyanide 1.97 and 0.09 mg/100 g, oxalate 5.10 and 2.64 mg/100 g, phytate 9.26 and 7.01 mg/100g and saponin 2.97 and 1.20 mg/100 g for the vegetables and fruits respectively.

*Corresponding author: Email: nkemumerah@yahoo.com;

The results showed that the fruits and vegetables of *Ficus capensis* is an underutilized fruits and vegetable with high nutrient profile especially iron which when incorporated to the diet could help to fight iron in a community with high prevalence of anemia.

Keywords: *Ficus capensis*; phytochemicals; anti-nutrient; anemia.

1. INTRODUCTION

High consumption of plant foods are associated with numerous health benefits rooted in their various physiological effects as a result of their antioxidant, phytochemical and nutritional composition. The use of plants as medicine in treatment of some disease is an ancient and reliable practice. Plant materials and products continue to play an important role in the maintenance of human health since antiquity. Several plants are now being used in part or as a whole to treat many diseases [1].

High intake of fruit and vegetables has been linked epidemiologically with reduced risk of many non-communicable diseases. The important of antioxidants which fruits and vegetables are good sources of, play a role of scavengers cleaning up free radicals before they cause detrimental health effects [2]. The high fiber content of fruits and vegetables may play a role in calcium absorption and reduce the 'acid load' of the diet [3], enhancing bone formation and suppressing bone resorption which consequently result in greater bone strength [4]. Moreover, phytochemicals in fruits and vegetables such as isoflavones have shown to be protective against lens damage which occurs due to hyperglycemia [5] and certain flavonoids such as quercetin can prevent oxidative stress in the pathogenesis of glaucoma [6]. Sufficient fruits and vegetables intake is also associated with lower risk of cognitive decline hence proved beneficial for mental health due to the present of iodine [7].

Fig is a native of Western Asia and Africa. The first creature (Adam and Eve) ate the figs and made a covering (clothing) of their naked bodies using the leaves. It is called "Opoto" among Yoruba in Nigeria [8]. Fig tree belongs to the order of *Urticales* and family of *Moraceae*. Figs are used as an excellent source of minerals, vitamins, carbohydrates, and dietary fibre because it is fat and cholesterol free and contain high number of amino acids [9]. *Ficus capensis* are a widespread medicinal plant species of fruit and vegetable commonly grown, especially in warm, dry climates. It has moisture content of

80.2%, ash 4%, carbohydrates 16.3%, fat 0.53% and protein 0.53% [10].

2. MATERIALS AND METHODS

Ficus capensis vegetables and fruits were collected from the ornamental garden in Amankpume Ozzi-Edem Nsukka Local Government Area, Enugu state, Nigeria.

2.1 Sample Preparation

Ficus capensis leaves and fruits were separately plucked and sorted by removing extraneous materials and cleaned by washing with deionized water. They were milled using electric blender until the desired particle size was obtained (150 - 850 microns).

2.2 Chemical Analysis

2.2.1 Proximate composition

The moisture, protein, fat, fibre and ash content of the samples were determined using the method of AOAC [11,12].

2.3 Determination of Carbohydrate Content

Carbohydrate content was calculated by difference. The estimated percentages of crude protein, ash, fat, fibre and moisture was summed up and the value subtracted from 100%.

$CHO = 100\% - \% (\text{protein} + \text{fat} + \text{ash} + \text{fibre} + \text{moisture})$.

2.4 Mineral Determination

The mineral contents, namely: Na, K, Ca, Mg, Cu, Mn, Hg and Pb contents were determined by the method described by Pearson [13] using a Pye Unicam SP9 Atomic Absorption Spectrophotometer (AAS) connected to an SP9 computer (Pye Unicam Ltd, York Street, Britain). Total phosphorus was determined by the spectrophotometric molybdovanadate [13].

2.5 Vitamin Determination

2.5.1 Determination of β -carotene

The determination of carotenoids was carried out according to the method of Seo et al. [14] with slight modifications.

2.5.2 Determination of vitamin E profile

Vitamin E content was analysed by the method described by Burri [15] using High performance liquid chromatography (HPLC).

2.5.3 Vitamin C determination

Vitamin C determination by iodine titration as described by Anne Helmenstine was carried out [13].

2.6 Phytochemical Screening

A small portion of the extract was subjected to the phytochemical test using Traese and Evans [16] and Harbourne [17] methods to test for alkaloids, flavonoids, saponins, lycopene, phenol and cardiac glycoside. The Folin-Denis Spectrophotometer method was used to determine the tannin content of the foods. The method was described by Pearson [13].

2.7 Anti-Nutrient Determination

Cyanide was determined by Wang and Filled method [18]. Phytate was determined from duplicate samples of food using diluted HCL [19]. Oxalate determination was carried out as described by [20].

3. RESULTS AND DISCUSSION

3.1 Proximate Composition of *Ficus capensis* Fruits and Vegetables

Protein: The protein content of *Ficus capensis* vegetables and fruits were 6.11 and 8.02% respectively. The fruit had higher protein content. The protein content of the vegetable was slightly higher than the findings of Al-Snafi [21] with value of 5.90% of *Ficus capensis*. The protein content of the fruit was lower than the findings of [22] with the protein value of 15.03% on *Ficus capensis* fruit. The low protein content of *Ficus capensis* might be attributed to the fact that fruits and vegetables are not good source of protein. Protein is vital for various body functions such as

body development, maintenance of fluid balance, formation of hormones, enzymes and sustaining strong immune function [23].

Ash: The ash content of the vegetables and fruits were 7.63 and 2.92% respectively with the vegetables having higher ash content compared to that of fruits. The ash content of the test samples were higher than the value recorded by Al-Snafi [21] with 5.30%, and 4.65% for vegetable and fruits of *Ficus capensis* respectively. The variation in the ash content could be due to the nature and the age of the sample used. Ash is the measure of mineral matter in food. Measuring ash content is important because mineral matter may be the cause of a pharmacological effect [23].

Fibre: The fibre content of the vegetable and fruits were 9.82 and 6.33% which were slightly higher than the research conducted by Jyoti et al. [24] and Al-Snafi [21] with fibre mean values of 7.50 and 3.68% respectively on *Ficus capensis* vegetables. Crude fiber content of this plant could aid in the absorption of trace elements in the gut and therefore increases intestinal bowel movement [25]. Consuming vegetables in our diet could aid in managing constipation problems [26]. Dietary fibers also lower cholesterol, triglycerides and protect against cancer and digestive disorders.

Moisture: The moisture content of the *Ficus capensis* vegetable and fruit ranged from 43.28 to 45.20%. There was higher moisture in *Ficus capensis* fruits (45.20%) than its vegetables (43.28%). The values obtained in this research were lower than the findings of Onuekwe, [27] and Al-Snafi, [21] with values of 63.39% and 60.90% on *Ficus capensis* fruits and vegetables respectively. The moisture content of the samples shows that the most single constituent of fruits and vegetable is water. The high moisture content in these vegetables and fruits were not a surprise as Ene-Obong [28] noted that the most single constituent of fruits and vegetable is water, which accounts for more than 80% of the nutrients.

Carbohydrate: The carbohydrate content of the vegetable was lower than that of the fruit with the values of 31.48±0.01 and 35.61% respectively. This finding was higher than the research work recorded by Adebisi and Oyeleke, [8] with the *Ficus capensis* fruits and vegetables values of 25.9% and 30.93% respectively. Carbohydrates are known to produce energy required for the

body because they are essential nutrient required for adequate diet and supplies energy to cells such as brain, muscle and blood.

Fat: On the basis of the fat, the mean value of the vegetable and fruit were 1.68 and 1.92% with the fruit having the higher fat while the vegetable had the least fat content. These values were also higher than the values obtained by Al-Snafi [21] and Jyoti et al., [24] for fruits and vegetables of *Ficus capensis* with fat content of 0.56% and 0.96% respectively. Leafy vegetables and fruits are poor sources of lipids, therefore the increase consumption of fruits and vegetables would naturally lower fat intake and will not predispose the consumers to cardiovascular diseases associated with consumption of fatty foods.

3.2 Mineral Composition of *Ficus capensis* Vegetables and Fruits

Iron: The iron content of the vegetable was 14.24mg/100g and the fruits was 11.68 mg/100 g. However, these values agreed with the findings of Adebisi and Oyeleke [8] that recorded iron values of 14.72 and 16.60 mg/100 g for the *Ficus capensis* fruits and vegetables respectively. Iron plays numerous biochemical roles in the body, including oxygen binding in hemoglobin and acting as an important catalytic centre in many enzymes such as the cytochrome oxidase [29]. The RDA for iron is 8 mg/day indicating that the studied fruits and vegetable could be recommended in diets for reducing anemia, which affects over one million people worldwide [30]. It is also essential for hemoglobin formation and plays a role in energy transfer within the plant and also an essential constituent of certain enzymes and proteins [31]. This justifies the use of *Ficus capensis* in folklore medicine as a blood tonic because of its blood boosting effect [32]. The recommended daily requirement for iron is 10mg, as such 100g portion of *Ficus capensis* specie could supply substantial amount of iron to the body [27].

Calcium: This research also revealed that the calcium content of the *Ficus capensis* vegetable (25.46 mg/100 g) was higher than that of the fruit (24.20 mg/100 g) which was lower than the findings of Ihedioha et al. [22] with the mean values of 35.46 and 34.49mg/100 g for *Ficus capensis* vegetables and fruits respectively. Calcium is reported to be essential for blood clotting, bone and teeth formation and as a co-factor in some enzyme catalysis [33]. The current

daily requirement of calcium for children 4-8 years is (210 mg).

Magnesium: The mean value of magnesium vegetables and fruits were 21.48 and 28.10 mg/100 g respectively which was higher in the fruits than the vegetables. The mean values of *Ficus capensis* by Ihedioha et al. [22] were 18.80 and 22.48 mg/100 g for vegetables and fruits respectively. In humans, magnesium is required in the plasma and extracellular fluid, where it helps maintain osmotic equilibrium [31]. It can also prevent some heart disorders and lower blood pressure in humans.

Zinc: The Zinc content in *Ficus capensis* fruits was 5.22 mg/100 g and the vegetable mean value was 2.42 mg/100 g while the vegetable agreed with the findings of Ihedioha et al. [22] with the value of 3.60 mg/100 g but the vegetable was lower than the value 17.22 mg/100 g. However, Pathak and Kapil [34] reported that zinc is vital in protein synthesis, cellular differentiation and replication, immunity and sexual functions. One hundred grammes (100 g) portion of *Ficus capensis* fruit could supply up to half of the daily requirement of the nutrient. Also one hundred grammes (100 g) of the leaves could supply up to one quarter of the daily requirement of the nutrient.

Sodium: There were 0.51 and 1.25 mg/100 g mean value of sodium in the vegetables and fruits respectively. Adebisi and Oyeleke, [8] observed mean value of 10.53 and 8.60 mg/100 g in the sodium content of vegetables and fruits. Sodium regulates the body fluids and also transmits electric impulses within the human body. It is also one of the most significant constituents of nerves while it also helps in regulating muscle contractions. High consumption of sodium predisposes one to hypertension, therefore taking this plant would not raise the blood sodium of individuals.

Potassium: Potassium level in the vegetable was (126.80 mg/100 g) and the mean value in the fruits was (5.01 mg/100 g) while the findings of Ihedioha et al. [22] recorded mean value of 113.01 mg/100 g for the vegetable of *Ficus capensis*. However, the fruits mean value agreed with the findings of Adebisi and Oyeleke, [8] with the value of 5.70 mg/100 g for *Ficus capensis* fruits. Moderate quantities of sodium and potassium were present in the leaves fruits and vegetables of *Ficus capensis* and these are principal cations of extracellular and intra-cellular

fluids and aid in maintaining electrolyte balance in the body [33]. Potassium is essential for proper growth and plant reproduction.

Phosphorus: The Phosphorus level in the vegetable and fruits were 0.62 and 1.24 mg/100 g respectively. It was observed that the fruits had higher phosphorus than that of the vegetables. The values obtained in this research were in line with the mean value of phosphorus in the vegetables 0.35 mg/100 g [22] but lower than the findings of Adebisi and Oyeleke, [8] that had 187.25 mg/100 g. Phosphorous maintain blood sugar levels and normal heart contraction. It is also important for normal cell growth and repair, bone growth and kidney function. It plays an important role in maintaining the body's acid-base balance [23].

3.3 Vitamins Composition of *Ficus capensis* Fruits and Vegetables

Beta-carotene: The beta-carotene content of the fruits and vegetables were 2500.00 and 450.00 RE respectively. There was higher beta carotene in fruits than the vegetables. Beta carotene is invaluable for the promotion of growth of cells and tissues, resistance to diseases and for delaying the ageing process. It is also important for the maintenance of eye, skin, nails and hair health. The RDA for Beta-carotene for a normal healthy, active adult man and non pregnant woman is 0.3mg/day and 0.27 mg/day respectively [30]. The beta carotene content for fruits and vegetables in this study suggests that the foods may be capable of providing adequate levels of beta carotene for healthy living.

Vitamin E: Vitamin E content were 4.28 and 2.11 mg/100 g for the vegetable and fruits respectively. According to FAO [35] the RDA requirement for vitamin E is 10 mg/day for normal healthy adult men between the ages of 19-65 years while that of adult non-pregnant women within same age range is 7.5 mg/day. Vitamin E is a very potent antioxidant that helps to protect body cells from damage due to reactive oxygen species. It is very important for the formation and normal function of erythrocytes and muscles [36].

Vitamin C: The results of vitamin C content of the samples were 3.45 and 5.26 mg/100 for vegetables and fruits respectively. Vitamin C is a potent antioxidant that facilitates nonhaem iron transport and uptake at the intestinal mucosa, the reduction of intermediates of folic acid as well

as the synthesis of cortisol. It also aids in the purification of blood [37]. The recommended daily requirement for Vitamin C according to FAO [30] is between 45.83 mg/day to 68.50 mg/day for both male and female adults between the ages of 19 to 65 years.

3.4 Phytochemicals and Antioxidant Content of *Ficus capensis* Vegetables and Fruits

Alkaloid: The alkaloid contents were 0.32 mg/100 g and 0.67 mg/100 g for the vegetables and fruits respectively. Alkaloids are used as medicinal agents for analgesic, antispasmodic, and bactericidal effects [38]. The presence of alkaloids in the fruits and vegetables is of interest because of its health benefit.

Glycosides: Glycosides content of vegetable was (2.38 mg/100 g) and that of fruits was (0.42 mg/100 g). These values were lower than the critical level 20 mg/100 g [39]. High level of glycoside causes toxicity which shows symptoms of diarrhea, vomiting, and heart failure in human. Studies showed that high level of glycoside could be reduced during processing such as soaking, malting, fermentation, boiling or frying [40]. The health benefit of glycoside is the fight against cancer and it helps in reducing pain associated with arthritis and also, in lowering high blood pressure (www.brighthub.com/health/alternative).

Lycopene: The lycopene level in the fruits and vegetables were 0.72 and 6.24 mg/100 g respectively. Consumption of *Ficus capensis* fruits with high lycopene level should be advocated to harness the health benefit of the fruits. Lycopene is an antioxidant that protects the body from damages caused by free radical, stress which can hurt DNA and other cell structures [41].

Flavonoids: The flavonoid values for the vegetable was 0.35% while the values for the fruits was 2.58%. The use of *Ficus capensis* fruits and vegetables in food consumption is of immense benefit not only for food diversification but also due to their health benefit. Flavonoids lower high blood pressure and have strong anti-inflammatory properties [42]. Flavonoids are potent anti-oxidants. They also inhibit low density lipoprotein (LDL) by free radicals and reduce the risk of cancer and Cardiovascular diseases [43]. Flavonoids are also involved in platelet aggregation, antimutagenic and antiproliferative properties [44].

Phenol: The level of phenol in the studies vegetable and fruit were 0.48 and 3.21 mg/100 g respectively. The antioxidant activity of phenolic compounds is attributed to the capacity of scavenging free radicals, donating hydrogen atoms, electrons, or chelate metal ion [45].

3.5 Anti-Nutrient Composition of *Ficus capensis* Fruits and Vegetables

Tannin: The tannin content of the vegetables and fruits were 4.28 and 1.67 mg/100 g. Although the tannins were higher in the vegetable than that of the fruits. The findings were lower than the values recorded by Achi et al. [23] that had the tannin content of 687.64 mg/100 g for the vegetables of *Ficus capensis* while Onuekwe [27] reported the tannin content of 0.05 mg/100 g. Tannin act as antinutrient when the value is above safe level but below safe level (0.15-0.20%) it functions as phytochemicals. These fruits and vegetables should be subjected to different food processing methods to reduce the tannin level and extend their food uses. Holz and Gibson [46] suggested that many traditional methods of food preparation such as fermentation, cooking and malting increases the nutritive quality of plant foods through reducing certain anti nutrients such as phytic acid, tannins, polyphenols and oxalic acid. Subjecting the vegetables and fruits to these processes will reduce the toxic level and at the same time boost the phytochemical properties of the vegetables [36]. Tannins may be employed medically in anti-diarrheal, haemostatic and antihemorrhoidal treatment. The anti-inflammatory effects of tannins help to control all indications of gastric enteritis and irritating bowel disorders. Tannins not only heal burns and stop bleeding, but they also stop infection while they continue to heal the wound internally.

Cyanide: The Cyanide content of the vegetables and fruits were 1.97 mg/100 g and 0.09 mg/100 g respectively. The cyanide levels (1.97 mg) and (0.09 mg) in the vegetables and fruits were below the toxic limit for cyanide (35 mg) [47]. Cyanide is a toxin affecting the host when consumed in large quantity. The low levels of cyanide in the vegetables and fruits studied suggest that cyanide content of these food may not pose a threat to the consumers.

Oxalate: The oxalate content of the vegetables was 5.10 mg/100 g while that of the fruits was

2.64 mg/100 g. The mean values in this study were similar with the research work of 8.54 and 2.74 mg/100 g recorded by Ihedioha et al. [22] on vegetables and fruits of *Ficus capensis* respectively. It is known that oxalate chelates calcium to precipitate its deficiency thereby causing it not to be bioavailable. The values obtained in this study were higher than the toxic limit for oxalate (2.20 mg) [48]. Holz and Gibson [46] suggested that many traditional methods of food preparation such as fermentation, cooking and malting increases the nutritive quality of plant foods through reducing certain anti nutrients such as phytic acid, polyphenols and oxalic acid.

Phytate: The phytate content of *F. capensis* vegetables and fruits were 9.26 mg/100 g and 7.01 mg/100 g respectively. The values obtained in this research were similar to the findings of Ihedioha et al. [22] with the phytate value of 8.54 mg/100 g and 8.64 mg/100 g for the *F. capensis* vegetables and fruits respectively. The toxic limit for phytate is (5.00 mg/100 g) [47]. It will be necessary to employ some processing techniques on the fruits and vegetable before consumption since their levels are beyond the toxic limit for phytate.

Saponin: The result showed that the saponin content was (1.20 mg/100 g) for the fruits and (2.97 mg/100 g) for vegetables. Ihedioha et al. [22] observed that the saponin level in *Ficus capensis* fruits was 0.50 mg/100 g while Achi et al. [23] reported that the mean value of saponin in the vegetables of *Ficus capensis* was 0.27 mg/100 g. Saponin is an antinutrient as well a phytochemical at safe level. The saponin level in the fruits and vegetable were below 3.00 mg which was reported by Kumar [49] to be responsible for cattle losses when they grazed on alfalfa. Saponins have expectorative, anti-inflammatory and immune stimulating activity. They also demonstrate antimicrobial properties particularly against fungi, bacteria and protozoa [48]. There is evidence of the presence of saponins in traditional medicine preparations [50,51,52]. Saponins are bitter and reduce the palatability of food and increase excretion of cholesterol concentration by free radicals that are bond with cholesterol and other pathogens in the body. Saponin decreases tumor size and improves cognitive ability [53].

Table 1. Proximate composition of *Ficus capensis* fruits and vegetables (%/100 g)

Samples	Protein	Ash	Fibre	Moisture	Carbohydrate	Fat
F. Vegetable	6.11±0.01	7.63±1.01	9.82±0.42	43.28±0.67	31.48±0.01	1.68±0.45
F. Fruits	8.02±0.24	2.92±0.33	6.33±0.05	45.20±0.22	35.61±0.36	1.92±0.18

Values are mean ± standard deviation of 3 replicate

Table 2. Mineral composition of *Ficus capensis* fruits and vegetables (mg/100 g)

Samples	Fe	Ca	Mg	Zn	Na	K	P
F. Vegetable	14.24±0.18	25.46±0.27	21.48±0.27	2.42±0.34	0.51±0.02	126.80±0.26	0.62±0.05
F. Fruits	11.68±0.14	24.20±0.08	28.10±0.06	28.10±0.06	1.25±0.61	5.01±0.72	1.24±0.12

Values are mean ± standard deviation of 3 replicate

Table 3. Vitamin composition of *Ficus capensis* fruits and vegetables

Samples	Beta-Carotene	Vitamin C	Vitamin E
F. Vegetable	450.00±1.26	3.45±0.23	4.28±0.11
F. Fruits	2500.00±2.01	5.26±0.28	2.11±0.07

Values are mean ± standard deviation of 3 replicate

Table 4. Phytochemical composition of *Ficus capensis* fruits and vegetables (mg/100 g)

Samples	Flavonoids	Lycopene	Alkanoid	Phenol	Glycoside
F. Vegetable	0.35±0.06	0.72±0.38	0.32±0.01	0.48±0.06	2.38±0.70
F. Fruits	2.58±0.07	6.24±0.21	0.67±0.11	3.21±0.24	0.42±0.18

Values are mean ± standard deviation of 3 replicate

Table 5. Anti-nutrient composition of *Ficus capensis* fruits and vegetables (mg/100 g)

Samples	Tannin (mg/100 g)	Cyanide (mg/100 g)	Oxalate (mg/100 g)	Phytate (mg/100 g)	Saponin (mg/100 g)
F. Vegetable	4.28±0.77	1.97±0.46	5.10±0.11	9.26±0.92	2.97±0.25
F. Fruits	1.67±0.21	0.09±0.24	2.64±0.82	7.01±0.68	1.20±0.02

Values are mean ± standard deviation of 3 replicate

4. CONCLUSION

The results showed that the fruits and vegetables of *Ficus capensis* is an underutilized fruits and vegetable with high nutrient profile especially iron which when incorporated to the diet could help to fight iron in a community with high prevalence of anemia.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ajeigbe KO, Enitan SS, Omotoso DR, Oladokun OO. Acute effects of aqueous leaf extract of *Aspilia africana* c.d. adams on some haematological parameters in rats. African Journal Traditional Complement Alternative Medicine. 2013; 10(5):236-243
2. Kaur C, Kapoor HC. Antioxidants in fruits and vegetables: the millennium's health. Int J Food Sci Technol. 2001;36: 703–725. [Google Scholar]
3. New S. Fruit and vegetable consumption and skeletal health: is there a positive link? Nutrition Foundation. Nutr Bull. 2001;26: 121–125. [Google Scholar]

4. Shen CL, Bergen VV, Chyu MC, et al. Fruits and dietary phytochemicals in bone protection. *Nutr Res.* 2012;32:897–910. [PubMed] [Google Scholar]
5. Agte V, Gite S. Diabetic Cataract and Role of Antigliating Phytochemicals. In: *Handbook of Nutrition, Diet and the Eye.* Ed, Preedy VR. 1st ed, Elsevier Inc; Burlington. 2014;131–140. [Google Scholar]
6. Miyamoto N, Izumi H, Tawara A, Kohno K. Quercetin and Glaucoma. In: *Handbook of Nutrition, Diet and the Eye.* Eds, Preedy VR. 1st ed, Elsevier Inc; Burlington. 2014; 97–103. [Google Scholar]
7. Mc Martin SE, Jacka FN, Colman I. The association between fruit and vegetable consumption and mental health disorders: Evidence from five waves of a national survey of Canadians. *Prev Med.* 2013;56: 225–230. [PubMed] [Google Scholar]
8. Adebisi GA, Oyeleke GA. Studies on *Ficus capensis* (fruit and leaf): Proximate and mineral compositions. *International Journal of Chemical Science.* 2009;7(3): 1761-1765.
9. Guarrera PM. Traditional phytotherapy in Central Italy (Marche, Abruzzo, and Latium), *Fitoterapia.* 2005;76(1):1–25.
10. Ambika C, Beenu T, Intelli C. Influence of Processing on Physicochemical, Nutritional and Phytochemical Composition of *Ficus capensis* (Fig) Fruit. *Research Journal of Pharmaceutical, Biological and Chemical Sciences.* 2015;6(6):1474-1489.
11. Association of Official Analytical Chemist (A.O.A.C). Official method of analysis, 16th edition. Association of Official Analytical Chemist. Washington, D.C. 2010;536–541.
12. AOAC. Official Methods of Analysis. Association of Official Analytical Chemists, Washington, DC; 201022 USA; 2005.
13. Pearson IO. *Fundamental of food biochemistry*, 2nd Ed. Atlanta, Georgia, 30322 USA; 1976. Available:www.en.wikipedia.org/wiki/specia l
14. Seo JS, Burri BJ, Quan Z, Neidlinger TR. Extraction and chromatography of carotenoids from pumpkin. *American Journal of Chromatography.* 2005;1073: 371-375
15. Burri BJ. Analysis of vitamin E by HPLC. *The Encyclopedia of Vitamin E*; 2007.
16. Traese GE, Evans WC. *Pharmacognocny.* 14th Ed. Brown Publication. London. 1989; 30-38.
17. Harbourne JB. *Phytochemical methods: A guide to modern technique of plants analysis.* Chapman and Hall: London. 1983;60-64.
18. Wang JK, Filled SM. *Food quality evaluation, an approach.* Lan. Lancet Press. 1980;123–130. Available:http://jds.fass.org/cgi/pmidlookup ?view=long&pmid=17106105
19. Suree N, Surat K, Akekachai N. Phytate and fiber content in Thai fruits commonly consumed by diabetic patients. *Journal of Medical Association, Thai.* 2004;87(1): 1444-6. Available:http://www.medassocthai.org/journal
20. AOAC. Association of official analytical chemist. *Official Methods of Analysis*, Washington, D.C.; 2000.
21. Al-Snafi AE. Nutritional and pharmacological importance of *Ficus capensis* - A review.*Journal of Pharmacy.* 2017;7(3):33-48 33.
22. Ihedioha JN, Ekere NR, Ibezim A, Okoye COB. Nutritional Evaluation of Leaves and Fruits of *Ficus Capensis*.*Journal Of Chemical Society of Nigeria.* 2015;40(2): 60-66
23. Achi NK, Onyeabo C, Ekeleme-Egedigwe CA, Onyeonula JC. Phytochemical, Proximate Analysis, Vitamin and Mineral Composition of Aqueous Extract of *Ficus capensis* leaves in South Eastern Nigeria. *Journal of Applied Pharmaceutical Science.* 2017;7(03):117-122.
24. Jyoti DV, Dipak V, Sneha RP, Aditi UP, Sheeba S. Biochemical, Organoleptic Assessment of Fig (*Ficus capensis*).*Journal of Biotechnology and Biochemistry.* 2017;3(2):95-104.
25. Abolaji OA, Adebayo AH, Odesanmi OS. Nutritional qualities of three medicinal plant parts (*Xylopi aethiopic a*, *Blighia sapida* and *Parinari polyandra*) commonly used by pregnant women in the Western Part of Nigeria. *Pakistan Journal of Nutrition.* 2007;6(6):665-668.
26. Olowokudejo JD, Kadiri AB, Travih VA. An ethnobotanical survey of herbal markets and medicinal Plants in Lagos State of Nigeria.*Ethnobotanical Leaflets.* 2008; 12(3):851-865.
27. Onuekwe ME. Chemical composition of some lesser-known wild fruits and

- vegetables consumed in Ayamelum Local Government Area of Anambra State. A Project Report for the Degree of Masters of Science in Human Nutrition, Department of Home Science, Nutrition and Dietetics, Faculty of Agriculture, University of Nigeria, Nsukka, Enugu State, Nigeria. 2012;55-56.
28. Ene-Obong HN. Native species in National food consumption system. A paper presented during the inauguration of the R and D Team for the National Programme on indigenous crops and animals by the Federal Ministry of Science and Technology at the conference Hall of National centre for Genetic Resources and Biotechnology, Moor Plantation, Ibadan. 1998;5-34.
 29. FAO. Diet, nutrition and the prevention of chronic diseases. Scientific Background Papers of the Joint WHO/FAO Expert Consultation. Geneva, Switzerland. 2002; 60-63.
 30. Emmanuel O, Ugbogu AE, Odungide AA, Atani CS. Proximate, antinutritional and mineral estimation of some selected consumed green vegetables in Afaha Eket, Akwa-Ibom State, Nigeria. *American Journal of Food Science Technology*. 2017;5(5):182–191.
 31. Thomas RA, Krishnakumari S. Proximate analysis and mineral composition of *Myristica fragrans* seeds. *Journal of Pharmacognosy and Phytochemistry*. 2015;3(6):39-42.
 32. Njoku-Oji NN, Nwike CV, Dimkpa U, Ifegwu NO, Anike LC, Maduka SO, Sobanke OA, Uchefuna RC. Hematological changes following oral administration of aqueous leaf extract of *Ficus capensis* in albino rats. *International Blood Research and Reviews*. 2016;5(1):1-7.
 33. Robert KM, Daryl KG, Peter AM, Victor WR. Harper's Illustrated Biochemistry. In Benders and Mayes Vitamins and Minerals, Lange Medical Books/McGraw-Hill, Medical Publishing Division, New York, USA. 2003;496.
 34. Pathak P, Kapil U. Role of trace elements zinc, copper and magnesium during pregnancy and its outcome. *Indian Journal Paediatric*. 2004;71(2):1003-1005.
 35. Rumeza H, Zafar I, Mudassar I, Shaheena H, Masooma R. Use of vegetables as nutritional food: Role in human health.
 36. Dias JS. Nutritional quality and health benefits of vegetables: A review. *Food and Nutrition Sciences*. 2012;3(10):1354-1374.
 37. FAO. Human vitamin and mineral requirements. Report of a Joint WHO/FAO Expert Consultation. Bangkok, Thailand. 2001;34-40.
 38. Schiavone A, Guo K, Tassone S. Effects of a natural extract of chestnut wood on digestibility, performance traits, and nitrogen balance of broiler chicks. *Poultry Science*. 2008;87(3):521–7. DOI:10.3382/ps.2007-00113 PMID:18281579 Available:<http://ps.fass.org/cgi/pmidlookup?view=long&pmid=18281579>
 39. Edeoga HO, Okwu DE, Mbaebia BO. Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnol*. 2005;4(7):685–92.
 40. WHO. Dietary intake of fruits and vegetables and risk of diabetes mellitus and cardiovascular diseases, WHO Report Geneva. 2005;23-25.
 41. Dashiell E. The health benefits of lycopene (www.verywellhealth.com); 2020.
 42. CSIRO. Reduction of root flavonoid level and its potential; 2004. Available: www.publish.csiro.au
 43. Verena S, Mario L, Karl S. The role of tea and tea flavonoids in cardiovascular health. *Journal of Nutritional Food Resources*. 2006;50:218-228.
 44. Subramani S, Casimir CA. Flavonoids and antioxidant activity of Georgia; 2002.
 45. Afanas'ev IB, Dcrozshko AI, Brodskii AV, Kistyuk VA, Potapovitch AI. Chelating and free radical scavenging mechanisms of inhibitory action of rutin and quercetin in lipid peroxidation. *Biochem Pharmacol*. 1989;38:1763-1769. Doi: 1016-2952 (89) 90410-3.
 46. Hotz C, Gibson RS. Traditional foodprocessing and preparation practice to enhance the bioavailability of micronutrients in plant-based diets. *Journal of Nutrition*. 2007;137(4):1097–100. [PMID 17374686]
 47. Munro A, Bassir O. Oxalate in Nigeria vegetables. *West African Journal of Biological, Agriculture and Chemistry*. 1969;12:14-17.
 48. Abbot WS. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol*. 1925;18:265-267. Available:www.rayahelian.com

49. Kumar K. Conducting focus group interviews in developing countries. A.I.D. Program Design and Evaluation Methodology Report No. 8. Washington, D.C. U.S. Agency for International Development; 1987
50. Asl MN, Hosseinzadeh H. Review of pharmacological effects of Glycyrrhiza spp and its bioactive compounds. *Phytother Resources*. 2008;22(6):709-724.
51. Hostettan K, Marston A. Saponins: Chemistry and pharmacology of natural products. Cambridge University Press, Cambridge, UK; 1995.
52. Hussain ZJ, Muhammad R, Ullah FU, Khan IU, Khan N, Khan J, Ali, Jan S. Evaluation of the chemical composition of *Sonchus eruca* and *Sonchus asper*, *Journal of American Science*. 2010;6(9): 231-235.
53. Malinow MR, McNaughty AL, Kohler GO. Effects of synthetic glycosides on cholesterol absorption. *Ann. N.Y. Acad. Sci*. 1985;23:454.

© 2022 Umerah et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/62391>