

Vacciplant and Calliete Cotreatment Synergically Induces Phenolic Compounds Production in Plantain (*Musa x paradisiaca* L.) and Trigger the Natural Defense Activation for Protection to Black Leaf Streak Disease

Gogbeu Dan Gbongué Lucien ^{a*}, Yeo Navigué Abou ^a, N'cho Achi Laurent ^b, Traore Siaka ^b, Kouakou Tanoh Hilaire ^c and Kone Daouda ^a

^a Université Félix Houphouët-Boigny, UFR Biosciences, 22 BP 582 Abidjan 22, Côte d'Ivoire .

^b Centre National de Recherche Agronomique, 01 BP 1536 Abidjan, Côte d'Ivoire.

^c Université Nangui Abrogoua, UFR Sciences de la Nature, 02 BP 801 Abidjan 02, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. Author GDGL designed the study, wrote the protocol and the first draft of the manuscript. Author YNA and NAL performed the statistical analysis. Authors KTH and ZM managed the literature review, curation and editing. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJA/2022/v15i330158

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86648>

Original Research Article

Received 14 February 2022

Accepted 26 April 2022

Published 27 April 2022

ABSTRACT

Aims: To evaluate the synergistic action of two elicitors or stimulators of natural plant defense, vacciplant and calliete, on the synthesis of defense metabolites and their impact on tolerance to *Mycosphaerella fijiensis*, causal agent of the black leaf streak disease (BLS) in plantain (*Musa x paradisiaca* L.).

Place and Duration of Study: The study was conducted in Abidjan (southern Côte d'Ivoire) in 2019.

Methodology: The ability of plants co-treated with vacciplant+calliete to product phenolic compounds was compared to vacciplant and the control. Subsequently, the treated plants were inoculated with the pathogenic filtrate of *Mycosphaerella fijiensis* in order to induce the black leaf

streak disease. Afterwards, the content of phenolic compounds was determined and then their analysis by ultra-high performance liquid chromatography was performed (U-HPLC). Finally, foliar pigments were quantified to have an indication of the sanitary state of elicited and infected plants.

Results: plants treated with vacciplant and inoculated, as well as plants co-treated with vacciplant+calliete and inoculated, accumulated more phenolic compounds than the control plants. However, the phenol content of the co-treated plants (191.10 mg/g FL) was higher than that of the plants treated with vacciplant alone (142 mg/g FL). In addition, qualitative analysis by U-HPLC revealed that treated plants synthesized de novo two compounds including *trans*-resveratrol, a stilbene with significant antifungal activity, in plants co-treated with vacciplant+calliete. In addition, it allowed the plants to acquire a higher leaf greening state than those treated with vacciplant alone, indicating a more intense photosynthetic activity and thus a good sanitary state

Conclusion: Vacciplant+calliete has a synergistic effect on the activation of the plantain's defense reactions and thus on the good tolerance to *M. fijiensis*, an agent of plantain BLSD.

Keywords: Black leaf streak disease; elicitor; *Mycosphaerella fijiensis*; natural defense; polyphenol; plantain; tolerance.

1. INTRODUCTION

Plantain is a food crop that is grown in the countries of the humid tropics for its fruit called plantain. Worldwide, plantain is the fourth most important crop after rice, wheat and maize [1]. Sub-Saharan Africa produces almost 70% of the world's plantain production. Uganda, Cameroon, Ghana, Côte d'Ivoire, Rwanda and the Democratic Republic of Congo are the main producers of plantain [2]. In Côte d'Ivoire, plantain is the second most consumed food after rice [3]. Moreover, it is the 4th most important food crop after rice, cassava and yams [4], with 1,600,000 tonnes produced each year by 2,500,000 active farmers, 80% of whom are women [5]. It thus contributes to the diversification of farmers' incomes and the fight against poverty. However, bananas are subject to numerous parasitic constraints, among which fungal diseases contribute significantly to the drop in yields [6]. Indeed, the fungus *M. fijiensis*, the causal agent of black leaf streak disease (BLSD), is one of the main limiting factors in banana cultivation. The economic consequences of MRN are serious, as losses can represent 50% of the crop [7]. Moreover, the incidence of parasitism is such that the chemical control recommended for intensive cropping systems is inappropriate for plantain because it is a small-scale crop. Indeed, large quantities of pesticides are required to control BLSD, up to 50 fungicide applications per year [7]. Thus, the ecological costs to the environment of using fungicides to control BLSD are high, as are the resulting economic costs. Moreover, Stover and Simmonds [8] evaluated the costs of pesticides at 27% of production. Consequently, it is

necessary to look for effective and accessible alternatives to chemical control for sustainable ecoproduction in plantain. One of these alternatives is to give plants the ability to defend themselves or to increase their defenses instead of fighting the pest directly by using elicitors or natural plant defenses stimulators [9,10]. Moreover, elicitors activate the synthesis of defense compounds without the plant being attacked by pathogens; and among them, phenolic compounds occupy a prominent place [9,11]. The plant is ready to retaliate in the event of a subsequent attack and simultaneously resist a wide spectrum of pathogens [12]. This study aimed to evaluate the synergistic action of two elicitors or stimulators of natural plant defense, vacciplant and calliete, on the synthesis of defense metabolites and their impact on tolerance to *M. fijiensis*, responsible for the black leaf streak disease (BLSD) in banana.

2. MATERIALS AND METHODS

2.1 Plant Material

The plant material consists of vivoplants of plantain cv. French 2. This cultivar is susceptible to BLSD caused by *M. fijiensis* [13]. The plants were provided by the Centre National de Recherche Agronomique (CNRA).

2.2 Fungal Material

The fungal material used is a virulent strain of *M. fijiensis* ST-W-D98.B1C1. It was isolated from the leaves of a plantain tree from the CNRA Wanita experimental plantation in Bimbresso (Côte d'Ivoire).

2.3 Production and Maintenance of Plantain Vivoplants

Plantain vivoplants were produced in a germander from shoots taken from dehulled stumps. After 14 days, they were transferred under shade at room temperature and 80-90% relative humidity. The vivoplants were then grown in perforated black polyethylene bags containing forest floor previously autoclaved at 121°C for 30 min under a pressure of 1 bar, as a substrate [14]. The vivoplants were maintained under shade for three months.

2.4 Preparation of Elicitation Solutions

Vacciplant (45 g/L laminarin) and calliete 80WP (alliete) were prepared respectively at 3 and 2%. A mixture of both solutions (v/v) was also prepared for co-treatment and all the eliciting solutions were supplemented with 20% Tamoil® (SAE 5W-40 oil) at 5 mL/L. Distilled water was used as a control.

2.5 Preparation of the *M. fijiensis* Filtrate

For the preparation of *M. fijiensis* toxic filtrate, 10 mL of inoculum (2.10^5 spores/mL) was added to 100 mL of liquid culture medium contained in Erlenmeyer flasks for the production of toxic metabolites according to Pinkerton and Strobel's method modified by Amari [15,16]. The Erlenmeyer flasks were placed in an enclosure for 28 days. The culture broths were separated from the mycelium by filtration through a sieve 80 µm in diameter. The extract obtained constituted the culture filtrate of *M. fijiensis*. Preparations without inoculation of fungal spores were used as control.

2.6 Treatments for Banana Leaves

The treatment of banana leaves was done according to Belhadj's technique [17]. After three months of cultivation of banana plants, 40 plants of the cultivar French 2 including control-plants (10 plants per treatment) were treated. The solutions were sprayed on the leaves using a hand sprayer and then the plants were incubated in a controlled environment for 48 h and 72 h respectively for the plants co-treated with vacciplant+calliete and treated with vacciplant.

2.7 Foliar Spray Inoculation of Banana with *M. fijiensis* Filtrate

The French 2 banana plants were separated into three distinct groups (untreated, untreated then infected and treated then infected). After the incubation period, the leaves of the banana plants were slightly punctured with a sterile needle. Then, 25 mL of the fungal inoculum was sprayed on the upper and lower surfaces of each banana leaf. Three treatments and controls were carried out for each banana cultivar as (T0) plants not treated by elicitor and not inoculated (PNTNI), (T1) plants not treated with elicitor and inoculated (PNTI), (T2) plants treated by vacciplant and inoculated (PTVI) and, (T3) plants treated with vacciplant+calliete and inoculated PT(V+Ct)I. For each treatment, three plants were used, i.e. 12 plants, and on each plant, three leaves were considered, i.e. 36 leaves. The experiment was thrice repeated.

The inoculated banana plants were placed under shade at room temperature. The plants were watered according to the humidity of the substrate. The leaves of the control plants were sprayed with sterile distilled water instead of elicitors. Two weeks after fungal inoculation, the second leaf after the cigar leaf of each banana cultivar was harvested. The leaves were then freeze-dried and stored at -20°C.

2.8 Quantitative Determination of Phenolic Compounds by Spectrophotometer

2.8.1 Extraction and Phenolic Compounds Essay

The extraction of total phenolic compounds was carried out according to the method of Kouakou [19]. Briefly, 10 mL of methanol was added to 50 mg of freeze-dried leaf and placed overnight at 4°C. The mixture was then centrifuged at 2000 rpm for 10 min. The supernatant was filtered through a Millipore membrane (0.45 µm) and the filtrate was used to determination of phenols content. Briefly, 0.5 mL of Folin-Ciocalteu reagent and 0.9 mL of water were added to 0.1 mL of phenolic extract. After stirring at room temperature, 1.5 mL of a 17% (w/v) sodium carbonate solution was added. After 20 min of incubation at 25 °C in the dark, the mixture was monitored at 765 nm. Phenol content, expressed in milligrams of gallic acid equivalents per gram of extract, was determined using a calibration

line with gallic acid ($y = 0,021x + 0,053$; $R^2 = 0,999$; where y is absorbance and x is the concentration of gallic acid).

2.9 Qualitative Analysis of Phenolic Compounds by Ultra-High Performance Liquid Chromatography

2.9.1 Extraction and Purification of Phenolic Compounds

Extraction and purification of phenolic compounds were carried out as above [18]. Approximately 100 mg of leaf lyophilisate from each cultivar was placed in a haemolysis tube and 20 mL pure methanol was added. The mixture was then placed at 4°C for 15 h. After sonication for 5 min in ultrasound, the mixture was centrifuged at 5,000 rpm for 10 min. Then, 4 mL of supernatant was evaporated using a Speed Vac mini-concentrator. The dry residue obtained was dissolved in 1 mL of 30% methanol and then placed on a C18 grafted silica mini-column in the Supelco Visiprep™ refill system [19]. The resulting eluate was evaporated using Speed Vac; residue was taken up in 1 mL of 50% methanol and filtered through a Millipore membrane at 0.45 µm. The filtrate was used as polyphenols extract.

2.9.2 Analytical Conditions in Ultra-High Performance Liquid Chromatography

The analyses were performed using Ultra High-Performance Liquid Chromatography [20]. An aliquot of the methanolic solution was injected into Ultra High-Performance Liquid Chromatography (UHPLC). The analysis of the samples was performed using the Agilent LC system (1100-1200 series). The UHPLC was coupled to a nuclear magnetic resonance spectrometer (Bruker Avance III) with an operating frequency of 600 MHz for one proton. Separation of phenolic compounds was performed on a reverse-phase C18 silica column (Zorbax Eclipse XDB-C18, 150 x 4.6 mm, 5 µm, Agilent). Distilled water filtered through the millipore membrane (0.45 µm) was acidified with 0.1% trifluoroacetic acid (TFA) to be solvent A and acetonitrile (HPLC grade) acidified by TFA to be solvent B. The elution gradient profile was performed as follows: 2% B (0-1 min), 50% B (1-5 min), 40% B (5-10 min), 20% B (10-15 min), 10% B (15-16 min) and 2% B (16-20 min).

2.9.3 Separation and Identification of Phenolic Compounds

The separation of phenolic compounds was carried out under a pressure of 550 to 1000 bar. The hydromethanolic extract was diluted with filtered distilled water (50/50, v/v). Approximately 10 µL of the purified phenolic compound extract was directly injected into U-HPLC with a flow rate of 1.3 mL/min and detection was performed at 284 nm. A reference library was previously made with commercially available phenolic compounds identified by ¹H-NMR [21,22]. The retention time and/or the NMR spectra of the separated compounds were compared with those of the reference library for their identification.

2.9.4 Determination of Leaf Pigments

The extraction of leaf pigments was performed on the freeze-dried leaves. Thus, 50 mg of freeze-dried leaves were placed in a test tube and 10 mL of acetone was added and then tightly closed with a cap to avoid evaporation. The mixture was then placed on a rotary shaker for 10 min for 2 h. After 5 min sonication with ultrasound, the debris was removed by centrifugation (5000 rpm for 15 min). The absorbance of the crude extract was measured at 470, 647 and 663 nm with a spectrophotometer. Pigment contents are calculated in mg/g of freeze-dried leaves according to the following equations [22]: chlorophyll a (chl a) = 12.25 OD₆₆₃-2.79 OD₆₄₇, chlorophyll b (chl b) = 21.50 OD₆₄₇-5.10 OD₆₆₃, total chlorophylls (Chl t) = 7.15 OD₆₆₃+ 18.71 OD₆₄₇. The indicator of functional pigment equipment and green state of the leaves of elicited banana plants was calculated according to the method of Lichtenthaler [23]. Thus, the indicator was determined by chl a/chl b. In addition, the ratio of total chlorophylls to carotenoids (chl t/car), which indicates the greenness of the plant leaves, was determined.

2.10 Statistical Analysis

The data obtained were analyzed using Statistica 7.1 software. The data were subjected to the test of normality by an analysis of variance. In case of significant differences, the means were classified into homogeneous groups using the Newman-Keuls test at 5%. All experiments were triplicate.

3. RESULTS AND DISCUSSION

Fig. 1 shows the evolution of phenolic compounds in leaves of the banana cultivar French 2 at different treatment stages. The treatment stage (PNTNI) is the phenolic control stage. Indeed, the phenolic content in the control leaves (PNTNI) was 30.50 mg/mg FL. When the leaves were inoculated without elicitation (PNTI), this content decreased from 30.50 to 9.10 mg/mg FL, i.e. a decrease of 70.16% compared to the control. When the leaves were treated and inoculated, an increase in content was observed compared to the control. Thus, in the leaves treated respectively with vacciplant and the vacciplant+calliete mixture and then inoculated, PTVI and PT(V+Ct), 142 and 191.10 mg /mg FL were obtained, corresponding to respective increases of 407 and 526.56% compared to the control. These results determine that without elicitation, the plant is susceptible to the establishment of infection. In addition, vacciplant and vacciplant+calliete accumulated phenolic compounds auguring an acquisition of resistance to *M. fijiensis* [24]. Indeed, the parietal polysaccharides would be elicitors of defensins of phenolic nature. It is worth mentioning that calliete accumulated around 92.40 mg/g FL of phenolic compounds (data not shown) which is very low compared to the other both elicitors. In fact, these elicitors rich in parietal polysaccharides influence the activation of phenolic biosynthetic pathways. Indeed, according to several studies, polysaccharides would hydrolyze into reducing sugars that stimulate the accumulation of phenolic phytoalexins and thus activate defense reactions [25,26]. In addition, phenol production from co-treated plants remains better and is even better when treated and inoculated with *M. fijiensis*.

PNTNI, Untreated and Inoculated Plants; PNTI, Untreated and Inoculated Plants; PTVI, Vacciplant Treated and Inoculated Plants; PT(V+Ct), Plants Treated with Vacciplant+Calliete and Inoculated. Histograms topped with the same letter are not significantly different at 5% (Newman-Keuls test); values represent the mean of triplicate. Bars are standard errors.

These results seem to indicate that vacciplant and its co-treatment with calliete would act as an early salicylic acid to mobilise phenolics in the leaves after elicitation [27]. As a result, a hypersensitivity reaction (HR) would be set up as reported by Beckers and Spoel [28]. HR relies on

the rapid recognition of the pathogen by the plant and would aim to confine the toxic agent to its site of entry. Thus, in many interactions, it appears that the important contribution of the hypersensitive response to host resistance is directly related to the speed with which it is deployed [29]. The accumulation of phenolic compounds suggests that plants treated with these elicitors are able to develop a systemic immunity phenomenon, i.e. a systemic acquired resistance (SAR) against the black leaf stripe disease. The plantain has consequently been "vaccinated" and is ready to fight back in case of a subsequent attack by pathogens such as *M. fijiensis* [30]. In addition, it has long been recognised that plant responses to pathogen attack are characterised by a rapid accumulation of phenolic compounds at the site of infection. In addition, application of elicitor confers enhanced pathogen resistance in many plants was reported [31,32]. Vacciplant and co-treatment would relay the distress signal caused by the presence of elicitors to the nucleus of the attacked cell to induce defence responses specific to the pathogenic attack and this accumulation of phenolic compounds may therefore be an important signal in the acquisition of pathogen tolerance as mentioned in several studies [33,34]. Other studies on the involvement of phenolic compounds in plant resistance to fungi have already been demonstrated [35,36]. Similarly, their involvement in plant tolerance to certain diseases has been proven by recent studies [33,37]. Moreover, phenolic compounds accumulate in tissues adjacent to necrotic areas, suggesting that these compounds may be defensive [38].

The analysis of the phenolic compounds identified (Fig. 2) in plantain leaves reveals that eight constitutive compounds were identified in the non-elicited plant (gallic acid, protocatechic acid, ferulic acid, p-coumaric acid, chlorogenic acid, vanillin, 3-p-coumaroylquinic acid and 3-hydroxybenzoic acid). However, when inoculated, there is no induction of new compounds, only four compounds are identified (gallic acid, protocatechic acid, p-coumaric acid and vanillin). In addition to these constitutive phenolic compounds. Elicited and inoculated-plant induced six new compounds (tryptophan, vanilic acid, caffeic acid, rutin, salicylic acid and *trans*-resveratrol). This plurality of phenolic metabolite biosynthesis has already been reported in cotton [27,39]. Vacciplant and vacciplant+calliete are elicitors that allow a good biosynthesis of phenolic

compounds. Co-treatment induced trans-resveratrol, a stilbene with antifungal action reported in grapevine and cotton [40-41]. They are therefore key molecules in the induction and accumulation of phytoalexins. So, the metabolism of phenolic compounds orientation in banana would be elicitor-dependent

[9,42]. In addition, phenolic compounds were reported to have beneficial actions on plant protection against pathogens [43,44]. Thus, vacciplant and vacciplant+calliote induce systemic acquired resistance (SAR) and potentiate phytoalexins (phenolic nature) accumulation.




Fig. 1. Phenolic compound content of elicited and inoculated leaves of the plantain cultivar French 2




Fig. 2. Chromatographic profiles of phenolic compounds extracted from the leaves of plantain cultivar French 2 at 284 nm

(1) gallic acid (1.486 min); (2) protocatechic acid (2.703 min); (3) tryptophan (3.143 min); (4) vanillic acid (5.067 min); (5) caffeine acid (5.203 min); (6) ferulic acid (6.554 min); (7) p-coumaric acid (7.905 min); (8) rutin (10.011 min); (9) chlorogenic acid (10.405 min); (10) trans-veratrol (12.077 min); (11) vanillin (12.851 min); (12) salicylic acid (13.920 min); (13) 3-p-coumaroylquinic acid (11.082 min); and (14) 3-O-caffeoylquinic acid (13.378 min). PNTNI: untreated and non-inoculated plant; PNTI, untreated and inoculated plant, PTVI, Vacciplant-treated and inoculated plant; PT(V+Ct)I, Vacciplant-treated+calliete and inoculated plant.

Phytoalexins are plant antibiotics synthesised during the hypersensitivity reaction or during HR [35,36]. They accumulate and are thus thought to actively participate in plant defence [45,46]. Their mode of action is linked to their antifungal power, their participation in the reinforcement of plant cell walls and their capacity to modulate and induce host defence reactions [25,47]. In banana, methyl jasmonate was showed to strongly stimulate the accumulation of phenolic phytoalexins, which can induce BLS resistance [48]. Similarly, the involvement of phenolic compounds in the tolerance of plants to parasitic diseases has been reported [11,37].

Table 1 shows the results on indicators of functional pigment equipment and leaf greenery. The Chla/Chlb ratio, which gives the indication of the functional pigment equipment of the leaves, remains lower than that of the control (PNTNI) and untreated and inoculated (PNTI). The same observation was observed in the Chlt/Car ratio, which gives an indication of the state of greenery of the leaves where the value of the PNTI remains below all values but also below 1.00 (reference value of the state of greenery). Thus, the controls (PNTNI) accumulate at the level of the indicator of the functional pigment equipment (Chl a/Chl b) a value of 3.02 while that of the green indicator (Chlt/Car) is 3.7. When the untreated plants are inoculated (PNTI), their Chl a/Chl b ratio is equal to 1.43, i.e. a decrease of 50.17%, while that of Chl t/Car is 0.63, i.e. a decrease of 77.42% compared with the control (PNTNI). When plants were treated with vacciplant® and vacciplant+ calliete followed by inoculation (PTVI and PT(V+Ct)I), Chl a/Chl b ratio was equal to that of the control for PTVI (3.02) and 3.31 for PT(V+Ct)I, an increase of 9.60% compared to the control. However, the green indicator (Chlt/Car) remained statistically identical to that of the control for PTVI, i.e. 3.72.

On the other hand, in PT(V+Ct)I, this Chl t/Car ratio increased significantly to reach 5.56, i.e. a 50% increase. The ratio of chlorophyll a to chlorophyll b was always greater than 1.00 except in unelicited and infected plants (Chl a/Chl b = 0.97). Infection of the plants with the pathogen caused a 67.88% reduction in the functional pigment equipment of the leaves compared to the control.

Chl a, chlorophyll a; Chl b, chlorophyll b; Chl t, total chlorophyll; Car, carotenoids; PNTNI, untreated and non-inoculated plant; PNTI, untreated and inoculated plant, PTVI, plant treated with vacciplant and inoculated; PT(V+Ct)I, plant treated with vacciplant-calliete combination and inoculated; within a column, means followed by the same letter are not significantly different (Newman-Keuls test at 5%).

Analysis of this result suggests a degradation of chlorophyll a under the action of the pathogen in contrast to chlorophyll b. Indeed, the active site of chlorophyll biosynthetic enzymes such as magnesium chelatase and chlorophyllase would be located on chlorophyll a [49]. Thus, a bioconversion of chlorophyll a to chlorophyll b would be plausible as reported by several authors [50-53]. The elicitors used in this study (vacciplant and the vacciplant+calliete co-treatment) would therefore have the role of protecting these enzymatic sites and also chlorophyll a with a view to ensuring important photosynthetic activity to maintain the plant in a good physiological state and thus a good health status. Moreover, Gamon *et al.* [54] reported that the photosynthetic activity of plants is significantly related to the ratio of chlorophyll pigments (total chlorophylls) and carotenoids (Chl t/Car). This ratio is an indicator of the greenness of the leaves [23]. For the elicited plants, the ratio is higher than 3.70 (PNTI), showing a positive effect of the treatment on the green state of the leaves. Moreover, the vacciplant+calliete co-treatment, which gives a value of 5.56, shows that this treatment is more favourable for the green state, which augurs a good photosynthesis, contrary to the vacciplant treatment (3.72), which gives a similar result to the untreated and uninfected plant (3.70). In fact, in the non-elicited plants, the inoculation of the *M. fijiensis* filtrate caused a yellowing of the leaves (data not shown) and thus a very reduced green state (Chl t/Car=1.74). Thus, *M. fijiensis* filtrate appears to cause damage to the plant's photosynthetic apparatus, including chlorophyll a, the major photosynthetic pigment [55].

Table 1. Indicators of functional pigment equipment and leaf green status in French 2 plantain elicited and inoculated with *Mycosphaerella fijiensis*

Treatment	Functional pigment equipment indicator	Indicator of leaf greening status
PNTNI	3.02 ± 0.03 ^b	3.70 ± 0.04 ^b
PNTI	0.97 ± 0.05 ^c	1.74 ± 0.09 ^c
PTVI	3.02 ± 0.01 ^b	3.72 ± 0.04 ^b
PT (V+Ct)I	3.31 ± 0.05 ^a	5.56 ± 0.01 ^a

Consequently, the leaves appear to begin a state of yellowing that would lead to their senescence. The results obtained in these experiments show that chlorophyll a is the key pigment to evaluate the green state of leaves. It seems to provide a good appreciation of the photosynthetic activity of leaves and would better express the physiological state of the plants [56,57]. Therefore, chl a/chl b and chl t/car could be regarded as indicators of plantain protection against black leaf streak disease (BLSD). Thus, vacciplant+calliete have a synergistic action on the protection of plantain against BLSD.

4. CONCLUSION

In the context of agriculture that is more respectful of the environment and human health, this study has been carried out in order to find new strategies for the protection of plants by exploiting their natural defense capacities. This research was conducted on French 2 plantain cultivar. To this end, vacciplant and calliete were applied as co-spray to the leaves and the plants were subsequently infected in vivo with virulent inoculum of *Mycosphaerella fijiensis*. The action of this co-treatment was compared to the action alone of vacciplant, the results of which were reported to be better for calliete. Vacciplant-treated and inoculated plants and co-treated and inoculated plants synthesized more total phenolic compounds than control plants. However, phenolic compounds produced by co-treated plants (191.10 mg/g FL) were greater than those treated with vacciplant alone (142 mg/g FL). Also, U-HPLC analysis revealed the *de novo* synthesis of four new compounds (phytoalexins) compared to the control (PNTNI) of eight compounds (constituent phenols or phytoanticipins). This study shows that the vacciplant+ calliete has a synergistic effect on the activation of the natural defenses in plantains against black line disease. Moreover, co-treatment has made it possible to induce *trans*-resveratrol, a stilbene with proven antifungal action. In addition, the level of synthesis of the constituent and induced phenolic compounds is

greater than in the vaccine alone. In addition, vacciplant+calliete allowed the plants to acquire a higher leaf greenery than those treated with vacciplant alone. These results show that vacciplant+ calliete co-treatment is synergistic and protects plantain against *M. fijiensis*-induced black leaf streak disease.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Lassois L, Busogoro JP, Jijakli H. La banane : de son origine à sa commercialisation. Biotechnology, Agronomy, Society and Environment. 2009;13(4): 575-586.
- Kwa M, Temple L. Le Bananier Plantain : Enjeux Socio-économiques et Techniques. Quæ CTA Presses Agronomiques de Gembloux, Belgique. 2019 :199.
- Soler A, N'da Adupo A. Réduction des pertes post-récolte du plantain. Projet PNUD/FAO IVC/87/003. Rapport final. Abidjan, Côte d'Ivoire: CIRADIRFA.1991 ;57.
- Traoré S, Kobenan K, Kouassi KS, Gnonhouri G. Systèmes de culture du bananier plantain et méthodes de lutte contre les parasites et ravageurs en milieu paysan en Côte d'Ivoire. Journal of Applied Biosciences. 2009;19:1094-1101.
- Silue M. Analyse de la demande des producteurs pour une offre de formation adaptée sur la banane plantain. 2021. Accessed on 13/3/2022. Available :https://agritrop.cirad.fr/599201/3/FABA_rapport%20Cote%20ivoire_besoins%20en%20formation_SMehin_2021.pdf.
- Daniells J, Hermanto C, Sutanto A, Jumjunidang, Edison W, O'Neill V, Sinohin A, Molina P, Taylor H. Incidence and distribution of fusarium wilt disease of banana in Indonesia. Acta Horticulturae. 2009;897: 313-322.

7. Lassoudière A. Le bananier et sa culture. Editions Quae, Paris, France. 2007;383 .
8. Stover R, Simmonds N. Bananas. John Wiley & Sons, Inc. New York, USA. 1987:468.
9. Konan Y, Kouassi K, Kouakou K, Koffi E, Kouassi K, Sékou D, Koné M., Kouakou T. Effect of Methyl jasmonate on phytoalexins biosynthesis and induced disease resistance to *Fusarium oxysporum* f. sp. *vasinfectum* in cotton. International Journal of Agronomy. 2014;1: 1-11.
10. N'goran KD. Régulation de la biosynthèse des polyphénols par des éliciteurs oligosaccharidiques et mycéliens d'origine fongique: effet sur la tolérance du cotonnier à la fusariose. Thèse de l'Université Nangui Abrogoua, Abidjan, Côte d'Ivoire. 2019 :197.
11. Hidalgo W, Chandran NJ, Menezes RC, Otalvaro F, Schneider B. Phenylphenalenones protect banana plants from infection by *Mycosphaerella fijiensis* and are deactivated by metabolic conversion. Plant, Cell and Environment. 2016; 39:492–513.
12. Abad P, Favery B. L'arsenal immunitaire des plantes. Pour la Science. 2012; 77:24-31.
13. Amari LD, Dick E, Koné D, Traoré S, Abo K, Kobénan K. Response of various banana genotypes to juglone toxic effect. Agronomie Africaine. 2011;23(3): 247-258.
14. Banerjee N, Delanghe E. A tissue-culture technique for rapid clonal propagation and storage under minimal growth-conditions of *Musa* (banana and plantain). Plant Cell Reports. 1985;4(6): 351-354.
15. Pinkerton F, Strobel G. Serinol as an activator of toxin production in attenuated cultures of *Helminthosporium sacchari*. PNAS. 1976;73(11):4007-4011.
16. Amari LDGE. Stratégies d'évaluation et de gestion par stimulation des défenses naturelles des bananiers à l'infection de la maladie des raies noires causée par *Mycosphaerella fijiensis* Morelet en Côte d'Ivoire. Thèse de l'Université Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire, 2012;237.
17. Belhadj A. Stimulation des défenses naturelles de la vigne par le méthyle jasmonate : impact sur la biosynthèse des polyphénols et sur la résistance aux champignons. Thèse de l'Université de Bordeaux 2, France. 2005:188.
18. Kouakou TH. Embryogenèse somatique chez le cotonnier: variation des composés phénoliques au cours de la callogenèse et de la culture des suspensions cellulaires. Thèse d'Etat, de l'Université Abobo-Adjamé, Côte d'Ivoire. 2009:137.
19. Siriwoharn T, Ronald E, Chad E, Chifford B. Influence of cultivar, maturity, and sampling on blackberry (*Rubus* L. Hybrids) anthocyanins, polyphenolics, and antioxidant properties. Journal of Agricultural and Food Chemistry. 2004;56(26):8021-8032.
20. Verdu C. Cartographie génétique des composés phénoliques de la pomme. Thèse de l'Université d'Angers, France. 2013 ;213.
21. Cruz-Cruz CA, Ramírez-Tec G, García-Sosa K, Escalante-Erosa F, Hill L, Osbourn AE, Peña-Rodríguez LM. Phytoanticipins from banana (*Musa acuminata* cv. Grande Naine) plants, with antifungal activity against *Mycosphaerella fijiensis*, the causal agent of black Sigatoka. European Journal of Plant Pathology. 2010;126:459–463.
22. Lichtenthaler H, Buschmann C. Chlorophylls and carotenoids: measurement and characterization by UV-VIS. In: Wrolsta et al. Current Protocols in Food Analytical Chemistry. 2001:431-438.
23. Lichtenthaler H. Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. Methods in Enzymology. 1987;148:350-382.
24. N'cho A, Yapou S, N'goran A, Kouakou T. Effect of the exogenous application of methyl and ethephon jasmonate on the accumulation of phenolic compounds and efficacy against fusarium wilt in cotton. Chemical Science International Journal. 2018;25(2):1-16.
25. Lattanzio V, Lattanzio V, Cardinali A. Role of phenolics in the resistance mechanisms of plants against fungal pathogens and insects. Advances in Research. 2006;661:23-67.
26. Yamaner O, Erdag B, Gokbulut C. Stimulation of the production of hypericins in *in vitro* seedlings of *Hy [2] Pericum adenotrichum* by some biotic elicitors. Turkish Journal of Botanic. 2013;37:153-159.
27. Konan YKF. Stimulation des défenses naturelles du cotonnier par le méthyle

- jasmonate et l'éthéphon : Effet sur la biosynthèse des composés phénoliques et sur la résistance à *Fusarium oxysporum* f. sp. *vasinfectum*, agent causal de la fusariose. Thèse de l'Université Nangui Abrogoua, Abidjan, Côte d'Ivoire. 2015 :207.
28. Beckers GJ, Spoel SH. Fine-Tuning Plant Defence Signalling: Salicylate versus Jasmonate. *Plant Biology*. 2006;8(1) :1-10.
 29. Smart CD, Myers KL, Restrepo S, Martin GB, Fry WE. Partial resistance of tomato to *Phytophthora infestans* is not dependent upon ethylene, jasmonic acid or salicylic acid signaling pathways. *Molecular Plant Microbe Interactions*. 2003;16:141-148.
 30. Jourdan E, Marc C, Phillipe T. Caractéristiques moléculaires de l'immunité des plantes induite par les rhizobactéries non pathogènes. *Biotechnologie, Agronomie, Société et Environnement*. 2008;12(4):437-449.
 31. Malamy J, Carr JP, Klessig DF, Raskin I. Salicylic acid: a likely endogenous signal in the resistance response of tobacco to viral infection. *Science*. 1990;50:1002-1004.
 32. Walters MB, Kruger EL, Reich PB. Relative growth rate in relation to physiological and morphological traits for northern hardwood tree seedlings: species, light environment and ontogenetic considerations. *Oecologia*. 1993;96:219-231.
 33. Lambert C. Etude du rôle des stilbènes dans les défenses de la vigne malade contre les maladies du bois. Thèse de l'Université de Bordeaux 2, France. 2011 :179.
 34. Bellow S. Etude des composés phénoliques impliqués dans la réponse des feuilles de vigne au mildiou. Thèse de l'Université de Paris sud, France. 2012 :134.
 35. Pieterse C, Dieuwertje V, Christos Z, Antonio L, Saskia W. Hormonal modulation of plant immunity. *Annual Review of Cell and Developmental Biology*. 2012 ;28:489-521.
 36. Kazunori S, Tomoyuki O, Yumi K, Yoichi N, Shigenori O. Effect of repeated harvesting on the content of caffeic acid and seven species of caffeoylquinic acids in sweet potato leaves. *Bioscience, Biotechnology and Biochemistry*. 2015;79(8):1308-1314.
 37. Yin Z, Sadok A, Sailem H, McCarthy A, Xia X, Li, F, Garcia MA, Evans L, Barr AR, Perrimon N, Marshall CJ, Wong ST, Bakal C. A screen for morphological complexity identifies regulators of switch-like transitions between discrete cell shapes. *Nature Cell Biology*. 2013;15(7): 860-871.
 38. Dufour MC, Lambert C, Bouscaut J, Méryllon JM, Corio-Costet MF. Benzothiadiazole-primed defence responses and enhanced differential expression of defence genes in *Vitis vinifera* infected with biotrophic pathogens *Erysiphe necator* and *Plasmopara viticola*. *Plant. Pathology*. 2013;62:370-382.
 39. N'goran AR, Kouakou TH, Konan FKH, Camara B, Kouassi NK, Kone D. Effet de la fraction oligosaccharidique de *Fusarium oxysporum* f. sp. *vasinfectum*, sur la protection du cotonnier contre la fusariose. *Agronomie Africaine*. 2016; 28(3):1-10.
 40. Belhadj A, Saigne C, Telef N, Cluzet S, Bouscaut J, Corio-Costet MF. Methyl jasmonate induces defense responses in grapevine and triggers protection against *Erysiphe necator*. *Journal of Agricultural and Food Chemistry*. 2006;54(24):911-925.
 41. Kouakou TH, Koné M, Koné D, Kouadio YJ, Amani NG, Teguo WP, Decendit A, Merillon JM. Trans-resvératrol as phenolic indicator of somatic embryogenesis induction in cotton cell suspensions. *African Journal of Biochemistry Research*. 2008; 2(1):15-23.
 42. Kouakou TH, Due EA, Kouadio NEJP, Niamke S, Kouadio YJ, Waffo TP, Decendit A, Merillon JM. Purification and biochemical characterization of polyphenol oxidases from embryogenic and nonembryogenic cotton cells. *Applied Biochemistry and Biotechnology*. 2009a;158:285-301.
 43. Kouakou TH, Due EA, Kouadio NEJP, Niamke S, Kouadio YJ, Waffo TP, Decendit A, Merillon JM. Purification and characterization of cell suspensions peroxidase from cotton *Applied Biochemistry and Biotechnology*. 2009b;157:575-592.
 44. Faurie B, Cluzet S, Corio-costet MF, Merillon JM. Methyl jasmonate/ethephon cotreatment synergically induces stilbène production in *Vitis vinifera* cell suspension but fails to trigger resistance to *Erysiphe necator*. *Journal International Sciences Vigne Vin*. 2009;43(2): 99-110.
 45. Ahuja I, Kissen R, Bones AM. Phytoalexins in defense against pathogens. *Trends in Plant Sciences*. 2012; 17:73-90.

46. N'goran AR. Stimulation des défenses naturelles du cotonnier par des éliciteurs oligosaccharidiques extraits des suspensions de *Fusarium oxysporum* f. sp. *vasinfectum*, intérêt phytopathologique des composés phénoliques. Thèse de doctorat de l'Université Félix Houphouët Boigny, Abidjan, Côte d'Ivoire. 2015 :169.
47. Bi Z, Zhao Y, Morrell JJ, Lei Y, Yan L. The antifungal mechanism of konjac flying powder extract and its active compounds against wood decay fungi. *Industrial Crops and Products*. 2021 ;164:113406.
48. Clerivet A, Alami I, Breton F, Garcia D, Sanier C. Les composés phénoliques et la résistance des plantes aux agents pathogènes. *Acta Botanica Gallica*. 1996;143:531-538.
49. N'cho XE. Élicitation du bananier par le méthyle jasmonate et l'acide salicylique : impact sur les composés phénoliques, efficacité sur *Mycosphaerella fijiensis* responsable de la maladie des raies noires. Thèse de l'Université Nangui Abrogoua, Abidjan, Côte d'Ivoire. 2017:198.
50. Wang F, Feng G, Chen K. Defense responses of harvested tomato fruit to burdock fructooligosaccharide, a novel potential elicitor. *Postharvest Biology and Technology*. 2009;52(1):110-116.
51. Porra RJ. Recent progress in porphyrin and chlorophyll biosynthesis. *Photochemistry and Photobiology*. 1997; 65:492-516.
52. Folly P. Catabolisme de la chlorophylle b Structures, mécanismes et synthèses. Thèse de de l'Université de Fribourg, Suisse. 2000 :192.
53. Djapic N. Chlorophyll catabolism in *Prunus serrulata* autumnal leaves. *Facta Universitatis Series: Physics, Chemistry and Technology*. 2012;10(1): 21-26.
54. Gamon JA, Serrano L, Surfus JS. The photochemical reflectance index: an optical indicator of photosynthetic radiation use efficiency across species, functional types and nutrient levels. *Oecologia*. 1997;112:492-501.
55. Barry KM, Newnham GJ. Anthocyanin and epicuticular waxes affect quantification of chlorophyll and carotenoid pigments in eucalyptus foliage with the radioactive transfer model Prospect 5. *Geospatial Science Research*. 2012;1328:1-7.
56. Zarco-Tejada PJ, Miller JR, Mohammed GH, Noland TL, Sampson PH. Vegetation stress detection through chlorophyll a + b estimation and fluorescence effects on hyperspectral imagery. *Journal of Environmental Quality*. 2002;31:1433-1441.
57. Garrity SR, Eitel JUH, Vierling LA. Disentangling the relationships between plant pigments and the photochemical reflectance index reveals a new approach for remote estimation of carotenoid content. *Remote Sensing of Environment*. 2011;115:628-635.

© 2022 Lucien et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86648>