

## Annual Research & Review in Biology

32(4): 1-9, 2019; Article no.ARRB.46992  
ISSN: 2347-565X, NLM ID: 101632869

# Comparative Studies on the Effect of Fermentation on the Nutritional Compositions and Anti-nutritional Levels of *Glycine max* Fermented Products: Tempeh and Soy-Iru

T. R. Omodara<sup>1\*</sup> and E. Y. Aderibigbe<sup>1</sup>

<sup>1</sup>Department of Microbiology, Ekiti State University, P.M.B. 5363, Ado-Ekiti, Nigeria.

### Authors' contributions

This work was carried out in collaboration between both authors. Author TRO performed the bench work, statistical analysis, wrote the first draft of the manuscript and managed the literature. Author EYA initiated the study, provided the protocol, managed the analysis of the study and vetted the final manuscript. Both authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/ARRB/2019/v32i430094

#### Editor(s):

- (1) Paola Angelini, Department of Applied Biology, University of Perugia, Perugia, Italy.
- (2) Prof. George Perry, Dean and Professor of Biology, University of Texas at San Antonio, USA.

#### Reviewers:

- (1) Takeshi Nagai, Graduate School of Yamagata University, Japan.
  - (2) Zakaria Fouad Fawzy Hassan, National Research Centre, Egypt.
- Complete Peer review History: <http://www.sdiarticle3.com/review-history/46992>

Original Research Article

Received 10 December 2018  
Accepted 17 February 2019  
Published 06 August 2019

## ABSTRACT

**Aims:** A comparative study of fungi and bacteria fermentation of soybean (*Glycine max*) was carried out to determine the effect of fermentation on the nutritional composition of their fermented products: tempeh and 'soy-iru'.

**Study Design:** The experiment was carried out in the Department of Microbiology, Ekiti State University, Ado Ekiti, Nigeria, between August, 2017 and July 2018.

**Methodology:** Soybean was processed into 'soy-iru' (bacterial fermentation) and tempeh (fungal fermentation) and the microbial load, physico-chemical properties, proximate composition, levels of anti-nutritional components (trypsin inhibitor and phytic acid), anti-oxidants (total phenol, total flavonoid and DPPH), in-vitro protein digestibility and vitamins (A, B, C, D, and E) were analyzed.

\*Corresponding author: E-mail: omodaratolani@yahoo.com;

**Results:** The microbial load, pH increased progressively during fermentation, while there was a decrease in the titratable acidity (TTA) of the two products. The protein(%), ash(%) and fat(%) contents of the *Glycine max* cotyledons increased from 29.56, 1.86 and 24.36 in unfermented substrate to 33.61, 2.21 and 26.90, respectively, after 24hrs of fermentation to produce tempeh. However, there was a reduction in the crude fibre(%) and carbohydrate(%) content from 2.94 and 41.29 in unfermented substrate to 2.53 and 32.57, respectively, after 24hrs of fermentation. Similar trends were observed during the production of 'soy-iru', however the change in proximate composition was not as significant as observed in tempeh. There was significant decrease in the trypsin inhibitor and phytic acid levels of the two products. The levels of anti-oxidants, vitamins B, D, E and protein digestibility increased significantly, in both bacterial and fungal-fermented products.

**Conclusion:** This research has therefore shown that fungal fermentation of *Glycine max* seeds into tempeh may be a better alternative to 'soy-iru' which was obtained from bacterial fermentation, because of the significant lower level anti-nutritional factors in the former.

**Keywords:** *Glycine max*; vitamins; tempeh; soy-iru; anti-nutritional factors.

## 1. INTRODUCTION

Soybean (*Glycine max*) is a plant legume, known for more than 3000 years in Southeastern Asia [1]. Soybean is one of the widely consumed foods in the world due to its high nutritional value and low cost [2]. It is a legume that has high level of protein, appreciable amount of minerals, vitamins and fibres, some amount of antioxidants, small amounts of saturated fat and absence of cholesterol [2].

Some of the health benefits of soybean include: improved metabolic activities, healthy weight gain, prevention of cancer, boost heart health, relieves menopausal symptoms, boost digestion and improve bone health. However, raw soybean is toxic to non-ruminants due to high concentration of anti-nutritional factors such as trypsin inhibitors and high level of phytic acids. [3]. Most of these anti-nutritional factors present in the raw seeds chelate some important vitamins and minerals, thereby preventing their absorption into the body. Due to the high level of anti-nutritional factors, processing is required before the seeds can be consumed by non-ruminant, since the goal of eating is to get adequate amount of nutrients in the diet [4]. Fermentation is one of the processing methods that can be employed in the processing of soybean into soy-iru. Bacterial fermentation (using *Bacillus subtilis* strains) leads to production of 'soy-iru', natto, thua-nao; while fungal fermentation (using *Rhizopus oligosporus*) leads to production of tempeh [5]. This research aims at comparing the bacterial fermented product ('soy-iru') of soybean, with the fungal fermented product (tempeh), on the bases of

nutritional factors, anti-nutritional factors and anti-oxidant levels.

## 2. MATERIALS AND METHODS

### 2.1 Sources of Materials

The *Glycine max* seeds were purchased from Oja Oba in Ado-Ekiti. The pure cultures of *Bacillus subtilis* strains and *Rhizopus oligosporus* were obtained from the stock cultures kept in the Laboratory of Department of Microbiology, Ekiti State University, Ado-Ekiti, Nigeria.

### 2.2 Processing of the Seeds

The method described by [4] on the production of 'soy-iru' from soybean (*Glycine max*) seeds was adopted. Five hundred grams (500 g) of soybean seeds were sorted washed and boiled for 2h. The boiled seeds were dehulled to remove the seed coat, washed and boiled again for 1 hour. The water was drained off and the beans were fermented in an incubator at 35°C for 36 h. Samples were taken at every 12 h and analyzed for microbial load, physico-chemical properties, proximate, anti-nutritional content, antioxidant level, vitamins content and protein digestibility.

### 2.3 Preparation of Spores' Suspension for 'tempeh' Production

The procedure described by [5] was adopted to prepare spore suspension. Five grams (5 g) of Malt Extract Agar (MEA, Oxoid) was weighed and dissolved in 100 ml of distilled water in a 250 ml conical flask. The medium was homogenized

and sterilized in an autoclave at 121°C for 15 minutes. One gram of bacteriological peptone (Lab M) was weighed and dissolved in 100 ml of distilled water in a 250 ml conical flask. This was also sterilized by autoclaving. The sterile MEA was poured into sterile plates and allowed to solidify. One gram of *Rhizopus oligosporus* NRRL 2710 powder was added aseptically into 5ml sterile peptone water in a 100 ml conical flask and it was mixed together to disperse the powdered inoculum. One millilitre (1 ml) was inoculated into the MEA plate. The agar plates were inverted and incubated at 30°C for 72 h. After incubation, the spores were harvested by pouring 5 ml sterile peptone water into each of the sporulated culture in the Petri dishes and scrapped, using wire loop. The harvested culture was filtered through sterile non-absorbent cotton wool into a sterile conical flask to obtain the spores' suspension.

#### **2.4 Laboratory Production of 'tempeh' from Soybean (*Glycine max*) Seeds**

'Tempeh' was prepared by fermenting soybean according to the procedure of [5]. The soybeans (*Glycine max*) were washed and boiled partially for 30 mins. The soybeans were dehulled, cleaned and soaked in clean water overnight. The soaked soybeans were then boiled for 45 mins. The moist cotyledons were drained properly air-dried and cooled for 20-30 minutes after which they were inoculated with spores' suspension of *Rhizopus oligosporus* NRRL 2710 with ratio 1:50 (v/w). The cotyledons were lightly packed into sterile perforated baking tins covered with perforated aluminum foil paper and incubated for 24 h at 35°C. Samples were taken at every 12 h and analyzed for microbial load, physico-chemical properties, proximate, anti-nutritional content, antioxidant level, vitamins content and protein digestibility.

#### **2.5 Microbiological Analysis**

The microbial load (viable counts) was determined using serial dilution and plating technique on nutrient agar (NA) plates. The bacterial isolates were partially characterized on the bases of cultural, morphological and biochemical properties [6].

#### **2.6 Determination of pH**

Five grams (5 g) of each sample was homogenized and mixed with 100 ml of distilled

water. The pH of each homogenate was determined with a Pye Unicam pH meter (Model PW9409). The determination was carried out in triplicates.

#### **2.7 Total Titratable Acidity Determination**

The suspension from the pH determination was filtered and 20 ml of the filtrate was titrated against 0.1M NaOH using 1 drop of phenolphthalein as indicator [7].

#### **2.8 Moisture Content Determination**

Five grams (5 g) of each sample was weighed separately into pre-weighed aluminum foil. The foil paper and its content was put in oven at 80°C overnight and weighed intermittently until a constant weight was achieved. The new weight was subtracted from the weight of the wet sample. The percentage moisture content was calculated [8].

#### **2.9 Proximate Analysis**

The proximate compositions of the fermented and unfermented samples were determined using standard procedures of [8]. The parameters determined were protein, ash, crude fibre, fat and carbohydrate.

#### **2.10 Determination of Anti-nutritional Factors**

##### **2.10.1 Phytic acid**

The method of [9] was employed in the determination of phytic acid. Four grams (4 g) of finely ground sample was soaked in 1 L of 2% HCl inside conical flask for 3 h and was filtered. Five milliliters (5 ml) of 0.03% NH<sub>4</sub>SCN was added as indicator and 50 ml of distilled water also added. This was titrated against ferric chloride solution which contained 0.05 mg of iron (Fe) per ml of FeCl<sub>3</sub>. The iron equivalent was obtained and the phytate content in mg/100 mg of dried sample was calculated.

##### **2.10.2 Trypsin inhibitor**

The trypsin inhibitor activity (TIA) in the sample was determined according to the method of [10]. The digest contained 1.0 g of the sample, 40 µg of trypsin and 2 mg of Nalpha-benzoyl-

DL-Arginine-P-nitroanilidehydrochloride. The absorbance was read at 410 nm.

## 2.11 Determination of Anti-Oxidants

### 2.11.1 Total phenol

The total phenol contents of the samples were determined using the method reported by [11], while total flavonoids content and 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical-scavenging ability of the samples were determined by the method of [12] and [13], respectively.

## 2.12 Determination of Vitamins

Vitamin A was determined by the method of [14]; vitamin B by the method of [15] vitamin C by the method of [16], while vitamins D and E were determined by the methods of Pearson [14].

## 2.13 Determination of Multi-Enzyme *In vitro* Protein Digestibility

The method of Singh and Krikorian [17] was adopted in the determination of multi-enzyme *in vitro* protein digestibility of the samples, using procaine pancreatic trypsin as enzyme. The absorbance was read at 700 nm against reagent blank. The standard calibration (STD) curve was prepared using 100 µg/ml of Bovine Serum Albumen (BSA).

## 3. RESULTS

Fig. 1 shows the microbial load of the samples during fermentation of *Glycine max* to 'tempeh' and 'soy-iru'. The microbial load increased progressively at different periods of fermentation, from 4.55 log CFU/g to 8.74 log CFU/g ('tempeh') and 4.75 to 7.67 log CFU/g ('soy-iru'), respectively. The pH of the substrate increased significantly during the fermentation (Fig. 2) from 5.50 to 6.94 (tempeh) and 5.50 to 8.079 ('soy-iru'). The total titratable acidity (TTA) (Fig. 3) of *Glycine max* reduced from  $3.09 \times 10^{-2}$ N to  $2.17 \times 10^{-2}$ N (tempeh) and from  $2.57 \times 10^{-2}$ N to  $1.10 \times 10^{-2}$  ('soy-iru'). As shown in Fig. 4, the moisture content of the substrate decreased from 20.3% to 16.53% in tempeh; but increased from 45.33% to 59% in 'soy-iru'.

The proximate compositions of 'tempeh' and 'soy-iru' during fermentation are shown in Table 1. The protein content of the *Glycine max* cotyledons increased from 29.56% to 33.61%


during fermentation of tempeh. There were also increases in the ash and fat contents. However, the crude fibre and carbohydrate content decreased from 2.94% to 2.53% and 41.29% to 32.57%, respectively. Similar trends in the values of the parameters assessed were observed during 'soy-iru' fermentations.

Table 2 shows the anti-nutritional factors and the anti-oxidants level of the fermenting substrate and products. The trypsin inhibitor level decreased significantly from 55.84 mg/g to 44.33 mg/g (tempeh) and from 64.35 mg/g to 45.02 mg/g ('soy-iru'), respectively. Similarly, phytic acid content decreased significantly from 38.45 mg/g to 8.43 mg/g and 55.76 mg/g to 9.89 mg/g in 'tempeh' and 'soy iru', respectively, after fermentation. There was significant increase in the anti-oxidants levels of the substrate during fermentation. The total flavonoids contents increased from 0.04 mg/g to 0.15 mg/g in 'tempeh' and 0.03 mg/g to 0.21 mg/g in 'soy-iru'. A similar trend was observed in the contents of total phenol and diphenyl-picrylhydrazyl (DPPH) radical scavengers during the fermentation of both the tempeh and 'soy-iru'. The vitamins and the *in vitro* protein digestibility of the fermenting substrate and fermented products during fermentation of *Glycine max* to tempeh and 'soy-iru' are presented in Table 3.

The vitamins A and C contents reduced during the fermentation processes. Vitamin A contents reduced from 10.73 mg/g to 3.28 mg/g ('tempeh') and 14.95 mg/g to 4.17mg/g ('soy iru'); while vitamin C contents reduced from 0.18mg/g to 0.10 mg/g in 'tempeh' and 0.46 mg/g to 0.06mg/g in 'soy-iru'. However, vitamins B, D and E increased significantly during the fermentation. There was a significant increase in the *in vitro* protein digestibility from 28.78% to 62.02% in tempeh; similar result was observed for 'soy-iru'.


## 4. DISCUSSION

The steady increase in microbial load during the fermentation might be due to availability of nutrients released from the cotyledons by the action of fermentation and the utilization of these nutrients by the fermenting organisms for their metabolic activities. This is in agreement with the previous result gotten by Omodara and Aderibigbe [18] when working on 'iru'. The increase in the protein, ash, fat and anti-oxidants might be attributed to secretion of hydrolytic enzymes by the fermenting organisms [19]. The



**Fig. 1. Microbial load (log CFU/g) of 'tempeh' and 'soy-iru' during fermentation of *Glycine max* seeds**

Key: UF= unfermented substrate, T1= sample at 12 h of fermentation, T2 = sample at 24 h of fermentation, T3= sample at 36 h of fermentation



**Fig. 2. pH of 'tempeh' and 'soy-iru' during fermentation *Glycine max* seeds**

Key: UFS= unfermented substrate, T12= sample at 12h of fermentation, T24 = sample at 24h of fermentation, T36= sample at 36h of fermentation




Fig. 3. Total titratable acidity (TTA) 'tempeh' and 'soy-iru' during fermentation of *Glycine max* seeds

Key: UFS= unfermented substrate, T12= sample at 12h of fermentation, T24 = sample at 24h of fermentation, T36= sample at 36h of fermentation




Fig. 4. Moisture content (%) of 'tempeh' and 'soy-iru' during fermentation of *Glycine max* seeds

Key: UFS= unfermented substrate, T12= sample at 12 h of fermentation, T24 = sample at 24 h of fermentation, T36= sample at 36 h of fermentation

**Table 1. Proximate composition (%) of ‘tempeh’ and soy-iru during fermentation of *Glycine max* seeds**

Samples	Proximate Composition (%)									
	Protein		Ash		Fibre		Fat		Carbohydrate	
	Tempeh	Soy-iru	Tempeh	Soy-iru	Tempeh	Soy-iru	Tempeh	Soy-iru	Tempeh	Soy-iru
UFS	29.56 <sup>c</sup> ±0.48	24.51 <sup>d</sup> ±0.01	1.86 <sup>d</sup> ±0.01	0.98 <sup>d</sup> ±0.01	2.94 <sup>a</sup> ±0.01	3.86 <sup>a</sup> ±0.10	24.36 <sup>c</sup> ±0.02	20.06 <sup>b</sup> ±0.10	41.29 <sup>a</sup> ±0.44	50.12 <sup>a</sup> ±1.17
T12	31.79 <sup>b</sup> ±0.23	26.97 <sup>c</sup> ±0.02	1.98 <sup>c</sup> ±0.01	1.34 <sup>c</sup> ±0.00	2.84 <sup>b</sup> ±0.01	3.78 <sup>a</sup> ±0.17	25.24 <sup>b</sup> ±0.09	21.97 <sup>c</sup> ±0.01	38.16 <sup>b</sup> ±0.31	46.26 <sup>b</sup> ±0.47
T24	31.24 <sup>b</sup> ±0.45	29.00 <sup>b</sup> ±0.58	2.13 <sup>b</sup> ±0.02	1.53 <sup>b</sup> ±0.05	2.63 <sup>c</sup> ±0.00	3.61 <sup>b</sup> ±0.06	26.85 <sup>a</sup> ±0.02	23.60 <sup>b</sup> ±0.13	34.79 <sup>c</sup> ±0.50	42.36 <sup>c</sup> ±0.00
T36	33.61 <sup>a</sup> ±0.00	31.27 <sup>a</sup> ±0.06	2.21 <sup>a</sup> ±0.02	1.74 <sup>a</sup> ±0.04	2.53 <sup>d</sup> ±0.02	3.46 <sup>b</sup> ±0.06	26.90 <sup>a</sup> ±0.10	32.57 <sup>a</sup> ±0.04	32.57 <sup>d</sup> ±0.04	38.73 <sup>d</sup> ±0.21

Key: UFS= unfermented substrate, T12= sample at 12 h of fermentation, T24 = sample at 24 h of fermentation, T36= sample at 36 h of fermentation.  
 Values that have same superscript in a column are not significantly different at P = 0.05

**Table 2. Anti-nutritional factors (mg/g) and antioxidant levels (mg/g) of ‘tempeh’ and soy-iru during fermentation of *Glycine max* seeds**

Samples	Antinutritional factors (mg/g)				Antioxidants (mg/g)					
	Trypsin inhibitor		Phytic acid		Total phenol		Total flavonoids		Free radical scavengers	
	Tempeh	Soy-iru	Tempeh	Soy-iru	Tempeh	Soy-iru	Tempeh	Soy-iru	Tempeh	Soy-iru
UFS	55.84 <sup>a</sup> ±0.29	64.35 <sup>a</sup> ±0.28	38.45 <sup>a</sup> ±0.48	55.76 <sup>a</sup> ±0.47	0.23 <sup>d</sup> ±0.01	0.46 <sup>d</sup> ±0.00	0.04 <sup>d</sup> ±0.00	0.03 <sup>c</sup> ±0.03	66.45 <sup>d</sup> ±0.65	68.82 <sup>d</sup> ±1.00
T12	51.11 <sup>b</sup> ±0.07	52.43 <sup>b</sup> ±0.00	19.23 <sup>b</sup> ±0.95	29.36 <sup>b</sup> ±0.47	0.43 <sup>c</sup> ±0.00	0.49 <sup>c</sup> ±0.00	0.07 <sup>c</sup> ±0.00	0.08 <sup>b</sup> ±0.00	73.33 <sup>c</sup> ±0.75	74.47 <sup>c</sup> ±1.00
T24	46.29 <sup>c</sup> ±0.00	49.36 <sup>c</sup> ±0.00	13.71 <sup>c</sup> ±0.00	17.30 <sup>c</sup> ±0.00	0.49 <sup>b</sup> ±0.00	0.56 <sup>b</sup> ±0.01	0.09 <sup>b</sup> ±0.00	0.10 <sup>b</sup> ±0.01	75.91 <sup>b</sup> ±0.38	85.16 <sup>b</sup> ±1.30
T36	44.33 <sup>d</sup> ±0.14	45.02 <sup>d</sup> ±0.00	8.43 <sup>d</sup> ±0.00	9.89 <sup>d</sup> ±0.00	0.62 <sup>a</sup> ±0.00	0.63 <sup>a</sup> ±0.01	0.15 <sup>a</sup> ±0.01	0.21 <sup>a</sup> ±0.02	86.45 <sup>a</sup> ±0.00	88.17 <sup>a</sup> ±1.34

Key: UFS= unfermented substrate, T12= sample at 12h of fermentation, T24 = sample at 24h of fermentation, T36= sample at 36h of fermentation.  
 Values that have superscript in a column are not significantly different at P = 0.05

**Table 3. Vitamins (mg/g) and protein digestibility levels (%) of tempeh and ‘soy-iru’ during fermentation of *Glycine max* seeds**

Samples	Vitamins (mg/g)										Protein digestibility (%)	
	A		B		C		D		E		Tempeh	Soy-iru
	Tempeh x10 <sup>2</sup>	Soy-iru x10 <sup>2</sup>	Tempeh	Soy-iru	Tempeh	Soy-iru	Tempeh	Soy-iru	Tempeh	Soy-iru		
UFS	10.73 <sup>a</sup> ±0.69	14.95 <sup>a</sup> ±0.23	0.26 <sup>d</sup> ±0.00	0.15 <sup>d</sup> ±0.00	0.18 <sup>a</sup> ±0.02	0.46 <sup>a</sup> ±0.02	0.18 <sup>d</sup> ±0.00	0.41 <sup>d</sup> ±0.00	0.28 <sup>d</sup> ±0.03	0.46 <sup>d</sup> ±0.00	28.78 <sup>d</sup> ±0.13	32.96 <sup>d</sup> ±0.21
T12	5.81 <sup>b</sup> ±1.10	8.62 <sup>b</sup> ±0.04	0.41 <sup>c</sup> ±0.00	0.41 <sup>c</sup> ±0.00	0.14 <sup>b</sup> ±0.01	0.23 <sup>b</sup> ±0.00	0.46 <sup>c</sup> ±0.02	0.49 <sup>c</sup> ±0.06	0.58 <sup>c</sup> ±0.00	0.58 <sup>c</sup> ±0.00	46.10 <sup>c</sup> ±0.10	40.23 <sup>c</sup> ±0.20
T24	4.43 <sup>c</sup> ±0.00	5.34 <sup>c</sup> ±0.37	0.56 <sup>b</sup> ±0.00	0.56 <sup>b</sup> ±0.00	0.12 <sup>c</sup> ±0.00	0.13 <sup>c</sup> ±0.01	1.28 <sup>b</sup> ±0.02	1.40 <sup>b</sup> ±0.00	1.14 <sup>b</sup> ±0.03	1.14 <sup>b</sup> ±0.03	54.90 <sup>b</sup> ±0.10	47.70 <sup>b</sup> ±0.65
T36	3.28 <sup>d</sup> ±0.00	4.17 <sup>d</sup> ±0.00	1.09 <sup>a</sup> ±0.21	1.09 <sup>a</sup> ±0.21	0.10 <sup>d</sup> ±0.01	0.06 <sup>d</sup> ±0.00	1.59 <sup>a</sup> ±0.00	1.84 <sup>a</sup> ±0.00	1.51 <sup>a</sup> ±0.00	1.51 <sup>a</sup> ±0.00	62.02 <sup>a</sup> ±0.02	57.13 <sup>a</sup> ±0.61

Key: UFS= unfermented substrate, T12= sample at 12h of fermentation, T24 = sample at 24h of fermentation, T36= sample at 36h of fermentation.  
 Values that have superscript in a column are not significantly different at P = 0.05

decrease in the level of phytic acid and trypsin inhibitor may be attributed to the metabolic activities of the fermenting organism. It may also be due to breaking down of these complexes by the enzymes produced the fermenting organisms [18]. The increase in the Vitamin B, D and E with increase in fermentation might be due to the release of this vitamin from their bond state by the activities of the fermenting organisms while the decrease in Vitamin A and C might be due to the metabolic activities of the fermenting organisms. It was found that fermentation had a significant increase in in- vitro protein digestibility of the two products. The microorganisms involved in the fermentation produce proteolytic enzymes which degrade complex proteins, hence increase in digestibility [20].

## 5. CONCLUSION

In conclusion fermentation was found to enhance of the nutritional qualities of *Glycine max* seeds when fermented into tempeh (using *Rhizopus oligosporus* NRRL 2710) and 'soy-iru' (using *Bacillus subtilis* 3A); as both have significant reduction in the anti-nutritional contents (phytic acid and tpsin inhibitor). However, tempeh may be better alternative to process the 'soybean' because of its lower anti-nutritional factors.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Lusas EW, Rias MN. Soy protein products: Processing and use. *J Nutr.* 1995;125:573-580.
2. Ciabotti S, Silva AB, Juhasz AP, Mendonça CD, Tavano OL, Mandarino JG, et al. Chemical composition, protein profile, and isoflavones content in soybean genotypes with different seed coat colors. *International Food Research Journal.* 2016;23(2):621-629.
3. Li HD, Zagorski J, Fournier MJ. Depletion of small nuclear RNA (snR128) disrupts production of 18S rRNA in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* 1910;3:1145-1152.
4. Fabiyi EB. Soyabean processing, utilization and health benefits. *Pakistan Journal of Nutrition.* 2006;5(5):453-457.
5. Aderibigbe EY, Adebayo CO. Fermentative utilization of soybean in Nigeria I. Laboratory production of 'tempeh'. *NISEB Journal.* 2003;2:31-35.
6. Olutiola PO, Famurewa O, Sonntag HG. Introduction to general microbiology: A practical approach. 2<sup>nd</sup> ed. Bolabay Publications, Ikeja, Nigeria; 2000.
7. Joslyn MA. Methods in food analysis. Academic Press, New York. 1970;49: 615.
8. AOAC. Official methods of analysis. 17<sup>th</sup> Edition Association of Official Analytical Chemist Washington D.C. 2005;5-10.
9. Wheeler EL, Ferrel RE. A method for phytic acid determination in wheat fractions. *Chem.* 1971;48:312-316.
10. Smith C, Megen WV, Twaalfhoven C, Hitcheock N. The determination of trypsin inhibitor levels in food stuffs. *J of Food Science and Agriculture.* 1980;3:341-350.
11. Singleton VL, Othorfor R, Lamuela-Raventos RM. Analysis of totalphenols and other oxidation substrates and antioxidants by means of folin gocattan reagent. *Methods in Enzymology.* 1999;152-178.
12. Meda A, Lamien CE, Romito M, Millogo J, Nacoulma OG. Determination of the total phenolic, total flavonoid and proline content in Burkinafaso honey, as well as their radical scavenging activity. *Food Chemistry.* 2005;91:571-577.
13. Gyamfi MA, Yonamine M, Aniya Y. Free radical scavenging action of medicinal herbs from Ghana: Thonnigfi sanguinea on experimentally induced liver injuries *General Pharmacology.* 1999;32:66-667.
14. Pearson D. Chemical analyses of foods. 7<sup>th</sup> ed. Churchill Livingstone, London. 1976;6-25
15. Okwo DE, Josiah C. Evaluation of chemical composition of two Nigerian medicinal plants. *African Journal of Biotechnology.* 2006;5(4):357-361.
16. Benderitter M, Mavpou V, Vergely C, Daltoz F, Briot F, Rochette L. Studies by electron paramagnetic resonance of the importance of iron in hydroxyl scavenging properties of ascorbic acid plasma: effects of iron chelators. *Fundamental and of Clinical Pharmacology.* 1998;12(5):510-516.
17. Singh M, Krikorian AD. Inhibition of trypsin activity *in-vitro* by phytate. *J. Agric. Food Chem.* 1982;30:799-800.



18. Omodara TR, Aderibigbe EY. Effects of the use of starter culture on the quality of fermented *Parkia biglobosa* seeds. Intl. J. of Biotechnol. 2013;3(4)33-40.
19. Oboh G, Rocha O. Polyphenols in red pepper [*Capsicum annuum* var. *aviculere* (Tel)] and their protective effect on some pre-oxidants lipids peroxidation in brain and liver, Eur. Food Res. Technol. 2007; 225:239-247
20. Aderibigbe EY, Odunfa SA, Schink B. Extracellular proteinases of *Bacillus* species isolated from fermented African Locust bean seeds, 'iru'. Food Microbiol. 1990;7:281-293.

© 2019 Omodara and Aderibigbe; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/46992>