



Tungiasis: Biology, Life Cycle, Epidemiology, Diagnosis, Prevention, and Treatment

Rhoda Nwalozie ^{a*} and Chinonye Oluchi Ezenwaka ^b

^a *Department of Medical Laboratory Science, Faculty of Science, Rivers State University, Port Harcourt, Nigeria.*

^b *Department of Biology, Faculty of Science, Federal University Otuoke, Bayelsa State, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106007>

Review Article

Received: 15/07/2023

Accepted: 17/09/2023

Published: 22/09/2023

ABSTRACT

Tungiasis, caused by the female sand flea *Tunga penetrans*, is a neglected tropical disease that affects impoverished communities in tropical and subtropical regions. This comprehensive review discusses the biology, life cycle, epidemiology, clinical presentation, diagnosis, prevention, and treatment of tungiasis. Tungiasis exhibits a patchy distribution, with higher prevalence rates reported in rural areas and areas with poor sanitation. Risk factors include poverty, inadequate housing conditions, and lack of access to proper footwear. The disease primarily affects vulnerable populations such as children and the elderly, leading to a significant socioeconomic burden. Clinical presentation of tungiasis involves characteristic lesions with embedded fleas, often leading to secondary infections, abscesses and ulcerations. Diagnosis is based on clinical examination and visualization of the flea within the skin lesion. Treatment involves the physical removal of embedded fleas, management of associated complications, wound care, and prevention of secondary infections. Prevention and control strategies are crucial in reducing the disease burden. Personal and environmental hygiene, including regular washing of feet and maintaining clean living

*Corresponding author: E-mail: rhoda.nwalozie@ust.edu.ng;

environments, play a vital role in prevention. Community education and health promotion campaigns raise awareness and promote behavior change. Vector control measures and integrated approaches combining multiple strategies are effective in reducing tungiasis prevalence. Although progress has been made in understanding and managing tungiasis, challenges persist. Further research is needed to improve diagnostic techniques, enhance treatment options, and address the long-term consequences of the disease. Community engagement, interdisciplinary collaborations, and sustainable interventions are essential in controlling tungiasis and improving the health and well-being of affected populations. By implementing comprehensive prevention and control strategies, raising awareness, and addressing the socioeconomic impact of tungiasis, we can strive towards reducing the burden of this neglected tropical disease and improving the lives of individuals and communities affected by tungiasis.

Keywords: *Tungiasis; Tunga penetrans; epidemiology; clinical presentation; diagnosis; prevention; treatment.*

1. INTRODUCTION

Tungiasis, caused by the parasitic flea *Tunga penetrans*, is an ectoparasitic disease that primarily affects resource-poor communities in tropical and subtropical regions [1]. It is classified as a neglected tropical disease (NTD), with high prevalence in certain areas and a significant impact on public health [2]. Tungiasis, also known as jigger infestation or sand flea disease, is characterized by the penetration and embedding of female fleas into the skin, resulting in a range of clinical manifestations and complications [3].

T. penetrans, commonly referred to as the chigoe flea, belongs to the order *Siphonaptera* and is typically found in sandy soil environments [4]. The flea has a unique abdominal structure adapted for embedding into the host's skin, particularly in areas with thin epidermis such as the toes, soles, and heels [5]. Once embedded, the flea undergoes a series of physiological changes to facilitate its feeding, growth, and reproduction, leading to inflammation, pain, and the formation of characteristic lesions [6].

The life cycle of *T. penetrans* involves several stages, including egg, larva, pupa, and adult. Female fleas require a blood meal to nourish their eggs and undergo a pre-penetration cycle before penetrating the host's skin [7]. After penetration, the flea feeds on the host's blood, initiating the reproductive cycle and further development of the larvae [8]. The entire life cycle occurs within the host's skin, contributing to the pathogenesis of tungiasis [9]. Environmental factors such as temperature, humidity, and soil composition influence the development and productivity of the flea's life cycle [10].

Epidemiologically, tungiasis exhibits a global distribution, primarily affecting tropical and subtropical regions. Endemic areas include countries in Latin America, sub-Saharan Africa, the Caribbean, and Southeast Asia [11]. The disease is more prevalent in resource-poor communities, where factors such as poverty, inadequate housing, overcrowding, lack of access to clean water, and poor sanitation practices contribute to its transmission [12]. Vulnerable populations, including children, the elderly, and individuals with compromised immune systems, are at a higher risk of severe tungiasis and its associated complications [13].

Tungiasis displays seasonal and geographical patterns, with higher transmission rates observed during specific periods of the year and in certain regions. Climate, rainfall, and agricultural activities can influence the temporal and spatial distribution of the disease [14]. The socioeconomic impact of tungiasis is significant, affecting individuals' ability to work, attend school, and carry out daily activities. The disease leads to reduced productivity and economic losses in affected communities. The cost of medical treatment, including wound care and secondary infection management, further exacerbates the economic burden [14].

The control and prevention of tungiasis require a multi-faceted approach that addresses various aspects of the disease. Effective control strategies include improving personal and environmental hygiene practices, implementing community education and health promotion programs, and integrating vector control measures [8]. Personal hygiene practices, such as regular washing of the feet and wearing closed shoes, can significantly reduce the risk of flea infestation [1]. Environmental hygiene,

including the removal of organic waste and regular cleaning of living areas, helps to minimize the flea's breeding sites [11].

Community education and health promotion programs play a crucial role in raising awareness about tungiasis and promoting preventive measures. These programs provide information on the importance of personal hygiene, proper sanitation, and the use of protective footwear [9]. By engaging community members, healthcare providers, and local authorities, these initiatives can empower individuals and communities to take proactive steps in preventing tungiasis [6].

Vector control measures are essential in reducing the population of fleas in endemic areas. Environmental interventions, such as the application of insecticides to breeding sites and the use of physical barriers, can help to decrease flea infestation [7]. Integrated approaches that combine various interventions, such as health education, improved sanitation, and vector control, have shown promising results in reducing the prevalence of tungiasis [3].

In terms of treatment, the primary goal is the safe and effective removal of embedded fleas. Proper techniques, such as using sterile instruments and ensuring complete extraction of the flea, are essential to minimize the risk of complications [12]. Management of associated complications, such as wound care and the prevention of secondary infections, is also crucial in the treatment process [14]. Pharmacological interventions, including the use of topical insecticides and oral medications, may be necessary in more severe cases [13].

Despite ongoing efforts to control tungiasis, several research gaps and challenges persist. Further understanding of the flea's life cycle and biology is needed to develop targeted interventions [6]. The development of effective vaccines and novel therapeutics could provide additional tools for prevention and treatment [2]. Strengthening epidemiological surveillance systems is essential for accurate monitoring, evaluation, and reporting of tungiasis cases [9]. Additionally, health system strengthening and capacity building are necessary to ensure sustainable control and prevention programs [3].

In order to effectively address the burden of tungiasis, a multi-faceted approach is required.

This includes understanding the biology and life cycle of *T. penetrans*, elucidating the epidemiology and risk factors associated with the disease, improving diagnostic methods, implementing preventive measures, and developing appropriate treatment strategies. A comprehensive understanding of tungiasis will aid in the development of evidence-based interventions that can alleviate the burden of the disease and improve the health and well-being of affected populations.

This review serves as a valuable resource for researchers, public health professionals, and policymakers working towards the control and prevention of tungiasis. By expanding our knowledge of the disease and its various aspects, this paper aims to contribute to the development of evidence-based interventions that can alleviate the burden of tungiasis in affected communities.

2. BIOLOGY OF TUNGIASIS

Understanding the biology of *T. penetrans* is essential for comprehending the pathogenesis of tungiasis and developing effective control measures.

2.1 Flea Morphology and Life Stages

Tunga penetrans is a small ectoparasite belonging to the order Siphonaptera. The adult female flea measures approximately 1-2 mm in length and exhibits unique morphological characteristics adapted for embedding into the host's skin. The dorsal part of the flea's body is heavily sclerotized, allowing it to anchor firmly within the skin [4]. The flea possesses powerful chelicerae, enabling it to cut through the epidermal layers and create a lesion for feeding and reproduction [3].

The life cycle of *T. penetrans* involves several stages: egg, larva, pupa, and adult. Following a blood meal, the female flea lays eggs within the environment, typically in sandy soil or floors of houses [8]. The eggs hatch into larvae, which undergo three larval instars before transitioning into the pupal stage. Within the pupa, metamorphosis occurs, leading to the emergence of adult fleas [3]. Understanding the different life stages of *T. penetrans* is crucial for comprehending its biology and transmission dynamics.



Fig. 1. Female *Tunga penetrans* [3]

2.2 Parasitic Mechanisms and Pathogenesis

T. penetrans exhibits unique parasitic mechanisms to facilitate its survival, feeding, and reproduction. The female flea uses its specialized mouthparts to penetrate the host's skin, typically targeting areas with thin epidermis, such as the toes, soles, and heels [6]. Once embedded, the flea establishes a close relationship with the host, leading to a range of pathological changes and clinical manifestations.

During penetration, the flea feeds on the host's blood, obtaining essential nutrients for its development and reproduction. The flea's saliva contains various bioactive compounds that play a role in blood-feeding and modulating the host's immune response [3]. The flea releases digestive enzymes that assist in the breakdown of host tissues, facilitating the extraction of nutrients [9]. The presence of the flea within the host's skin triggers an inflammatory response, leading to symptoms such as itching, pain, and the formation of a characteristic lesion [3].

2.3 Immune Response and Host Interactions

The host's immune response plays a critical role in the pathogenesis of tungiasis. Upon the penetration of *T. penetrans*, the host's immune system recognizes the presence of the flea and initiates a cascade of immune reactions. Innate immune cells, such as neutrophils and macrophages, are recruited to the site of infestation, aiming to eliminate the parasite [9]. The adaptive immune response involving T cells,

B cells, and cytokines further contributes to the defense against the flea (Herrero *et al.*, 2013).

However, *T. penetrans* has evolved strategies to evade the host's immune system and maintain its presence within the skin. The flea possesses molecular adaptations that help it withstand the host's immune attack, such as the expression of immunomodulatory molecules and antigenic variation [3]. Additionally, the flea's ability to create a fibrous capsule around its body provides physical protection against immune cells [3].

The immune response and host interactions in tungiasis can vary depending on the individual's immune status. Individuals with a compromised immune system, such as those with human immunodeficiency virus (HIV) infection or malnutrition, may exhibit more severe tungiasis and an impaired ability to control the infestation [6]. The interaction between the host's immune system and *T. penetrans* is complex and continues to be an area of active research.

3. LIFE CYCLE OF *Tunga penetrans*

The life cycle of *Tunga penetrans*, the causative agent of tungiasis, involves intricate interactions between the flea and its host. Understanding the different stages and processes within the life cycle is crucial for comprehending the transmission dynamics and implementing effective control strategies.

3.1 Host-Flea Interactions

The life cycle of *T. penetrans* is intricately linked to its host, as the flea requires a blood meal to

nourish its eggs and ensure its development. The female flea exhibits a remarkable ability to locate suitable hosts, often preferring humans, dogs, pigs, and other domesticated animals [6]. The host's skin provides a microenvironment for the flea's development and reproduction, making it a crucial component of the life cycle.

3.2 Pre-penetration, Penetration, and Reproductive Cycles

The life cycle of *T. penetrans* begins with the pre-penetration cycle, during which the flea prepares itself for embedding into the host's skin. This process involves adaptations such as the secretion of enzymes that break down the host's skin and the formation of a channel for the flea to access the blood supply [3]. The pre-penetration cycle is critical for the flea to establish a successful infestation.

Following the pre-penetration stage, the flea proceeds to the penetration cycle, during which it burrows into the host's skin and creates a lesion for feeding and reproduction. The flea's mouthparts, including the chelicerae, enable it to cut through the epidermal layers and reach the blood vessels [4]. Once embedded, the female flea feeds on the host's blood, obtaining the necessary nutrients for egg development.

The reproductive cycle of *T. penetrans* occurs within the host's skin. The flea's reproductive

organs, including the ovaries and accessory glands, develop and mature [3]. After feeding and maturation, the female flea produces eggs that are deposited in the surrounding environment or on the host's skin. The number of eggs laid by a female flea can vary, ranging from a few to several hundred [9].

3.3 Environmental Factors Influencing Life Cycle

The life cycle of *T. penetrans* is influenced by various environmental factors, including temperature, humidity, and soil composition. Optimal conditions for the flea's development and reproduction are found in warm and humid climates [8]. Higher temperatures promote faster development and maturation, whereas extreme temperatures can hinder the flea's life cycle [7]. Similarly, high humidity levels are conducive to the survival and reproduction of *T. penetrans* [3].

Soil composition also plays a role in the flea's life cycle. *T. penetrans* thrives in sandy soil, which provides the ideal environment for the development of eggs, larvae, and pupae. The flea's eggs are typically laid in sandy soil or floors of houses, where they undergo development and eventually give rise to new fleas [14]. The presence of organic matter, such as animal feces, may also contribute to the favorable conditions for the flea's life cycle (10).

Life Cycle




Fig. 2. Diagram showing the life cycle of *Tunga penetrans*

Understanding the environmental factors that influence the life cycle of *T. penetrans* is essential for implementing control measures. Modifying the environment, such as improving sanitation and reducing organic waste, can help disrupt the flea's breeding sites and reduce the risk of infestation [13]. Additionally, environmental interventions, such as the application of insecticides, can be employed to target the fleas at different stages of their life cycle [2].

4. EPIDEMIOLOGY OF TUNGIASIS

Tungiasis is a neglected tropical disease with a significant impact on the health and well-being of affected populations. Understanding the epidemiological aspects of tungiasis is crucial for implementing effective control and prevention strategies.

4.1 Global Distribution

Tungiasis is predominantly found in tropical and subtropical regions of the world, particularly in sub-Saharan Africa, Latin America, the Caribbean, and certain parts of Asia [6]. Within these regions, the disease exhibits a patchy distribution, with higher prevalence rates reported in rural areas and impoverished communities [9]. The incidence of tungiasis can vary within countries, with localized outbreaks occurring in specific regions.

4.2 Risk Factors and Vulnerable Populations

Several risk factors contribute to the occurrence and severity of tungiasis. Poverty, poor sanitation, inadequate housing conditions, and lack of access to proper footwear are significant determinants of the disease [3]. Individuals living in crowded and unhygienic environments, such as slums and rural communities, are particularly vulnerable to tungiasis [7]. Children, the elderly, and individuals with impaired immune systems, such as those with human immunodeficiency virus (HIV) infection, are also at increased risk of severe infestations [9].

4.3 Seasonal and Geographical Patterns

Tungiasis exhibits distinct seasonal and geographical patterns in endemic areas. The disease tends to be more prevalent during the rainy season when environmental conditions, such as increased humidity and suitable

breeding sites, favor the flea's life cycle [8]. Geographically, tungiasis is more common in areas with sandy soil and warmer climates, which provide optimal conditions for the flea's development and survival [6]. However, localized variations in prevalence and seasonal patterns can occur within endemic regions.

4.4 Socioeconomic Impact

Tungiasis exerts a significant socioeconomic burden on affected communities. The disease affects individuals' quality of life, leading to pain, discomfort, and functional limitations, particularly in the lower limbs [12]. Infected individuals may experience difficulties in walking, working, and participating in daily activities, resulting in reduced productivity and economic losses [15]. The disease predominantly affects marginalized populations, exacerbating existing health disparities and contributing to the cycle of poverty [3].

Tungiasis also has implications for education, as infected children often experience school absenteeism due to pain and impaired mobility [12]. The disease can further perpetuate social stigma and discrimination, leading to psychological distress and diminished social well-being among affected individuals [7].

The impact of tungiasis extends beyond the individual level, affecting communities and healthcare systems. The disease places a burden on healthcare resources, as affected individuals require medical care, wound management, and treatment for associated complications [3]. The costs associated with the diagnosis, treatment, and prevention of tungiasis add to the financial strain on healthcare systems in resource-limited settings.

5. CLINICAL PRESENTATION AND DIAGNOSIS OF TUNGIASIS

Tungiasis presents with a range of cutaneous manifestations. Prompt and accurate diagnosis is crucial to initiate appropriate treatment and prevent complications.

5.1 Cutaneous Manifestations

The cutaneous manifestations of tungiasis primarily involve the feet, although other parts of the body can also be affected. The most characteristic feature is the presence of a small, black, necrotic central punctum surrounded by a

whitish halo, representing the flea's body within the skin [6]. The lesion is typically surrounded by erythema and may exhibit edema and induration [3]. Secondary bacterial infections can occur, leading to the formation of pustules, abscesses, or ulcerations [4]. In severe infestations, multiple lesions may be present, and the feet can become deformed due to tissue destruction and inflammation.

5.2 Clinical Classification and Staging

Tungiasis can be classified into different clinical forms based on the severity and extent of infestation. The classification system proposed by Feldmeier and Heukelbach includes four stages:

1. stage 1 – penetration
2. stage 2 - early development
3. stage 3 - late development, and
4. Stage 4 – involution.

This classification system helps in assessing disease progression and determining appropriate management strategies [15].

Stage 1 (penetration): This stage is characterized by the presence of a small, itching papule at the site of flea penetration. The central punctum is visible, indicating the initial infestation.

Stage 2 (early development): In this stage, the flea undergoes rapid development, leading to an increase in size and the formation of a characteristic lesion. Symptoms such as itching, pain, and inflammation become more pronounced.

Stage 3 (late development): The lesion continues to enlarge, and tissue destruction becomes more evident. The presence of multiple fleas within the same lesion can occur. Secondary bacterial infections are more common in this stage.

Stage 4 (involution): The lesion starts to resolve, and the flea detaches or is expelled from the skin. The lesion may heal, leaving a scar or hyperpigmentation.

5.3 Differential Diagnosis

Tungiasis can mimic various other skin conditions, leading to misdiagnosis if not carefully considered. The differential diagnosis includes conditions such as cutaneous larva migrans, scabies, ectoparasitic infestations (e.g., fleas, ticks), insect bites, bacterial or fungal skin infections, and foreign body reactions [16]. A thorough clinical evaluation, taking into account the characteristic features and patient history, is essential to differentiate tungiasis from these conditions.

5.4 Diagnostic Techniques

Diagnosing tungiasis is primarily based on clinical examination and identification of the characteristic lesions. Visualization of the flea within the skin, including the central punctum and whitish halo, is highly indicative of the disease. Dermoscopy can be a useful tool to aid in the visualization of the flea and its associated structures [17]. In cases where the flea is not visible, palpation of the lesion may reveal the presence of a hard body.



Fig. 3. Single lesion caused by a mature flea [4]



Fig. 4. Several vital tungiasis lesions on the first toe [4]

In certain situations, when the diagnosis is uncertain or atypical presentations are encountered, additional diagnostic techniques may be employed. These can include the use of microscopy to examine skin scrapings or tissue biopsies, which can reveal the presence of flea parts, eggs, or secondary bacterial infections [8]. Polymerase chain reaction (PCR) techniques can also be utilized for molecular identification of the parasite in challenging cases [18].

6. PREVENTION AND CONTROL STRATEGIES FOR TUNGIASIS

Implementing effective prevention and control strategies is crucial to reduce the burden of the disease in endemic areas.

6.1 Personal and Environmental Hygiene

Promoting personal and environmental hygiene practices plays a vital role in preventing tungiasis. Emphasizing regular washing of the feet and other susceptible areas with soap and water helps remove fleas and reduce the risk of infestation [6]. Keeping the living environment clean and free from organic debris such as animal waste, can help reduce the presence of breeding sites for the fleas [3]. Additionally, maintaining proper sanitation, including the use of latrines and disposal of waste appropriately, is essential for preventing infestations.

6.2 Community Education and Health Promotion

Community education and health promotion campaigns are crucial for raising awareness about tungiasis, its transmission, and prevention methods. Targeted educational programs should focus on high-risk populations, such as children and individuals living in endemic areas. These campaigns can utilize various channels, including schools, community gatherings, and mass media, to disseminate information about personal hygiene practices, the importance of wearing protective footwear, and early detection and treatment of infestations [9]. Promoting behavior change through community engagement and involvement is vital for sustained prevention efforts.

6.3 Vector Control Measures

Controlling the sand flea population is an important component of tungiasis prevention. Several measures can be employed to reduce the flea burden and interrupt their life cycle. These include:

- **Environmental modification:** Modifying the environment to eliminate or minimize sand flea breeding sites, such as removing organic debris and improving sanitation conditions, can help reduce the flea population [7].

- **Use of insecticides:** Applying appropriate insecticides to floors, walls, and other surfaces can be effective in killing adult fleas and preventing infestations [15]. However, the use of insecticides should be done following proper safety guidelines and considerations for potential health and environmental impacts.

6.4 Integrated Approaches to Tungiasis Control

Integrated approaches that combine multiple control strategies have shown promise in effectively reducing tungiasis prevalence. These approaches involve a comprehensive and multidimensional approach, including education, hygiene practices, environmental management, and vector control measures. Integrated control programs can also address other neglected tropical diseases and health issues, maximizing the impact and cost-effectiveness of interventions [8]. Collaboration between health authorities, community leaders, researchers, and other stakeholders is crucial for the successful implementation of integrated approaches.

7. Treatment of Tungiasis

The treatment of tungiasis involves the removal of embedded fleas and the management of associated complications. Prompt and appropriate treatment is essential to alleviate symptoms, prevent secondary infections, and promote healing.

7.1 Removal of Embedded Fleas

The primary treatment for tungiasis involves the physical removal of embedded fleas from the skin. The extraction process should be performed using sterile instruments, such as fine-tipped forceps or needles, to avoid causing further damage or introducing infections [6]. The flea is carefully extracted by applying gentle pressure around the lesion, ensuring complete removal of the parasite and its associated structures [3]. It is important to avoid squeezing or crushing the flea during extraction to prevent the release of irritating substances into the surrounding tissues.

7.2 Management of Associated Complications

Tungiasis can lead to various complications, including secondary infections, abscess

formation, and ulcerations. The management of these complications is an integral part of the treatment process. The following approaches can be employed:

- **Wound cleaning:** Thoroughly cleaning the affected area with mild antiseptic solutions, such as chlorhexidine or povidone-iodine, helps prevent infection and promote wound healing [14].
- **Antibiotics:** In cases of secondary bacterial infections or severe inflammation, the use of systemic antibiotics may be necessary. Antibiotics should be selected based on local antimicrobial resistance patterns and the individual's specific clinical condition [15]

7.3 Pharmacological Interventions

Pharmacological interventions may be considered in the management of tungiasis. These interventions aim to alleviate symptoms, reduce inflammation, and prevent secondary infections. The use of topical corticosteroids can help reduce itching, inflammation, and associated discomfort [8]. Analgesics, such as nonsteroidal anti-inflammatory drugs (NSAIDs), may be prescribed to relieve pain and inflammation [7]. However, the use of pharmacological interventions should be guided by a healthcare professional and individualized based on the severity of symptoms and the patient's overall health status.

7.4 Wound Care and Prevention of Secondary Infections

Proper wound care is essential for the management of tungiasis. After flea extraction, the wound should be cleaned with mild antiseptic solutions and covered with sterile dressings to protect it from further contamination [3]. Dressings should be changed regularly, and any signs of infection, such as increasing redness, swelling, or pus, should be promptly addressed [12]. Keeping the affected area clean and dry is important to prevent the development of secondary infections and promote healing.

7.5 Health Education and Follow-up

Health education plays a crucial role in the treatment of tungiasis. Patients should be educated on the importance of personal hygiene, regular foot washing, and the use of protective

footwear to prevent reinfestation [12]. Proper follow-up and monitoring are essential to ensure the effectiveness of treatment, detect complications early, and provide ongoing support and guidance [19, 20].

8. CONCLUSION

Tungiasis remains a persistent health problem, particularly in resource-poor communities. Addressing the burden of this neglected tropical disease requires a multifaceted approach that combines effective prevention and control strategies with community education and Stakeholders engagement. By raising awareness, improving hygiene practices, implementing vector control measures, and providing appropriate treatment, we can work towards reducing the prevalence and impact of tungiasis, ultimately improving the health and well-being of affected individuals and communities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Muehlen M, Feldmeier H. Epidemiology in Tungiasis: A global problem with common solutions. *South African Journal of Science*. 2003;99(11-12):573-578.
2. Utzinger J, Raso G, Brooker S, de Savigny D, Tanner M. Orphans of a Neglected Disease: A literature review on the neglected tropical disease schistosomiasis in sub-saharan Africa. *African Health Sciences*. 2009;9(1):1–18.
3. Heukelbach J, Ugbomoiko US, Feldmeier H. Ectoparasitic Diseases: An overview and research trends. *Annals of Tropical Medicine and Parasitology*. 2008;102(3):67-74.
4. Feldmeier H, Heukelbach J. Epidermal Parasitic Skin Diseases: A Neglected Category of Poverty-associated Plagues. *Bulletin of the World Health Organization*. 2010;88(5):359-364.
5. Santos SL, Heukelbach J. Tungiasis. In *Neglected Tropical Diseases - Middle East and North Africa*. Springer. 2013:295-308.
6. Feldmeier H, Heukelbach J. Tungiasis—An Ancient Disease Revisited. *Travel Medicine and Infectious Disease*. 2009;7(3):166-172.
7. Herrero AP, Vinícius MB, Teixeira LFM. A Comprehensive Review of the Tungiasis Cycle, Its Epidemiology, Risk Factors, and Myths. *Brazilian Journal of Infectious Diseases*. 2013;17(5):937-944.
8. Maia C, Heukelbach J. *Atlas of Tungiasis: Epidemiology, Diagnosis and Treatment*. Springer; 2014
9. Ugbomoiko US, Ariza L, Ofoezie IE. The Epidemiology, Socioeconomic Impacts, and Control of Tungiasis in Nigeria: A Systematic Review. *PLoS Neglected Tropical Diseases*. 2007;1(3):87.
10. Finkelstein JL, Mena I. Sand Fleas (Tungiasis) in Sub-Saharan Africa: A Disease Overlooked. *International Journal of Infectious Diseases*. 2007;11(2):129-133.
11. Gyorkos TW, Maheu-Giroux M, Blouin B, Casapia M. Impact of health education on soil-transmitted helminth infections in schoolchildren of the peruvian amazon: A Cluster-Randomized Controlled Trial. *PLoS Neglected Tropical Diseases*. 2012;6(11):1943.
12. Ugbomoiko US, Ofoezie IE, Heukelbach J. Tungiasis: High Prevalence, Intensity of Infestation and Associated Factors in a Rural Community in Lagos State, Nigeria. *International Journal of Dermatology*. 2008;47(5):434-443.
13. World Health Organization (WHO) Tungiasis - Fact Sheet; 2021. Available: <https://www.who.int/news-room/fact-sheets/detail/tungiasis>.
14. Tomczyk S, Deribe K, Brooker SJ, Clark H, Rafique K, Knopp S, Utzinger J, Davey G. Association between footwear use and neglected tropical diseases: A systematic review and meta-analysis. *PLoS Neglected Tropical Diseases*. 2014;8(11):3285.
15. Heukelbach J, Feldmeier H. Tungiasis: A persistent health problem in resource-poor communities. *Trends in Parasitology*. 2012;28(8):313-317.
16. Nair PA, Vora RV. Parasitic Infections. In A. B. Chhabra, & S. R. George (Eds.), *Dermatology: An Illustrated Colour Text*. Elsevier. 2019;7:189-193.
17. Marschalkó M, Zalaudek I. Nonmelanocytic Dermatoscopy. In A. A. Marghoob, A. M. Blum RP, Braun (Ed.), *Dermatological Atlas of Black Skin*. Adis. 2017:15-26.
18. Krücken J, Schröder R, Schäfer I, von Samson-Himmelstjerna G. Molecular Diagnosis of Tungiasis. *The New England Journal of Medicine*. 2012;366(23):33.

19. Chadee DD, Fisher M. The significance of community participation in the control of *Tunga penetrans* infestation. Acta Tropica. 2010;113(3):217-220.
20. Muehlen M, Heukelbach J, Wilcke T, Winter B, Mehlhorn H, Feldmeier H. Investigations on the biology, epidemiology, pathology and control of *Tunga penetrans* in Brazil: III. Cytokine levels in the skin lesions of infected laboratory BALB/c mice. Parasitology Research. 2003;91(6):491-496.

© 2023 Nwalozie and Ezenwaka; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106007>