



Development and Nutritional Analysis of Taro Powder [*Colocasia esculenta* (L.) Schott.] Enriched with Natural Colorants

Simanta Das ^{a*}, Ranjan Kandali ^{a++} and Samindra Baishya ^{a#}

^a Department of Biochemistry and Agricultural Chemistry, Assam Agricultural University, Jorhat, Assam, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author SD did the formal analysis, investigation, data visualization, wrote original draft of the manuscript. Author RK did study conceptualization, performed methodology, helped in administration and supervised the study. Author SB did the funding acquisition. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92523

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104104>

Original Research Article

Received: 25/05/2023

Accepted: 28/07/2023

Published: 04/08/2023

ABSTRACT

Aims: To prepare an anthocyanin rich food product from Indigenous plant species.

Study Design: Lab experiments based on CRD statistical analysis.

Place and Duration of Study: Department biochemistry and agricultural chemistry, Assam Agricultural university, Jorhat, Assam, India between June 2018 and January 2020.

Methodology: We collected anthocyanin rich plant samples viz. fruits of jamun (*S. cumini* L. and *S. fruticosum* Roxb. DC.), full-grown flowers of rose (*R. indica* L.) and studied the stability of total anthocyanin under pH, temperature and light. Taro powder was selected with low oxalic acid content to prepare an anthocyanin rich food product. The storage duration was also study biochemically for its commercial application.

⁺⁺ Professor;

[#] HOD and Professor;

*Corresponding author: E-mail: dsimanta650@gmail.com, simantad37@gmail.com;

Results: total anthocyanin content as 124.87±0.07 mg/100g (*S. cumini* L.) followed by 115.26±0.09 mg/100g (*R. indica* L.) and 91.41±0.09 mg/100 g (*S. fruticosum* Roxb. DC.). The anthocyanin pigments were found to be stable at an acidic pH (up to pH 5) and below 40° C temperature and activity was maintained under a light of 2500 lux for 6 hours. A novel food product was developed by immobilizing pigments extracted from *S. cumini* L., *S. fruticosum* Roxb. DC. and *R. indica* L. on *Colocasia esculenta* (L.) Schott. (variety: Ahina) powder.

Conclusion: Anthocyanins act as potent antioxidant but its stability after extraction is a major concern. By applying different storage condition and suitable carrier material selection it can be stored for long.

Keywords: Indigenous plants; food colorants; anthocyanin; stability; immobilization; food product.

ABBREVIATIONS

TAC : Total Anthocyanin Content
TPC : Total Phenolic Content
TFC : Total Flavonoid Content
TSS : Total Soluble Sugar
TSP : Total Soluble Protein
EMS : Error Mean Square
GAE : Gallic Acid Equivalent
QE : Quercetin Equivalent

1. INTRODUCTION

Food color industry is thought to be increased up to 15 % in recent years. Natural colors like anthocyanin, betacyanin, carotenoids, betalins and chlorophyll etc. are now considered as substitutes for synthetic colorants. In terms of natural colorants, secondary metabolites mainly phenolic compounds are found to be rich in different colors due to their unique structural features. Structurally, anthocyanins are glycosides of polyhydroxy and polymethoxy derivatives of 2-phenylbenzopyryryl salts and are composed of an aglycone known as anthocyanidin. A carbohydrate residue (glucose, xylose, galactose, arabinose, rhamnose or rutinose) usually connected to the anthocyanidin skeleton through the C3 hydroxyl organization in ring C. The functioning of anthocyanin is distinctly dependent on (i) Chemical structure, ring orientation and role of hydroxyl or methoxy groups (ii) Quantity of glycosyl, acyl groups, sugar acylation and the identification of acylating agent and (iii) Stability under varying pH, temperature, light and storage situations [1]. Anthocyanin as a compound is very unstable in natural condition but with proper selection of carrier material it can be stored for long duration. Taro powder does not have any anthocyanin compound if compared with high anthocyanin rich plant products like fruits and flowers of different indigenous. However, high carbohydrate content of taro can be used to bind anthocyanin

after proper extraction techniques and can be stored long with proper study of storage condition. Provide a factual background, clearly defined problem, proposed solution, a brief literature survey and the scope and justification of the work done.

2. MATERIALS AND METHODS

2.1 Chemical, Reagents and Instruments

HPLC grade methanol, Milli-Q water (0.42 mm membrane filtered), ethanol and standards of anthocyanins as Delphinidin Chloride (CAS 28-53-0) and Pelargonidin chloride (CAS 134-04-3) were purchased from Sigma-Aldrich, India. Aluminium chloride, ascorbic acid, Bovine Serum Albumin (BSA), Folin Ciocalteu Reagent (FCR), hydrochloric acid (HCL), quercetin, nitric acid (HNO₃), sulphuric acid (H₂SO₄), anthrone and sodium carbonate (Na₂CO₃) were procured from Merck (India). All other chemicals used were of analytical grade. Instruments like Eutech pH meter (pH700), Waters 2487 HPLC (UV-Vis detector); Toledo ME204 weighing balance; Chemito make: model Spectrascan 2600 spectrophotometer and Hermle centrifuge 236s were majorly used.

2.2 Materials

Freshly ripped fruits of Jamun (*S. cumini* L. and *S. fruticosum* Roxb. DC.), Malabar spinach (*B. alba* L. and *B. rubra* L.) and full grown flowers of rose (*R. indica* L., *R. damascena* Mill. and *R. bracteata* J.C. Wendl.) were collected from different places of Jorhat, Assam (GPS: 26° 45' 0.00" -N and 94° 13' 12.00" -E). Samples were washed with distilled water, wrapped up in blotting paper to remove water and stored in refrigerated condition (4°C). Matured tubers of *Colocasia esculenta* (L.) Schott. (variety: Ahina) and fruits of Lemon cv. Assam lemon (*Citrus limon* Burm.) were collected from Horticultural orchard of Assam Agricultural University, Jorhat, Assam.




Fig. 1. Basic structure anthocyanidin

2.3 Preliminary Analysis

2.3.1 Total moisture content

Moisture content was determined by of AOAC (1970) method. For this, approximately 10 g of sample was weighed in aluminium moisture boxes and dried in an oven at $54 \pm 2^\circ\text{C}$ till constant weight. Total moisture content was expressed by following formula.

Moisture Content (g/100g sample) =

$$\frac{\text{Initial weight (g)} - \text{Final weight (g)}}{\text{Weight of the sample (g)}}$$

2.3.2 Extraction of colorant

The extraction of colorant was done by grinding and maceration in 100 ml distilled water followed by filtration with muslin cloth [2]. Filtered samples were centrifuged at 10,000 rpm for 15 minutes. The supernatant was again filtered with Whatman no. 42 filter paper. The filtrate was stored in refrigerated condition (4°C) for further analysis

2.3.3 Determination of total phenolic content

Qualitative analysis of phenolic content was done by the ferric chloride. The total phenolic content of extracts was determined using the Folin–

Ciocalteu Reagent [3]. Gallic acid was used as standard and TPC was expressed as mg gallic acid equivalent weight per 100 g fresh mass.

2.3.4 Determination of total flavonoids

The total flavonoid content was determined by colorimetric method [4] and expressed in mg of Quercetin equivalent (QE)/100g of fresh weight.

2.3.5 Total monomeric anthocyanin pigment content

Total monomeric anthocyanin pigment content was measured by pH differential method, which is based on the structural change of the anthocyanin chromophore between pH 1.0 and 4.5 [5]. Potassium chloride buffer (KCl, 0.025 M, pH 1.0) and sodium acetate buffer ($\text{CH}_3\text{CO}_2\text{Na} \cdot 3\text{H}_2\text{O}$ 0.4 M, pH 4.5) were prepared and diluted with extracted colorants (5:1). HCl (0.2N) was used to adjust the pH. After 15 mins of equilibration, the absorbance of two dilutions was read at 510 nm and 700 nm using a spectrophotometer. Results of total monomeric anthocyanin pigment content expressed as cyanidin-3-glucoside equivalents in mg/100g by the following formula:

Total monomeric anthocyanin pigment content =

$$\frac{(A \times MW \times DF \times 1000)}{\epsilon}$$

Where,

A= (A₅₁₃-A₇₀₀) pH 1.0–(A₅₁₃-A₇₀₀) pH 4.5

MW= molecular weight of cyaniding-3-glucoside (449.2)

DF= Dilution factor

ε= Molar extinction coefficient (26,900 L mol⁻¹ cm⁻¹)

2.3.6 pH stability

The method by Selim *et al.* [6] with slight modifications was used to evaluate the stability of anthocyanin extract at varying pH. 2 ml extracted colorants mixed with buffers ranging from pH 1 to pH 10 were prepared and made up to 20 ml with the desired buffer. The test tubes were covered with aluminium foil and stored at room temperature for 2 hours.

2.3.7 Thermal stability

The thermal stability of extracted colorants was determined according to Amr and Al-Tamimi [7] with some modifications. 0.1 ml of extracted colorant was made up to 25 ml with 0.1 M (pH 3.5) citrate-phosphate buffer and kept in -4°C, 20°C, 40°C, 60°C, 80 °C and 100 °C for 6 hours. Absorbance was read at 520 nm and percentage retention of colorant was calculated using following equation:

$$\text{Pigment retention} = \frac{\text{Absorbance after heating}}{\text{Absorbance before heating}} \times 100$$

2.3.8 Light stability

Extracted colorants were prepared in equal proportion (1:1) in citrate- phosphate buffer (pH 3.5) in sealed test tubes [8]. To maintain average daylight condition (2500 lux), a square hardboard box (75 cm x 75cm x 75 cm) laden internally with aluminium foil. Two 40W bulbs were placed at 10 cm distance inside the box. Buffer mixed with colorants were incubated under artificial light at room temperature and readings were taken in-between 2h, 6h, 12h, 24 h, and 5days.

2.4 Development of an Anthocyanin-rich Solid Food Product

2.4.1 Preparation of *Colocasia esculenta* (L.) Schott. (variety: Ahina) powder

Colocasia esculenta (L.) Schott. (var. Ahina) powder was prepared by the method of Hazarika

[9]. Pilled tubers were washed, cut into small pieces, air-dried at room temperature, ground to a fine powder and stored in a desiccator for further analysis. To immobilize colorant on edible carrier food material, the method given by Debnath, 2016 was followed. Powdered *Colocasia esculenta* (L.) Schott. (variety: Ahina) was added to the concentrated dye in a ratio of 1:3 (w/v) to make it a viscous solution, then incubated at 4°C overnight, further ground to powder and stored in a sealed clear glass bottle covered with aluminum foil in refrigerated condition (4°C).

2.5 Nutritional Analysis of Carrier Materials

Nutritional analysis of carrier material *viz.* *Colocasia esculenta* (L.) Schott. (variety: Ahina) powder and Assam lemon beverage was done before and after immobilization of colorants extracted from selected plant samples.

2.5.1 Total phenol and total anthocyanin

Total phenolics and total anthocyanin content were determined by Folin Ciocalteu Reagent [3] and pH differential method [5] respectively.

2.5.2 Total soluble protein

Total soluble protein was determined by Folin-Ciocalteu Reagent [10]. To 0.1 ml of the sample extract, 0.9 ml distilled water was added to make up the volume. 5 ml of alkaline copper solution was added, mixed well by vortexing and incubated at room temperature for 10 mins. After addition of 0.5 ml of Folin-Ciocalteu Reagent and readings were taken at 650 nm. The protein concentration was calculated out from the standard curve of BSA (Bovine Serum Albumin) and expressed in mg of protein content per 100 g on a fresh weight basis.

2.5.3 Total starch content

Total starch content of *Colocasia esculenta* (L.) Schott. (variety: Ahina) powder was determined by the anthrone method [11]. 0.2 g of sample was homogenized in 80% ethanol and centrifuged. To the extracted residue 5.0 ml of water and 6.5 ml of perchloric acid (52%) were added. After centrifugation, the supernatant was made up to 50 ml and 4 ml of anthrone reagent was added followed by heating in boiling water for 8 to 10 minutes. After cooling, the reading was taken at 630 nm.

2.5.4 Total Ash and minerals (Phosphorous and Iron)

The ash content of *Colocasia esculenta* (var. *Ahina*) powder and peel of Lemon cv. Assam lemon (*Citrus limon* Burm.) was determined by the method of AOAC (1970) [12]. Mineral solutions were prepared and analyzed for the determination of Phosphorous and Iron [13].

2.6 Evaluation after Immobilization of Colorants

After immobilization of colorants, *Colocasia esculenta* (L.) Schott. (variety: Ahina) powder immobilized with colorants were evaluated for the change in total anthocyanin content (mg/100gm) and total starch content (%) with in storage period of one month (4°C). Similarly, change in total anthocyanin content of Lemon cv. Assam lemon (*Citrus limon* Burm.) beverage mixed with colorants was also evaluated for same period of storage period (4°C).

2.7 Statistical Analysis

The data obtained from laboratory experiments were analyzed statistically. The experiment was laid out in Complete Randomized Design (CRD). All analysis was performed in triplicate and the average has been reported. The data were analyzed by one-way analysis of variance (ANOVA). The standard error of the mean difference (S. Ed.±) was calculated. The treatment means were compared among themselves by calculating critical difference (CD at P<0.05) as per the method of Panse and Sukhatrne [1].

The analysis of variance (ANOVA) was done with treatments and replications. The critical differences were calculated by the formula:

$$CD = t_{0.05, \text{error d.f.}} \times S. Ed$$

$$\text{Where, } S. Ed = \sqrt{\frac{2EMS}{r}}$$

3. RESULTS AND DISCUSSION

The design of experiment for present investigation is based on selection of experimental design by ISO 3534-3:1985 [14]. Accuracy (trueness and precision) of measurement methods and results were as per the guidelines of ISO 5725-1:1994 [15]. For all the recorded data we have taken three

replications and standard deviation was in accordance with the replicated value. Out of seven selected plant samples flower of *Rosa indica* L. had highest total phenolic content of 1516.52±0.03 mg GAE/100g. In consideration of total flavonoid content among seven different plant samples, *Rosa damascena* Mill. and fruits of *S.cumini* L. had a higher amount of total flavonoid content of 262.87 ±0.07 mg QE/100g. (Table 1). Fruit extract of *S. cumini* L. (124.87±0.07 mg/100g) had the highest total anthocyanin concentration, followed by flower extract of *Rosa indica* L. (115.26± 0.09 mg/100g) and fruit extract of *S. fruticosum* Roxb. DC. (91.41± 0.09 mg/100g). The total anthocyanin content of Jamun fruit was found to be 195.58±6.15 mg/100g as reported by Ghosh et al. *Rosa indica* L. had total anthocyanin content of 115.26±0.091 mg/100g. Poonam [16] studied anthocyanin pigments in Indian rose types and discovered total anthocyanin concentration ranged from 2.14±0.12 mg/100g to 667.46±21.27 mg/100g. Characterization of anthocyanins in selected plant samples was done by the UHPLC method comparing with the retention time and absorbance peaks of two selected anthocyanin standards viz Delphinidin-3-glucoside (RT:10.713 minutes) and Pelargonidin -3-glucoside (RT:11.313 minutes). Pelargonidin-3-glucoside was present in a major amount followed by delphinidin-3-glucoside in extracted colorant of *S. cumini* L. and *S. fruticosum* Roxb. DC. fruits. On the other hand, delphinidin-3-glucoside was the major anthocyanin followed by pelargonidin-3-glucoside in the extracted colorant of *R. indica* L.

Both in presence or absence of light, the temperature had a significant impact on the stability of anthocyanin pigment. The anthocyanin concentration was tested for 6 hours at 20°C, 40°C, 60°C, 80°C and 100°C, and it was observed that anthocyanin degradation was less at lower temperatures (below 40°C). More than 50% pigment retention up was found for *R. indica* L. (111.65 mg/100g) at 40°C. However, at 60°C, 50% pigment retention was observed for *S. cumini* L. (68.72 mg/100g) and *S. fruticosum* Roxb. DC. (75.34 mg/100g) (Fig. 3). In contrast, fast degradation rate was found above 60°C for 6 hours. The rapid decomposition of anthocyanin at higher temperatures could be attributed to the hydrolysis of the 3-Glycoside structure, which protects unstable anthocyanin. Anthocyanin acylation contributes to their stability as

Table 1. Preliminary investigation of selected plant samples: 1) *S. cumini* L., 2) *S. fruticosum* Roxb. DC., 3) *Basella alba* L., 4) *Basella rubra* L., 5) *R. indica* L., 6) *R. damascena* Mill. and 7) *R. bracteata* J.C. Wendl

Species	Part used	Weight (g)	Length (cm)	Breadth (cm)	Moisture (%)	TPC (mg GAE/100g)	TFC (mg QE/100g)	TAC (mg/100g)
1	Fruit	6.31±0.02	2.21±0.09	2.04±0.08	81.92±0.16	1149.28 ±0.07	262.87±0.07	124.87 ± 0.07
2	Fruit	2.56±0.06	0.64±0.03	0.69±0.06	78.48±0.23	741.25± 0.05	176.23 ±0.17	91.41 ± 0.09
3	Fruit	0.79±0.03	0.52±0.03	0.78±0.02	84.12±0.42	548.98± 0.42	184.16± 0.12	5.96 ± 0.06
4	Fruit	0.76±0.04	0.54±0.05	0.77±0.03	83.88±0.09	938.47± 0.83	292.52± 0.24	8.95 ± 0.11
5	Flower	4.03±0.04	3.83±0.09	3.30±0.06	81.37±0.16	1516.52±0.03	172.91± 0.49	115.26 ± 0.09
6	Flower	4.73±0.02	4.07±0.08	3.47±0.02	76.65±0.17	1380.67 ±0.04	262.88± 0.05	2.36 ± 0.17
7	Flower	3.20±0.01	3.87±0.03	3.17±0.07	78.78±0.27	1347.23± 0.16	196.12± 0.89	0.66 ± 0.07

Table 2. Effect of temperature (a), pH (b) and light exposure duration (2500 lux) (c) on TAC of extracted colorants from fruits of *S. Cumini* L, *S. fruticosum* Roxb. DC. and flowers of *Rosa indica* L. Within optimum condition i.e., 40 °C (temperature), pH 4 and exposure of light (2500 lux) up to six hours resulted in more than 50 % pigment retention

a. Sample	4°C	20°C	40 °C	60 °C	80 °C	100 °C
<i>S. cumini</i> L.	124.87 (100)	119.68 (95.84)	77.29 (61.89)	68.72 (55)	35.65 (36.16)	19.02(15.23)
<i>S. fruticosum</i> Roxb. DC.	91.41 (100)	89.24 (97.58)	82.85 (90.63)	75.34 (82.41)	45.38 (49.64)	11.26 (12.31)
<i>R. indica</i> L.	115.26 (100)	113.58 (98.54)	111.65 (96.86)	8.23 (7.1)	7.63 (6.61)	3.68 (3.19)
CD _(0.05)	0.073 (0.08)	0.0124 (0.35)	0.073 (0.35)	0.06 (0.06)	0.08 (1.06)	0.074 (0.06)
SE(d)	0.034 (0.03)	0.006 (0.16)	0.033 (0.16)	0.025 (0.02)	0.037 (0.49)	0.035 (0.02)
SE(m)	0.024 (0.02)	0.004 (0.11)	0.023 (0.11)	0.018 (0.01)	0.265 (0.35)	0.024 (0.01)

b. Sample	pH 1	pH 2	pH 3	pH 4	pH 5	pH 6	pH 7	pH 8	pH 9	pH10
<i>S. cumini</i> L.	102.36 (81.97)	122.14 (97.81)	118.96 (95.26)	110.93 (88.83)	84.23 (67.45)	55.24 (44.23)	26.44 (21.17)	14.67 (11.74)	4.54 (3.63)	3.65 (2.92)
<i>S. fruticosum</i> Roxb. DC.	69.35 (75.86)	90.57 (99.08)	88.44 (96.75)	82.51 (90.26)	42.62 (37.16)	37.16 (40.65)	19.21 (21.01)	12.04 (13.17)	3.05 (3.33)	1.25 (1.36)
<i>R. indica</i> L.	91.47 (79.35)	99.38 (86.22)	112.45 (97.56)	106.86 (93.71)	85.98 (74.59)	74.88 (64.96)	55.02 (47.73)	18.86 (16.36)	7.44 (6.45)	1.93 (1.67)
CD _(0.05)	0.072 (0.051)	0.056 (0.045)	0.06 (0.067)	0.063 (0.12)	0.067 (0.78)	0.066 (0.098)	0.07 (0.098)	0.061 (0.53)	0.062 (0.064)	0.069 (0.028)
SE(d)	0.033 (0.017)	0.025 (0.089)	0.026 (0.032)	0.031 (0.051)	0.033 (0.049)	0.032 (0.073)	0.031 (0.049)	0.028 (0.018)	0.029 (0.064)	0.032 (0.048)
SE(m)	0.023 (0.011)	0.018 (0.13)	0.018 (0.011)	0.021 (0.017)	0.022 (0.007)	0.022 (0.014)	0.022 (0.014)	0.019 (0.008)	0.02 (0.01)	0.023 (0.014)

c. Sample	2500lux (2h)	2500lux (6h)	2500 lux (12h)	2500lux (24h)	2500lux (5days)
<i>S. cumini</i> L.	115.65 (92.61)	86.55 (69.31)	32.21 (25.79)	12.33 (9.87)	0.94 (0.75)
<i>S. fruticosum</i> Roxb. DC.	82.38 (90.12)	67.31 (73.63)	26.93 (29.46)	4.13 (4.51)	0.62 (0.67)
<i>R. indica</i> L.	101.21 (87.12)	81.07 (70.33)	53.08 (46.05)	9.33 (8.09)	0.07 (0.60)
CD _(0.05)	0.07 (0.09)	0.186 (0.067)	0.06 (0.071)	0.06 (0.019)	0.063 (0.011)
SE(d)	0.032 (0.101)	0.087 (0.043)	0.027 (0.121)	0.028 (0.021)	0.029 (0.07)
SE(m)	0.022 (0.013)	0.061 (0.02)	0.019 (0.013)	0.019 (0.015)	0.02 (0.01)

*Values were expressed in average of three replications

temperatures rise [17-20]. The per cent pigment retention of extracted colorants showed similar results. Per cent pigment retention of extracted colorants was found to be greater than 50% at 60°C temperature for both the fruits of jamun (55 and 82.41 %). Whereas, flower extract of *R. indica* L. have the percentage pigment retention reduced as the temperature increased (Table 2a).

After one hour of incubation at each pH (1-10), the total anthocyanin content was measured. At pH 2 the maximum total anthocyanin content for extracted colorants from plant samples was observed as 122.14 (*S. cumini* L.), 90.57 (*S. fruticosum* Roxb. DC.) and 99.38 (*R. indica* L.) expressed in mg/100g. With rising pH the total anthocyanin content steadily dropped, however more than 50% of total anthocyanin was retained up to pH 4. As the pH was raised higher, the overall anthocyanin content declined up to 2.92, 1.36 and 1.93 mg/100g (Table 2a). In acid solutions, some anthocyanins were red, violet or

purple in neutral solutions and blue in alkaline pH solutions. Anthocyanin molecules in solution were found to be in a state of equilibrium between the colored cationic form and the colorless pseudo base. It became protonated and formed a positive ion or cation at low pH, deprotonated as pH increased, and formed a negative ion or anion at high pH [21,22].

The stability of extracted colorants under light was investigated by incubating sample extracts for 2 hours, 6 hours, 12 hours, 24 hours, and 5 days under 2,500 lux light intensity (Table 2c). Light exposure for 2 hours (2500 lux) resulted in pigment retention of 92.61 % (115.61 mg/100g), 90.12% (82.38 mg/100g) and 87.12 % (101.21 mg/100g) in extracted colorants from *S. cumini* L., *S. fruticosum* Roxb. DC. and *R. indica* L [23,24]. However, with increasing light exposure duration, TAC fell drastically. 24 hours of light exposure results in less than 10% of pigment retention i.e., 9.87% (12.33 mg/100 g), 4.51 % (4.13 mg/100g) and 8.09% (9.33 mg/100g). To




Fig. 2. Effect of A. temperature B. pH and C. duration of light exposure (2500 lux) on extracted colorants obtained from fruits of *S. cumini* L., *S. fruticosum* Roxb.DC. and flowers of *R. indica* L.

Table 3. Nutritional comparison of final food product from *Colocasia esculenta* (L.) Schott. (variety: Ahina) with raw powder

Parameters	Nutritional analysis of raw Taro powder	Nutritional analysis of final product mixed with colorants		
		<i>S. Cumini</i> L	<i>S. fruticosum</i> Roxb. DC.	<i>Rosa indica</i> L.
Total phenol content (mg GAE /100g)	46.78±0.19	71.83±0.08	49.42±0.14	69.54±0.35
Total anthocyanin content (mg/100g)	0.84±0.32	44.76±0.70	35.66±0.67	48.71±0.39
Total soluble protein (%)	0.86±0.16	0.84±0.28	0.82±0.14	0.84±0.52
Total starch content (%)	92.63±0.64	91.98±0.42	92.14±0.06	91.86±0.09
Total ash content (%)	1.83±0.09	1.79±0.14	1.80±0.45	1.80±0.24
Phosphorus (mg/100g)	28.79±0.13	28.77±0.05	27.61±0.22	26.84±0.17
Iron (mg/100g)	7.63±0.38	7.42±0.12	6.99±0.08	7.09±0.32

*Values were expressed in average of three replications




Fig. 3. Concentrated colorants extracted from A. *S. cumini* l. B. *S. fruticosum* roxb. DC. C. *R. indica* L. which were later mixed and immobilized on taro powder. Fig. D, E and F represents the taro powder with concentrated colorants from *S. cumini* L., *S. fruticosum* roxb. DC. and *R. indica* L

retain the TAC, extracted colorants must be mixed with some carrier material that increase its self-life. Stability of anthocyanin under light rely upon the presence of oxygen, interactions with other components (sugars and ascorbic acid), hydrolyzation at glycoside linkages to produce chalcone, alpha diketones and velocity of free sugar formation [1,18]. *S. cumini* L. (124.87±0.07 mg/100g), *S. fruticosum* Roxb. DC.

(115.26±0.09 mg/100 g) and *Rosa indica* L. (91.41±0.09 mg/100g) were found with higher total anthocyanin content (Table 1) from the spectroscopic determination and UHPLC characterization of anthocyanins. Extracted colorants from these three samples were subjected for immobilization on inert carrier material to develop a solid anthocyanin rich food product (Fig. 2).

Table 4. Change in TAC (mg/100g), Total starch content (%) and per cent degradation of taro powder mixed with extracted concentrated colorants from *S. Cumini* L, *S. fruticosum* Roxb. DC. and *Rosa indica* L. during storage up to 30 days at 4°C

Colorants extracted from	1 day		2 days		4 days		6 days		12 days		30days	
	TAC (mg/100g)	TSC (%)	TAC (mg/100g)	TSC (%)	TAC (mg/100g)	TSC (%)	TAC (mg/100g)	TSC (%)	TAC (mg/100g)	TSC (%)	TAC (mg/100g)	TSC (%)
<i>S. Cumini</i>	44.60	91.97	44.57	91.93	44.43	91.84	43.98	91.76	42.72	90.68	39.42	90.07
L	(99.64)	(99.98)	(99.57)	(99.94)	(99.26)	(99.84)	(98.25)	(99.76)	(95.44)	(98.58)	(88.06)	(97.92)
<i>S. fruticosum</i>	35.63	92.12	35.58	92.08	35.47	92.01	35.12	91.98	34.75	91.52	31.33	89.96
Roxb. DC.	(99.91)	(99.97)	(99.77)	(99.93)	(99.46)	(99.85)	(98.48)	(99.82)	(97.44)	(99.32)	(87.86)	(97.63)
<i>Rosa indica</i>	48.70	91.86	48.67	91.85	48.59	91.83	48.56	91.76	46.23	91.25	44.51	90.84
L.	(99.97)	(100)	(99.91)	(99.98)	(99.75)	(99.96)	(99.69)	(99.89)	(94.91)	(99.35)	(91.38)	(98.88)
CD_(0.05)	0.066	NS	0.103	NS	NS	0.053	NS	NS	NS	0.061	0.082	0.065
	(NS)	(NS)	(NS)	(NS)	(0.110)	(0.064)	(0.032)	(NS)	(0.12)	(NS)	(NS)	(0.012)
SE(d)	0.027	0.018	0.029	0.014	0.017	0.021	0.016	0.019	0.017	0.021	0.033	0.026
	(0.012)	(NS)	(0.015)	(0.051)	(0.032)	(0.03)	(0.013)	(0.12)	(0.021)	(NS)	(0.078)	(0.08)
SE(m)	0.019	0.012	0.021	0.010	0.012	0.015	0.011	0.013	0.012	0.015	0.023	0.018
	(0.007)	(0.078)	(0.012)	(0.032)	(0.017)	(0.102)	(0.01)	(0.041)	(0.014)	(NS)	(0.021)	(NS)

*Values were expressed in average of three replications




Fig. 4. Graphical representation of change in TSC (%) and TAC (mg/100g) in final food products of taro powder mixed with concentrated colorants extracted from fruits of *S. cumini* L., *S. fruticosum* Roxb.DC. and flowers of *R. indica* L. during one month of storage period shows decreasing pattern of TSC and TAC with increasing time

Total phenol, total anthocyanin, total soluble protein, total starch, total ash, phosphorus and iron content in raw *Colocasia esculenta* (L.) Schott. (variety: Ahina) were found to be 46.78 ± 0.19 mg GAE /100g, 0.84 ± 0.32 mg/100g, $0.86 \pm 0.16\%$, $92.63 \pm 0.64\%$, $1.83 \pm 0.09\%$, 28.79 ± 0.13 mg/100g and 7.63 ± 0.38 mg/100g, respectively (Table 3). Total soluble protein, total starch content, total ash content as well as minerals viz. Phosphorus and iron were found to be in similar range in both raw and colorant mixed final product of *Colocasia esculenta* (L.) Schott. (variety: Ahina) (Table 3). The total phenol content and total anthocyanin content of the final food product was found to be higher than raw *Colocasia esculenta* (L.) Schott. (variety: Ahina). The final product isolated from *S. Cumini* L., *S. fruticosum* Roxb. DC., and *Rosa indica* L. found with increased total phenol content of 71.83 ± 0.08 mg GAE/100g, 49.42 ± 0.14 mg GAE/100g, and 69.54 ± 0.35 mg GAE/100g. In comparison to 0.84 mg/100g in raw *Colocasia esculenta* (L.) Schott. (variety: Ahina) powder, total anthocyanin concentration in final product blended with colorants was found to be 44.76 ± 0.70 mg/100g (*S. Cumini* L.), 35.66 ± 0.67 mg/100g (*S. fruticosum* Roxb. DC.) and 48.71 ± 0.39 mg/100g (*Rosa indica* L.) (Table 4).

4. CONCLUSION

Bioavailability and nutritional diversity in colorants of different indigenous and wild plant

sections have been indicated in previous publications. On the basis of chemical attributions, the source of colorants varies from plant to plant. In terms of food coloring, anthocyanin extraction and stability are key concerns. Although the majority of the colorants are phenolic compounds, extraction in an aqueous medium yields the highest extraction percentage for food usage. The total anthocyanin concentration of the fruit extracts of *S. cumini* L. was found to be the highest among seven plant samples, with a value of 124.87 ± 0.078 mg/100g, followed by *R. indica* L. (115.26 ± 0.091 mg/100g) and *S. fruticosum* Roxb. DC. (91.41 ± 0.097 mg/100g). High total anthocyanin concentration suggested a greater stability of total anthocyanin extracted from plant samples in the acidic range up to pH 4. Best stability was obtained with temperature treatment of 40°C for 6 hours. 2,500 lux light intensity applied up 2 hours of incubation resulted in lowest degradation of anthocyanin content. The starch content of *Colocasia esculenta* (L.) Schott. (variety: Ahina) in the final food product was unaffected by the addition of colorants. The total anthocyanin content of final products was determined to be 39.42 mg/100g (*S. cumini* L.), 31.33 mg/100g (*S. fruticosum* Roxb. DC.) and 44.51 mg/100g (*Rosa indica* L.) after one month of refrigerated storage. Similarly, enriched Assam lemon beverage with extracted colorants results in better stability and stable pH for color optimization.

ACKNOWLEDGEMENTS

This study is in part of Mr. Simanta Das's M.Sc. thesis and was funded by "Assam Agricultural University, Jorhat, Assam". Mr. Simanta Das would like to thank Department of Biochemistry and Department of Biotechnology, Assam Agricultural University, Assam, India for their support during his M.Sc. studies.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Miguel MG and Albano SM. Biological activities of extracts of plants grown in Portugal. *Industrial Crops and Products*. 2011;33:338-343.
2. Abdulrahman FI, Inyang I, Abbah, J, John AL, Amos S. & Gamaniel K. Effect of aqueous leaf extract of *Irvingia gabonensis* on gastrointestinal tract in rodents. *Indian J. Exp. Biol.*2004; 42:787-91.
3. Singleton VL & Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am J. Enol. Vitic.*1965;16:144-158.
4. Chia-Chi Chang, Ming-Hua, Y., Hwei-Mei, W. & Jiing-Chuan, C. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J. Food Drug Anal.*2002;10(3): 178-182.
5. Lee J, Durst R, Wrolstad R. Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants, and wines by the pH differential method: Collaborative Study. *J. of AOAC Int.* 2005;88:1269-1278.
6. Selim KA, Khalil KE, Abdel-Bary MS & Abdel-Azeim N.A. Extraction, encapsulation and utilization of red pigments from Roselle (*Hibiscus sabdariffa* L.) as natural food colorants. In: 5th Alexandria Conference of Food and Dairy Science and Technology, Alexandria, Egypt. 2018;7-20.
7. Amr AK and Al-Tamimi EK. Stability of the crude extracts of *Ranunculus asiaticus* anthocyanins and their use as food colourants. *Int. J Food Sci. Tech.* 2007;42(8):985 – 991.
8. Amr AK & Al-Tamimi EK. Stability of the crude extracts of *Ranunculus asiaticus*anthocyanins and their use as food colorants. *Int. J Food Sci. Tech.* 2007;42(8):985–991.
9. Hazarika I. Biochemical characterization of some *Colocasia (Colocasia esculenta L.)* germplasm. M.Sc. (Biochemistry and Agricultural Chemistry) thesis, Assam Agricultural University, Jorhat, Assam 2016.
10. Chitnis N, Pemberton P & Yukich J. Theory of reactive interventions in the elimination and control of malaria. *Malar J.* 2019;18:266-269.
11. Sadasivam S, Manickam A. *Biochemical methods for agricultural sciences*. Wiley Eastern Ltd., New Delhi; 1992.
12. AOAC official methods of analysis. 10th Ed. Association of Analytical Chemists, Washington D.C; 1970.
13. AOAC official methods of analysis, 13th Ed. Association of Official Analytical Chemists, Washington, D.C; 1980.
14. ISO 3534-3. Statistics — Vocabulary and symbols — Part 3: Design of experiments; 1985.
15. ISO 5725-6. Accuracy (trueness and precision) of measurement methods and results — Part 6: Use in practice of accuracy values; 1994
16. Kumari P. Isolation and characterization of anthocyanin pigments from Indian Rose varieties as a potential source of nutraceuticals. Division of floriculture and landscaping. ICAR - Indian Agricultural Research Institute New Delhi 110012; 2016. Available:<http://krishikosh.egranth.ac.in/handle/1/5810069960>.2016
17. Devi P, Saravanakumar M, Mohandas S. African the effects of temperature and pH on stability of anthocyanins from rd sorghum (*Sorghum bicolor*) bran. *J. of Food Sci.* 2012;6(24): 567-573.
18. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*. Indian Council of Agricultural Research Publication.1985;87-89.
19. Ameya G, Tsalla T, Getu F. Antimicrobial susceptibility pattern, and associated factors of *Salmonella* and *Shigella* infections among under five children in Arba Minch, South Ethiopia. *Ann. Clin. Microbiol. Antimicrob.* 2018;17:1. Available:<https://doi.org/10.1186/s12941-018-0253-1>.

20. Debnath I. Phytochemical analysis and antimicrobial activity of Aparajita (*Clitoria ternatea* L.) against rice pathogens. M.Sc. (Biochemistry and Agricultural Chemistry) thesis, Assam Agricultural University, Jorhat, Assam; 2016.
21. El-Gharably AMA.Characterization of anthocyanin pigments extracted from grape skins and its potential uses as antioxidant and natural food colorants. J. Home Econ. 2005;15:51–70
22. Ghosh P, Pradhan RC, Mishra S, Patel AS & Kar A. Physicochemical and nutritional characterization of Jamun (*Syzygium cumini*). Curr. Res. Nutr. Food Sci. 2017; 5(1):25-35.
23. Ramalingam R, Palmurugan S & Shanmugapriya D. Effect of temperature, light, pH on the stability of anthocyanin pigments in *Cocculus hirsutus* fruits. Int. J. Multidiscip. Res. Mod. Educ. 2016;2(2):91-96.
24. Roobha J, Marappan S, Aravindhana KM, Devi PS. The effect of light, temperature, pH on stability of anthocyanin pigments in *Musa acuminata* bract. Res. Plant Biol. 2011;1:5-12.

© 2023 Das et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104104>