



Changes in Physico-Chemical Properties of Mango Pickle during Storage

Hatneikim Haokip^{a*}, Tabuiliu Abonmai^b and Alok Singh^a

^a *Department of Horticulture, College of Agriculture, Central Agricultural University, Imphal-795004, Manipur, India.*

^b *Department of Agronomy, College of Agriculture, Central Agricultural University, Imphal-795004, Manipur, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i232499

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91353>

Original Research Article

Received 25 July 2022
Accepted 12 September 2022
Published 28 October 2022

ABSTRACT

The present study was carried out to investigate the effect of recipes and varieties of Mango (*Mangifera indica* L.) pickle on changes in physico-chemical properties during storage at the Post Graduate Laboratory of the Department of Horticulture, College of Agriculture, Central Agricultural University, Imphal during 2020-2021. The experiment was laid out in factorial randomized block design (FRBD) with eight treatments and three replications. The treatments consist of four pickled recipes and two varieties. Among all pickles, the salty pickle was found to have maximum pH whereas, the sweet mango pickle was found to be minimum. The TSS increases continuously during storage irrespective of the recipe and variety used. Sweet pickle showed maximum TSS. Among all pickles, oily mango pickle was found to have maximum moisture content whereas, salty mango pickle was found to be minimum. Sweet mango pickle has the highest score and is organoleptically more acceptable among other types of a pickle. The findings also showed that with the advancement of the storage period organoleptic quality improved in all mango pickles. The local mango was also found to be the better variety for pickling. It can be concluded that mango of the local kind was better suited for pickling.

Keywords: *Mango; pickle; pH; TSS; moisture; chlorophyll; acceptability.*

*Corresponding author: E-mail: hatneikim.hk@gmail.com;

1. INTRODUCTION

Mango (*Mangifera indica* L.) belongs to the family Anacardiaceae and originated in the Indo-Burma region. It is one of the choicest fruits in the world [1] and is also one of the most preferential fruit crops of the tropical and subtropical regions of the world for human consumption [2]. Due to its taste, importance and luscious flavour, *M. indica* is often named the “King of fruits”. Its social and economic impact is most relevant in developing and emerging countries, where mango is a high-valued component in the diet, rich in vitamins and minerals [3].

Mango fruits are usually eaten fresh because of their succulence, exotic flavour and delicious taste. Nutritionally, mango is a good source of bioactive compounds such as provitamin A, vitamin C, carbohydrates, organic acids, minerals, anti-oxidants, fibres and proteins [4]. Raw fruits of local varieties of mango trees are used for preparing various traditional products like raw slices in brine, amchur, pickle, murabba, chutney, sherbet, etc. Pickling, the commonly used method of preservation began as a way to preserve food for out of season use and for long journeys. Mango pickle is prepared by using a fully mature raw mango. It is also one of the oldest and most successful methods of food preservation known to humans. So far, not much research has been done in relation to mango pickling and its standardization of recipe in the Northeast region especially in Manipur. Considering the above factors, the study entitled “Effect of Recipes and Varieties on Mango (*Mangifera indica* L.) Pickle” was done. Considering the above factors, this study was done to investigate the effect of recipes and varieties of Mango (*Mangifera indica* L.) pickle on changes in physico-chemical properties during storage.

2. MATERIALS AND METHODS

Immature mango fruits of Totapuri and Local varieties free from blemishes, any visible disease infestation and physical injury were collected from Horticulture Research Farm, Central Agricultural University, Andro, Imphal East. Other

materials viz., chemicals and spices were procured from the local market of Imphal, Manipur. The mango fruits were washed with clean water and wiped dry with a muslin cloth. It was then cut into pieces of convenient sizes removing the stones. The mango pieces were mixed with salt and filled in large jars and kept for a week at room temperature, shaking it occasionally. It was then removed from the jar and excess brine solution drained off and dried in the air followed by pickle preparation. The salt concentration was 20 per cent for salty mango pickle (P1) and 15 per cent salt for sweet (P2), oily (P3) and vinegar (P4) mango pickles. The experiment included the combination of 2 mango varieties and 4 mango pickle recipes. The details of which are given below:

(a) Variety: 2

V1 - Local
V2 - Totapuri

(b) Recipe: 4

P1 - Salty mango pickle
P2 - Sweet mango pickle
P3 - Oily mango pickle
P4 - Vinegar mango pickle

2.1 Preparation of Mango Pickle

Four recipes for mango pickles were selected for standardization. Variations were made in the contents of each recipe. Uniform size slices were made from the composite samples of brined mango pieces which were taken with varying amounts of other ingredients for the preparation of pickles in these recipes. The following recipes as presented in Table 1 were used for the preparation of mango pickles. Fig. 1 represented the steps involved during the preparation of mango pickles. The resulting pickles were filled in glass jars and stored at room temperature for use in the estimation of the changes in various quality parameters. Pickle prepared by using different recipes were analysed at 0, 30, 60 and 90 days of storage for their chemical properties i.e., total titratable acidity [5], total sugar [6], ash content [5], microbial evaluation [5] and organoleptic quality [7].




Fig. 1. Flowsheet for preparation of mango pickle

Table 1. Different recipes used for preparation of mango pickle

Ingredients	Recipes			
	P ₁	P ₂	P ₃	P ₄
Unripe mango pieces	1 kg	1 kg	1 kg	1 kg
Turmeric	20 g	20 g	20 g	20 g
Mustard seed	50 g	50 g	50 g	50 g
Black pepper	5 g	5 g	5 g	5 g
Fenugreek	40 g	40 g	40 g	40 g
Red chilli powder	20 g	20 g	20 g	20 g
Fennel seeds	10 g	10 g	10 g	10 g
Cumin seed	20 g	20 g	20 g	20 g
Salt	200g	150g	150g	150g
Jaggery	-	500 g	-	-
Mustard oil	-	-	500 ml	250 ml
Vinegar	-	-	-	100 ml

Table 2. Effect of different recipes and varieties on pH, TSS (⁰B), moisture content (%) and chlorophyll content (mg/g) of mango pickle at different storage days

Treatment	pH				TSS (⁰ B)				Moisture content (%)				Chlorophyll content (mg/g)			
	0	30	60	90	0	30	60	90	0	30	60	90	0	30	60	90
Recipes																
P1	2.41	2.28	2.16	2.14	8.43	9.53	11.17	12.45	0.61	0.99	1.07	1.34	1.28	1.23	1.17	1.13
P2	2.17	2.11	2.01	1.94	10.27	11.45	13.60	14.57	0.56	0.72	0.90	1.28	1.44	1.37	1.22	1.16
P3	2.30	2.21	2.09	2.06	8.03	9.30	10.98	12.22	0.42	0.56	0.79	1.07	1.24	1.21	1.16	1.12
P4	2.25	2.19	2.06	2.01	8.82	10.00	12.02	12.85	0.48	0.64	0.85	1.17	1.31	1.26	1.19	1.15
S.E(d) (±)	0.02	0.02	0.02	0.03	0.11	0.07	0.07	0.09	0.01	0.03	0.01	0.02	0.01	0.01	0.00	0.00
C.D	0.04	0.05	0.05	0.06	0.24	0.15	0.15	0.20	0.03	0.07	0.02	0.04	0.01	0.02	0.01	0.01
Varieties																
V1	2.20	2.10	1.93	1.87	9.63	10.62	12.58	13.24	0.55	0.79	0.96	1.25	1.44	1.39	1.27	1.22
V2	2.36	2.29	2.23	2.20	8.14	9.53	11.30	12.80	0.49	0.67	0.85	1.19	1.20	1.15	1.10	1.06
S.E(d) (±)	0.01	0.02	0.02	0.02	0.08	0.05	0.05	0.07	0.01	0.02	0.01	0.01	0.00	0.01	0.00	0.00
C.D	0.03	0.04	0.03	0.04	0.17	0.10	0.11	0.14	0.02	0.05	0.02	0.03	0.01	0.01	0.00	0.00

Table 3. Effect of different recipes and varieties on flavour, texture and overall acceptability of mango pickle at different storage days

Treatment	9-point Hedonic rating											
	Flavour				Texture				Overall acceptability			
	0	30	60	90	0	30	60	90	0	30	60	90
Recipes												
P1	6.75	7.17	7.25	7.58	6.58	7.25	7.42	7.75	7.17	7.25	7.58	7.58
P2	7.83	8.08	8.25	8.67	7.25	7.92	8.08	8.25	7.92	8.42	8.58	8.75
P3	7.17	7.50	7.50	7.92	7.58	8.25	8.42	8.75	7.42	7.58	7.92	8.08
P4	7.58	7.75	7.92	8.33	6.92	7.42	8.00	8.08	7.75	7.92	8.25	8.42
S.E(d) (\pm)	0.30	0.25	0.28	0.30	0.21	0.29	0.21	0.24	0.21	0.22	0.22	0.24
C.D	0.60	0.51	0.57	0.22	0.43	0.58	0.19	0.48	0.42	0.45	0.44	0.20
Varieties												
V1	7.46	7.75	7.92	8.21	7.17	7.79	8.04	8.29	7.67	7.88	8.17	8.29
V2	7.21	7.50	7.54	8.04	7.00	7.63	7.92	8.13	7.46	7.71	8.00	8.13
S.E(d) (\pm)	0.21	0.18	0.20	0.21	0.15	0.20	0.15	0.17	0.15	0.16	0.15	0.17
C.D	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

3. RESULTS AND DISCUSSION

The pH value of the pickles decreases as the storage period increases in all the recipes and mango varieties. A decrease in pH can be attributed to increasing in acidity. Acidity increased in pickles due to lactic acid fermentation. Fermentation has been known to decrease pH and increase titratable acidity in several foods. Similar results of decrease in pH with storage period were also reported by Saroj and Singh [8] in mango pickles and Sharma [9] in lime pickles. There was a significant increase in TSS during 90 days of storage period irrespective of recipe and variety used. An increase in the TSS value of pickles may be possible due to the jaggery, salt and other ingredients being added and mixed well during the storage period. Increase in TSS could also be due to loss of moisture. A similar trend was reported by Singh et al. [10] in cauliflower pickles. The study revealed a considerable decrease in moisture content of all mango pickles throughout the storage period. The reduction in moisture content was due to adding of more salt and sugar which causes osmosis. So, the salty pickle showed a considerable decrease in moisture content showed a maximum decrease in moisture which might be due to its already dryness. Similar results were observed by Sharma [9] in lime pickles and Saroj and Singh [8] in mango pickles. There was a general decreasing trend in chlorophyll content of mango pickles during storage. Further, while comparing the overall effect of varieties it was retained higher in local variety and lesser in Totapuri mango during storage. The decrease in chlorophyll content of pickles might also be due to its photo-oxidation in transparent containers during storage. The same trend was observed in seedling mango pickle by Thakur et al. [11].

Data regarding the organoleptic quality of pickles revealed that the overall acceptability of pickles increased significantly with the increase in storage period. Flavour and texture showed improvement in their quality. Sweet pickles obtained the highest score with respect to their overall acceptability than that of other recipes. Mango oil pickle and vinegar pickle also scored higher which might be due to their gravy formation and salty pickle scored less due to its dryness. The increase in organoleptic quality in pickles throughout the storage period can be attributed to the conversion of the insoluble fraction of pectin to soluble form which might have resulted in softening of the pickle. Similar

results were obtained by Sharma [9] in lime pickles and Rekha (2004) in kachri pickles. The interaction between the recipes and mango variety could not show a significant effect on the mango pickle.

4. CONCLUSION

It can be concluded that the local variety mango was more suitable for pickle making due to low pH, lower moisture content and higher overall organoleptic scores than Totapuri. Among the recipes sweet mango pickle scored the highest organoleptic scores and is the best recipe for making pickles. The texture obtained out of oily mango pickles was also very well received. The organoleptic quality of pickles improved with the increase in storage period in all the recipes.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Joshi R, Kundu M, Singh CP. Efficient tool for identification on different mango cultivars. *Environ. Ecol.* 2013;31(1):385-388.
2. Vasugi C, Dinesh MR, Sekar K, Shivashankara KS, Padmakar B, Ravishankar KV. Genetic diversity in unique indigenous mango accessions (Apemidi) of the Western Ghats for certain fruit characteristics. *Curr. Sci.* 2012; 103(2):199-207.
3. Ribeiro RSM, Queiroz JH, Queiroz LRME, de Campos FM, Santana PHM. Antioxidant in mango (*Mangifera indica* L.) pulp. *Plant Foods Hum. Nutr.* 2007;6:13-21.
4. Masibo M, He Q. *Food Reviews International.* 2009;25:346-370.
5. Ranganna S. *Handbook of analysis and quality control for fruit and vegetable products.* 2nd Edn., Tata McGraw-Hill Publishing Company Limited, New Delhi, India; 2007.
6. Yemm EW, Willis AJ. The estimation of CH₂O₂ in plant extracts anthrone. *Biochem. J.* 1954;57:508-517.
7. Amerine MA, Pangborn RM, Roessler EB. *Principles of sensory evaluation of food.* Academic Press: London. 1965;236-268.
8. Saroj, Krishan KS. Standardization of recipe and method for mango pickle. *Int. J. Chem. Stud.* 2018;6(2):2033-2037.

9. Sharma A. Studies on preservation of products of sour lime (*Citrus aurantifolia* Swingle). M.Sc. Thesis, CCS Haryana University, Hisar; 2002.
10. Singh SN, Patel VA, Varma MG, Patel SG, Savaliya MV. Effect of processing methods on physico-chemical properties, sensory evaluations and storage studies of cauliflower (*Brassica oleracea* L. var. botrytis) Pickle. Int. J. Curr. Microbiol. Appl. Sci. 2021;10(01):1413-1428.
11. Thakur NS, Thakur A, Chauhan M, Kumar P. Studies on Development and Storage Quality of Seedling Mango (*Mangifera indica* L.) Pickle of Himachal Pradesh. Indian J. Ecol. 2017;44:690-696.

© 2022 Haokip et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91353>