



Varietal Evaluation of Hybrid Cabbage (*Brassica oleraceae* var. capitata L.) under Prayagraj Agro-climatic Conditions

R. Y. Ashik^{a*}, Vijay Bahadur^a, Tarence Thomas^a and V. M. Prasad^a

^a *Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj - 211007, Uttar Pradesh, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2131342

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89983>

Original Research Article

Received 22 May 2022
Accepted 27 July 2022
Published 29 July 2022

ABSTRACT

An experiment on cabbage was conducted during October (2021) to February (2022), in horticulture Research field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, and Technology and Sciences Prayagraj (U.P) India. The results of investigation, regarding the Evaluation of the 12 varieties of Cabbage *i.e.*, V1 (Cabbage F1 Hybrid NBH- Arun) , V2 (Cabbage F1 Hybrid royal vantage), V3 (Hybrid cabbage Kashi (22)), V4 (Cabbage NS 22), V5 (Green Hero), V6 (F1- Hybrid cabbage fieldman), V7(J K Green King F1-Hybrid Seed), V8 (Green Top- F1), V9(Hybrid Gayatri), V10(Cabbage- Bio Samrat), V11(Cabbage- F1 Lucky Ball) and V12(F1 Hybrid Indam-296) obtained from different sources to find out the best performance in terms of growth and yield in Prayagraj agro climatic conditions. The experiment was conducted in Randomized Block Design, were each hybrid replicated thrice the results from the present investigation Concluded that Cabbage varieties JK Green king F1-Hybrid was recorded with minimum days for germination (8.50), plant height (32.72), Number of leaves (16.92), Plant spread (2633.47), Marketable Head Yield (1571.95), Net Head Yield (1324.18), Head Yield (t/ha) (256.16), TSS (8.61), Ascorbic Acid (29.66), Fiber content (16.16), Head equatorial diameter (10.98), Head polar Diameter (12.71), Head Compactness (37.47) and with maximum Benefit cost ratio (3.64).

Keywords: Cabbage; varieties; agro-climatic conditions.

1. INTRODUCTION

Cabbage (*Brassica oleraceae* var. capitata) is one of the most important leafy vegetables worldwide [1]. It originated from a wild non heading type, 'cole wart' (*Brassica Oleracea* var. sylvestris) and considered the real headed cabbage evolved in Germany. Cabbage belongs to the family of Brassicaceae and genus of Brassica. Cabbage (*Brassica Oleracea* var. Capitata) is one of the most popular vegetables in the world because of its adaptability to a wide range of climatic conditions and soil types, ease of production and storage, and its food value. It is believed to have evolved from a wild form native to Europe, growing along the coast of the North Sea, the English Channel and northern Mediterranean. Cabbage has been ranked by the Food and Agriculture Organization among the top twenty vegetable crops grown, establishing it as an important food source, globally [2]. Area and production of cabbage was 397 ha and 9207 MT respectively (NHB, 2020) Cabbage is a cool season crop which grows best under cool, moist weather conditions [3]. The cabbage is a crop plant that is easily grown on a wide range of soil types and is adaptable to many different climatic conditions (Smith, 1995). It can be cultivated throughout the year if proper varieties are selected and planted. It favors in cold climate and soil pH with 5.5 to 6.5. Cabbage yield and yield characteristics are determined by several factors, such as variety, plant spacing, environmental conditions and market requirements, among others. Days to maturity, or earliness, in cabbage is largely determined by variety, and is a desirable characteristic when varieties with short growth cycles are required to meet market demands [4] Commercial cultivation of cabbage is very successful due to high market demand. In the recent years, hybrid cabbage cultivation is increasing. Cabbage is a good source of vitamin C and sulphur. Cabbage helps with constipation. Cabbage helps with gastric ulcers. There is a wide scope of increasing cabbage production with the introduction of new suitable cultivars from abroad. There are many cabbage varieties available in the market, which have been imported by different seed companies. Prior to recommendation for farmers, varietal evaluation needs to be determined. Otherwise farmers will be deprived of getting benefit from cabbage cultivation. The objective of this study was to evaluate the yield and growth characteristics of 12 cabbage varieties in order to

determine the best suited to growing conditions, particularly season and location.

2. MATERIALS AND METHODS

The experiment was conducted Central Horticulture Research farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, and Technology and Sciences Prayagraj (U.P) India during 2020-2021. The experiment was laid out in RBD (Randomized Block Design) having 12 varieties with 3 replications of hybrid varieties of cabbage tabulated-1. Cabbage varieties were sown in the field at a spacing of 60 x 45 cm in the plot of 2m x 1m size. The standard recommended dose of fertilizers (200: 120:80 NPK kg/ha + 15 ton FYM/ha) was applied and Normal cultural practices and plant protection measures were followed during the cultivation process.

The data recorded during the course of investigation were subjected to statistical analysis as per method of analysis of variance [5]. The significance and non-significance of the treatment effect were judged with the help of 'F' value (variance ratio) was compared with the table value at 5% level of significance. If calculated value exceeded then the value, the effect of considered to be significant. The significant difference between the means was tested against the critical difference at 5% level of significance.

2.1 Chemical Analysis of Soil

Composite soil samples are collected randomly before the layout of experiment was laid so as to determine the soil properties initially. The soil samples are collected from 0-15 cm depth and were dried under shade, then powdered with the help of a wooden pestle and mortar then sieved through a 2 mm sieve and was then subjected to further analysis. The physical properties of soil were evaluated by using the Bouyoucos hydrometer method outlined by Bouyoucos, (1952) and for organic carbon by Wet method (Walkely and Black, 1956). Available nitrogen (260.14 kg/ha) was estimated by alkaline permanganate method by Subbiah and Asija, [6], available phosphorus (133kg/ha) by Olsen method by Olsen et al., (1954), available potassium (145.23 kg/ha) was determined by use of Flame Photometric method [7].

Table 1. List of hybrid varieties

Sl.No	Varieties	Name of hybrid	Source
1.	V1	Cabbage F1 Hybrid NBH-arun	Noble Pvt.Ltd Seeds
2.	V2	Cabbage F1 Hybrid Royal Vantage	Sakata India (P) Ltd Seed
3.	V3	Hybrid Cabbage Kashi (22)	Akshay Pvt.Ltd Seeds
4.	V4	Cabbagens 22	Namdhari Seeds Pvt.Ltd
5.	V5	Green Hero	Takii-Seeds India Pvt.Ltd
6.	V6	F1-Hybrid Cabbage Fieldman	Kalash Pvt.Ltd Seeds
7.	V7	JK Geen King F1-Hybrid Seed	JK-Agri Genetics Ltd. East-west Seed
8.	V8	Green top-F1	International
9.	V9	Hybrid Gayatri	Doctor Seeds Pvt. Ltd
10.	V10	Cabbage-Biosamrat	Bioseed
11.	V11	Cabbage-F1 lucky Ball	Lucky-Seeds Private Limited
12.	V12	F1 Hybrid Indam-296	Indo-American Hybrid-Seed (India)Pvt.Ltd.

3. RESULTS AND DISCUSSION

3.1 Growth Parameters Days to Germination

The statistically analysed data on Days to Germination of cabbage seeds are represented in table 2. It is clear from the table that there is significant difference among various hybrids of cabbage. The Minimum days to germination was found in V7 (JK green King F1- Hybrid Seed) which was found 8.50 followed by V9 (Hybrid Gayatri) which was 9.00 and the maximum Germination days was found in varieties V6 (F1-Hybrid Cabbage Fieldman) which was 14.50 which are statically at par value respectively. The number of days to germination is an important character. which indicate earliness or lateness of the crop in general. Better germination in this hybrid may be due to genetic potential or having the better permeability of seed coat to water hence earlier initiation of germination. The earlier and late germination helps in the occurrence of early/late flush of crop which is advantageous for market to fetch the higher price. The view was supported by Kamal et al., [8], Narayan et al., [9] and (Chandra Leela et al., 2020) in Bottle gourd. Similar findings also [10] in broccoli.

3.1.1 Plant height

The statistically analysed data on days to Plant Height of cabbage seeds are represented in table 2. It is clear from the table that there is significant difference among various hybrids of

cabbage. The maximum Plant Height was found in varieties V7 (JK green King F1- Hybrid Seed) which was found 26.22 followed by V₉ (Hybrid Gayatri) which was 25.58, V₁₂ (F1 Hybrid Indam-296) (24.19) which are statically at par value and minimum was found in V6 (F1- Hybrid Cabbage Fieldman) which was 19.35 in 30 DAT. The maximum plant height was found in varieties V7 (JK green King F1- Hybrid Seed) 31.07 followed by V9 (Hybrid Gayatri) which was 30.26, V12 (F1 Hybrid Indam-296) (30.17), V5 (Green Hero) (29.42) which are statically at par value and minimum was found in V6 F1- Hybrid Cabbage Fieldman which was 27.72 in 60 DAT. The maximum plant height was found in varieties V7 (JK green King F1- Hybrid Seed) (32.72) followed by V9 (Hybrid Gayatri) which was 32.56, V12 (F1 Hybrid Indam-296) (31.81), V₅ (Green Hero) (31.88) which are statically at par value and minimum was found in V₆ F1- Hybrid Cabbage Fieldman which was 16.29 and minimum was found in V6 F1- Hybrid Cabbage Fieldman which was 30.68 in 90 DAT respectively. This variation in plant height could be due to variation in the genetic make- up of different varieties. Environmental conditions caused variation in the hormonal balance and cell division rate that result in changes in the plant height of the different varieties. The results of this study are in agreement with Gaurav, Srivastava., et al., [11].

3.1.2 Number of leaves

The statistically analysed data on days to Number of Leaves of cabbage seeds are

represented in Table 2. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum number of leaves was at 30 DAT found in varieties V7 (JK green King F1- Hybrid Seed) which was found 8.59 followed by V9 (Hybrid Gayatri) (8.29) which are statically at par value and minimum plant spread found in V6 (F1- Hybrid Cabbage Fieldman) which was 5.44 respectively. The maximum number of leaves was at 60 DAT found in varieties V7 (JK green King F1- Hybrid Seed) which was 14.63 and minimum number of leaves V6 (F1- Hybrid Cabbage Fieldman) which was 9.96 respectively. The maximum number of leaves was at 90 DAT found in varieties V7 (JK green King F1- Hybrid Seed) which was 16.92 and minimum was found in V6 F1- Hybrid Cabbage Fieldman which was 16.29 respectively. The variation in number of leaves per plant under different varieties, might be due to differences in their genetic inherent capacity, and suitability under this climate. The lower number of leaves in some cultivars was probably due to slow rate in leaf initiation. The mean Number of leaves of varieties of cabbage was found significant at different stages. Similar findings also reported by Thapa and Rai, [12] in broccoli.

3.1.3 Plant spread

The statistically analysed data on days to Plant Spread of cabbage seeds are represented in Table 2. It is clear from the table that there is

significant difference among various hybrids of cabbage. The maximum plant spread in 30 DAT was found in V7 (JK green King F1- Hybrid Seed) which was found 1040.45 which are statically at par value and minimum plant spread found in V6 (F1- Hybrid Cabbage Fieldman) which was 594.41 respectively. The maximum plant spread in 60 DAT was found in V7 (JK green King F1- Hybrid Seed) which was found 1873.67 which are statically at par value and minimum plant spread found in V6 (F1- Hybrid Cabbage Fieldman) which was 1126.42 respectively. The maximum Plant Spread in 90 DAT was found in varieties V7 (JK green King F1- Hybrid Seed) which was found 2633.47 followed by V9 (Hybrid Gayatri) which was 2632.33, V12 (F1 Hybrid Indam-296) (2607.89), V5 (Green Hero) (2604.35), V3 (Hybrid Cabbage kasha) (2526.32), V4 (Cabbage NS 22) (2414.83), V1 (Cabbage F1 Hybrid NBH-ARUN) (2333.55) which are also statically at par value and minimum plant spread was found in V6 (F1- Hybrid Cabbage Fieldman) which was 773.20 90 DAT respectively. Because of the capacity of genotype to intake more water and ability of the genotype to grow fast in the condition like sunlight, moisture, temperature and the number of days to second true leaf emergence indicate earliness or lateness of flowering. Based on the ability of genotype which induces both cell division and cell elongation [13], [14] and [8]. similarly, it can stimulate plant tissue results in enhanced vegetative growth.

Table 2. Performance of cabbage varieties in terms of days to germination, plant height (cm), number of leaves, plant spread (cm)

Varieties	Days to germination	Plant height (cm)	Number of leaves	Plant spread (cm)
V1	10.17	30.68	14.51	2333.55
V2	11.67	28.13	13.29	2117.46
V3	9.83	31.50	13.44	2526.32
V4	10.00	30.88	15.74	2414.83
V5	9.67	31.72	15.77	2604.35
V6	14.50	30.07	16.29	2002.75
V7	8.50	32.72	16.92	2633.47
V8	13.00	29.96	14.37	2037.18
V9	9.00	32.56	17.59	2632.33
V10	13.50	28.52	16.03	2017.73
V11	11.50	30.35	15.14	2165.22
V12	9.50	31.81	14.92	2607.89
F test	S	S	S	S
S. Ed (\pm)	0.40	0.53	0.39	2.05
C.D at 5%	4.53	2.11	3.17	0.10
C.V.	0.84	1.10	0.82	4.27

3.2 Yield Parameters

3.2.1 Marketable head yield

The statistically analysed data on days to Marketable Head Yield of cabbage seeds are represented in Table 3. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum Marketable yield was found in varieties V7 (JK green King F1- Hybrid Seed) which was found 11571.95 and minimum was found in V6 F1- Hybrid Cabbage Fieldman which was 773.20 90 DAT respectively. The yield attributes are significantly influenced by genetic potential of the plant attributed to higher biomass accumulation coupled with effective translocation and distribution of photosynthates from source to sink, which in turn resulted into elevated stature of yield attributes, which of course was due to favorable weather conditions such as rainfall distribution, evaporation and relative humidity prevailed during the crop growth period. Similar findings of the experiment was concluded by Surendra Lal Shrestha, [15] in cabbage.

3.2.2 Net head yield

The statistically analysed data on days to Net Head Yield of cabbage seeds are represented in table 3. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum Net Head yield was found in varieties V7 (JK green King F1- Hybrid Seed) which was found 1324.18 and minimum was found in V6 F1- Hybrid Cabbage Fieldman which was 524.59 at 90 DAT respectively. The yield attributes are significantly influenced by genetic potential of the plant attributed to higher biomass accumulation coupled with effective translocation and distribution of photosynthates from source to sink, which in turn resulted into elevated stature of yield attributes, which of course was due to favourable weather conditions such as rainfall distribution, evaporation and relative humidity prevailed during the crop growth period. Similar findings of the experiment was concluded by Surendra Lal Shrestha, [15] in cabbage.

3.2.3 Head yield

The statistically analysed data on days to Head Yield of cabbage seeds are represented in Table 3. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum Head yield was found in varieties V7 (JK green King F1- Hybrid Seed) which was

found 256.16 followed by V9 (Hybrid Gayatri) which was 255.66 which are also statically at par value and minimum was found in V6 F1- Hybrid Cabbage Fieldman which was 235.83 at 90 DAT respectively. The higher yield plant⁻¹ (kg) due to its inherent characteristics, better adoptability for the environmental conditions and efficiently all available factors viz. water, nutrient, light and CO₂. None of the treatments significantly influenced the plant stand. This indicates that whatsoever differences were observed in growth and yield attributes were mainly due to different hybrid effect. The results are in agreement with the finding of Kamal et al., [8], Shinde et al., [16] and Deepthi et al., [17] in Bottle gourd.

3.3 Quality Parameters

3.3.1 TSS brix

The statistically analysed data on days to TSS Brix of cabbage seeds are represented in Table 4. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum TSS Brix was found in varieties V7 (JK green King F1- Hybrid Seed) which was found 8.61 followed by V9 (Hybrid Gayatri) which was 8.34, V12 (F1 Hybrid Indam-296) (8.27), V5 (Green Hero) (8.22), V3 (Hybrid Cabbage kasha) (8.17), V4 (Cabbage NS 22) (8.10), V1 (Cabbage F1 Hybrid NBH-ARUN) (8.07), V11 (Cabbage-F1 Lucky Ball) (8.03) which are also statically at par value and minimum was found in V6 F1- Hybrid Cabbage Fieldman which was 7.55 at 90 DAT respectively. Increase of protein content is due to the seed inoculation increased the N content of pods which ultimately reflects TSS content. Also, the higher TSS value is due to the inherent characteristics of the hybrid and is due to favorable influence on various metabolic processes like photosynthesis, respiration, enzyme activity [18] which augments the production of metabolites and their translocation to different parts including seed which ultimately increases the concentration of nutrients in seed. The mean TSS Brix of varieties of cabbage was found significant at different stages. Similar findings were found by Madhuri, [19] in watermelon.

3.3.2 Ascorbic acid (mg/100g)

The statistically analysed data on days to Ascorbic Acid of cabbage seeds are represented in Table 4. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum Ascorbic Acid

Table 3. Performance of cabbage varieties in terms of Yield parameters

Varieties	Marketable head yield	Head polar diameter	Head equatorial diameter	Net head weight	Head yield
V1	1156.34	9.98	10.44	715.62	249.83
V2	1043.66	9.8	10.2	677.52	245
V3	1234.93	10.19	10.61	824.81	252.5
V4	1213.15	10.03	10.7	764.86	251.16
V5	1236.93	11.21	10.64	833.70	252.63
V6	773.20	9.16	8.41	524.59	235.83
V7	1571.95	12.71	10.98	1324.18	256.16
V8	953.43	9.52	9.79	661.95	242.16
V9	1336.44	12.67	10.87	917.14	255.66
V10	844.34	9.35	8.59	633.26	239
V11	1115.56	9.93	10.24	704.70	245.33
V12	1311.76	12.21	10.83	916.67	253
F test	S	S	S	S	S
S. Ed (\pm)	2.38	0.37	0.40	2.44	1.30
C.D at 5%	0.25	4.29	4.81	0.38	0.64
C.V.	4.95	0.76	0.83	5.07	2.70

(mg/100g) was found in varieties V7 (JK green King F1- Hybrid Seed) which was 29.66 followed by V9 (Hybrid Gayatri) which was 29.25, V12 (F1 Hybrid Indam-296) 28.86, V5 (Green Hero) (28.32), V3(Hybrid Cabbage kasha) (28.05), V4(Cabbage NS 22) (27.54) which are also statically at par value and minimum was found in V6 F1- Hybrid Cabbage Fieldman which was 21.03 at 90 DAT respectively. It was probably caused by higher sunlight intensity and also is due to favorable influence on various metabolic processes like photosynthesis, respiration, enzyme activity [18] which augments the production of metabolites and their translocation to different parts including seed which ultimately increases the concentration of nutrients in seed and stover. Positive relationship between light and ascorbic acid level in the plants was also stated. Similar findings were found by Lee and Kader, [20]. The mean Ascorbic Acid (mg/100g) of varieties of cabbage was found significant at different stages.

3.3.3 Fiber content

The statistically analysed data on days to Fiber Content of cabbage seeds are represented in Table 4. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum Fiber Content was found in the varieties V7 (JK green King F1- Hybrid Seed) which was 16.16 followed by V9 (Hybrid Gayatri) which was 15.76, V12 (F1 Hybrid Indam-296) 15.73, V5 (Green Hero) (15.69) which are also statically at par value and

the minimum Fiber Content was found in V6 F1- Hybrid Cabbage Fieldman which was 9.03. This could be due to favorable influence on various metabolic processes like photosynthesis, respiration, enzyme activity [18]. These significant differences with respect to Fiber content among different varieties is due to own genetic makeup and the suitability of varieties to the weather conditions of this zone. Similar findings also reported by El-Bassiony et al., (2014) in Khol-khol, Bhangre et al., [21] in broccoli.

3.3.4 Head equatorial diameter

The statistically analysed data on days to Head Equatorial Diameter of cabbage seeds are represented in Table 4. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum Head Equatorial Diameter was found in the varieties V7 JK green King F1- Hybrid Seed which was 10.98 followed by V9 (Hybrid Gayatri) which was 10.87, V12 (F1 Hybrid Indam-296) 10.83, V5 (Green Hero) (10.64), V3(Hybrid Cabbage kasha) (10.70), V4(Cabbage NS 22) (10.61), V1(Cabbage F1 Hybrid NBH-ARUN) (10.44), V11(Cabbage-F1 Lucky Ball) (10.24), V2(Cabbage F1 Hybrid Royal Vantage) (10.20) which are also statically at par value and the minimum head Equatorial Diameter was found in V6 (F1- Hybrid Cabbage Fieldman) which was 8.41. Increased fruit size attributed in different hybrid genotypes might be due to enhanced photosynthesis, accumulation of carbohydrates

and favourable effect on vegetative growth which increased the fruit variety besides increasing the fruit size. This variation might be due to fruit length, number of fruits per plant and effective node. Similar results have been reported Husnan et al., [22], and Mishra et al., [23] in Bottle gourd.

3.3.5 Head polar diameter

The statistically analysed data on days to Head polar Diameter of cabbage seeds are represented in table 4. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum Head Polar Diameter was found in the varieties V7 JK green King F1- Hybrid Seed which was 12.71 followed by V9 (Hybrid Gayatri) which was 12.67, V12 (F1 Hybrid Indam-296) 12.21 which are also statically at par value and the minimum head Polar Diameter was found in V6 (F1- Hybrid Cabbage Fieldman) which was 9.16. The diameter of fruits may be due to its hybrid vigour and adoptability to Allahabad agro-climatic conditions. Increased fruit size attributed in different hybrid genotypes might be due to enhanced photosynthesis, accumulation of carbohydrates and favourable effect on vegetative growth which increased the fruit variety besides increasing the fruit size. Similar results have been reported Husnan et al., [22], Damor et al., [24], Kumar et al., [25], and Mishra et al., [23] in Bottle gourd [26-36].

3.3.6 Head compactness

The statistically analysed data on days to Head Compactness of cabbage seeds are represented in Table 4. It is clear from the table that there is significant difference among various hybrids of cabbage. The maximum Head compactness was found in the varieties V7 (JK green King F1- Hybrid Seed) which was 37.47 followed by V9 (Hybrid Gayatri) which was 36.17 which are also statically at par value while the minimum head compactness was found in V6 (F1- Hybrid Cabbage Fieldman) which was 29.40. Increased fruit size attributed in different hybrid genotypes might be due to enhanced photosynthesis, accumulation of carbohydrates and favourable effect on vegetative growth which increased the fruit variety besides increasing the fruit size. This variation might be due to fruit length, number of fruits per plant and effective node. Similar results have been reported Husnan et al., [22], and Mishra et al., [23] in Bottle gourd.

3.3.7 Economics

In terms of economics Benefit cost ratio represented in Table 5 was found in the variety V7 JK green King F1- Hybrid Seed which was 3.64 and the minimum Benefit Cost ratio was found in V6 F1- Hybrid Cabbage Fieldman which was 3.35.

Table 4. Performance of cabbage varieties in terms of quality parameters

Varieties	TSS	Ascorbic Acid content	Fiber content	Head compactness
V1	8.07	26.38	13.43	32.30
V2	7.93	26.32	12.33	30.93
V3	8.17	28.05	15.05	33.40
V4	8.10	27.54	14.76	32.70
V5	8.22	28.32	15.69	34.43
V6	7.55	21.03	9.03	29.40
V7	8.61	29.66	16.16	37.47
V8	7.86	21.80	11.88	30.80
V9	8.34	29.25	15.76	36.17
V10	7.84	25.65	11.53	30.13
V11	8.03	26.34	12.8	31.80
V12	8.27	28.86	15.73	35.50
F test	S	S	S	S
S. Ed (\pm)	0.31	1.06	0.53	0.313
C.D at 5%	4.73	4.89	4.76	1.16
C.V.	0.65	2.20	1.10	0.65

In terms of economics Gross Return (INR/ha) represented in Table 5 was found in the variety V7 JK green King F1- Hybrid Seed which was (384240 INR/ha) is maximum and V₆ F1- Hybrid Cabbage Fieldman was found (353745 INR/ha).

Table 5. Performance of cabbage varieties in terms of economics

S No.	Variety	Gross Return (INR/ha)	Net Return (INR/ha)	Benefit Cost Ratio
1.	V1	374745	254843	3.552018
2.	V2	367500	246698	3.483346
3.	V3	378750	251648	3.589979
4.	V4	376740	256118	3.570928
5.	V5	378945	255443	3.591828
6.	V6	353745	231443	3.35297
7.	V7	384240	263138	3.642016
8.	V8	363240	240098	3.442968
9.	V9	383490	259988	3.634907
10.	V10	358500	234998	3.39804
11.	V11	367995	240293	3.488038
12.	V12	379500	254198	3.597088

In terms of economics Net Return (INR/ha) represented in Table 5 was found in the variety V7 JK green King F1- Hybrid Seed which was (263138 INR/ha) is maximum and V6 F1- Hybrid Cabbage Fieldman was found (231443 INR/ha).

4. SUMMARY AND CONCLUSION

The results from my experiment concluded that variety (JK green King F1- Hybrid Seed) found to be best in terms of Growth, Yield and Quality contributing traits along with highest gross return, net return and benefit cost ratio which also gave a significant result for all the characters which were recorded.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Talekar NS. Chinese cabbage. Proceedings of the 1st international symposium on Chinese cabbages. Shanhua, Tainan, Taiwan: Asian vegetable Research Development center. 2000;67-9.
- FAO. Traditional food plants. Food and agricultural organisation of the United Nations, Rome, Italy; 1988.
- Thompson JK. Yield evaluation of cabbage varieties. Agric Technol. 2002;5:15-9.
- Cervenski J, Gvozdanovic-Varga J, Glogovac S. Variance components and correlations of agronomic traits among cabbage (*Brassica oleracea* var. capitata L.) maturity groups. Genetika. 2012; 44(1):55-68.
- DOI: 10.2298/GENSR1201055C.
- Fisher RA. Technique of analysis of variance. Hand book of agriculture statistics. 1960;29-110.
- Subbiah BV, Asija GL. A rapid procedure for the estimate of available nitrogen international journal of Soil. Curr Sci. 1956;25:259-60.
- Toth SJ, Prince AL. Estimate of cation exchange capacity and exchangeable Ca, K, Na, content of soil by flame photometer technique soil. Sciencedirect; 1949;67:439-45.
- Kamal N, Verma S, Agrawal S, Rao SS. Genetic variability and correlation studies in bottle gourd grown as intercrop in coconut garden. Plant Arch. 2012;12(1):85-8.
- Narayan K. Genetic diversity and correlation studies in bottle gourd germplasm under Baster condition. J Agric Sci Chhattisgarh Young Sci Congress. 2013;XI:1.5: 15.
- Gogoi S, Millu R, Das P, Bora N, Das BK. Effect of sowing dates and spacing on broccoli (*Brassica oleracea* var. italica) seed production. Indian J Agric Res. 2016;50(4):350-3.
- Himanshi J. Amarjeet, Bhardwaj. Gaurav, Srivastava, and Mainak, Das. Nano pyrite driven root foraging increases production of the heavy feeders, viz., cauliflower, cabbage and tomato in nutrient deficient soil with no fertiliser application Advanced National Science Nanoscience Nanotechnology. 2019;10:035007:10.
- Thapa U, Rai R. Evaluation of sprouting broccoli (*Brassica oleracea* var. italica). Genotypes for growth, yield and quality. Int J Agric Sci. 2012;4(7):284-6.

13. Jones RL. The physiology of gibberellins induced elongation. In: Skoog F, editor. Plant growth substances. WI, NY: Madison Book Company. 1979;188-95. 14.
14. Jacob M, Hussein S, Alfred O. Correlation and path coefficient analyses of qualitative and quantitative traits in selected bottle gourd landraces. 2016;13. ISSN: 0906-4710.
15. Surendra L. Shrestha. Agric Food Sci (IJHAF). Performance Evaluation of Cabbage (*Brassica oleraceae* Capitata) Cultivars in Mid-Hills of Nepal for winter Season Production. International Journal of Horticulture. 2019;3(2). ISSN: 2456-8635.
16. Shinde RD, Vadodaria JR, Savale SV, Vasava HV. Effect of nature of cultivation and different varieties on flowering, yield attributes and yield of bottle gourd (*Lagenaria siceraria* Mol. Standl.). Trends Biosci. 2014;7(24):4340-5.
17. Deepthi BP, Syam S. Reddy. Satya raj Kumar, A, Ramanjaneya, Reddy. A. Plant Arch. Studies on phenotypic coefficient of variation, genotypic coefficient of variation, heritability and genetic advance in bottle gourd genotypes for yield and yield components. 2016;16(2):597-601. ISSN 0972-5210.
18. Ganie MA, Akhter F, Bhat MA, Malik AR, Junaid JM, Shah MA et al. A critical nutrient element for plant growth and productivity with reference to temperate fruits. Curr Sci. 2013;104(1):76-85. Growth, yield and quality of french bean (*Phaseolus vulgaris* L.) as influenced by sulphur and boron application on inceptisols of Kashmir.
19. Madhuri. performance of watermelon (*Citrullus lanatus*) under different growing conditions; 2016.
20. Lee SK, Kader AA. Pre harvest and postharvest factors influencing vitamin C content of horticultural crops. Postharvest Biol Technol. 2000;20:207-20.
21. Bhangre KK, Sonawane PC, Warade SD. Effect of different varieties and spacing on growth and yield parameters of broccoli (*Brassica oleracea* var *italica*) under Pune conditions, Asian. J Hort. 2011;6(1):74-6.
22. Husnan A, Mahmud F, Hossain MS, Begum S, SS, Hafiz TB. Multivariate analysis of Bottle gourd (*Lagenaria siceraria* L.). Int J Bio-Resource Stress Manag; 2013.
23. Mishra S, Pandey S, Kumar N, Pandey VP, Mishra N. Studies on gene action involved in inheritance of yield and yield attributing traits in kharif season bottle gourd [*Lagenaria siceraria* (Molina) standl.]. J Pharmacogn Phytochem. 2019;8(1):39-44.
24. Damor AS, Patil JN, Parmer HK, Vyas ND. Studies on genetic variability, heritability and genetic advance for yield and quality traits in bottle gourd [*Lagenaria siceraria* (Molina) Standl.] genotypes. Int J Environ Sci Technol. 2016;5(4):2301-7.
25. Kumar S, Thakur V, Tiwari R, Chormule SR. Evaluation of genotypes for quantitative traits in bottle gourd (*Lagenaria siceraria* (Mol.) standl.). J Pharmacogn Phytochem. 2018;7(3):841-3.
26. Ranjitha SR, Kumar Mahender, Jayasree G. Evaluation of rice (*Oryza sativa* L.) varieties and hybrids in relation to different nutrient management practices for yield, nutrient uptake and economics in SRI Annals of Biological Research. 2013;4(10):25-8.
27. Singh R, Kumar S, Kumar S. Performance and preference of broccoli varieties grown under low-hill conditions of Himachal Pradesh. Indian Research Journal. 2014;14(1):112-4.
28. Singh J. Response of cluster bean [*Cyamopsis tetragonoloba* (L.) Taub.] Varieties of different spacing during summer season. Unpublished M.Sc. (Agri.) Thesis. Gujarat Agricultural University, Sardar krushi Nagar; 2000.
29. Tamil Nadu Agricultural University. Origin, Area, Production, Varieties, Package of practices for cole crops cabbage. Available:[http://eagri/](http://eagri;); 2011. Tnau [cited May 2 2017]. Available:<http://ac.in/eagri50/HORTI281/lec12.html>.
30. Thakor DP. Influences of different cultivars and plant spacing on growth, yield and quality of garden pea. Unpublished M.Sc. (Agri.) Thesis, submitted to Sardarkrushinagar Dantiwada Agricultural University. Sardarkrushinagar; 2008.
31. Thakur S, Thakur R, Mehtha DK. Evaluation of different genotypes of broccoli in dry temperate conditions of Kinnur district of Himachal Pradesh in India. International Journal of science. Environ Technol. 2016;5(3):1673-9.
32. VDD. Vegetable development directorate. Khumaltar, Lalitpur; 2016.

33. Wakley A, Black IA. Critical examination of rapid method for determining organic carbon in soils, effect of variance in digestion conditions and inorganic soil constituents. *Soil Sci.* 1947;632:251.
34. Wilcox LV. Electrical Conductivity Am. Water Work Assoc J. 1950;42:775-6.
35. Zaki MF, Saleh SA, Tantawy AS, El-Dewiny CY. Effect of different rates of potassium fertilizer on the growth, productivity and ulaity of some broccoli cultivars under new reclaimed soil conditions. *Int J Chem Tech Res.* 2015;8(12):28-39.
36. Znidarcic, Marsic, Osvald, Pozrl, and Tradan. Yield and quality of cabbage inresponse to within row plant spacing. *Acta Agric.Slovenia.* 2007;89:15-23.

© 2022 Ashik et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/89983>