



## A Dual Method of Carroll's Generalized Canonical Correlation Analysis

Rodnellin Onesime Malouata <sup>a\*</sup>, Gelin Chedly Louzayadio <sup>b</sup>,  
Michel Koukouatikissa Diafouka <sup>c</sup> and Leonard Niere <sup>a</sup>

<sup>a</sup> *Departement of Licence, Higher Institute of Management, Congo.*

<sup>b</sup> *Departement of Licence, Faculty of Economics, Congo.*

<sup>c</sup> *Departement of Licence, Teachers' College, Congo.*

### Authors' contributions

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### Article Information

DOI: 10.9734/JAMCS/2021/v36i1130414

#### Editor(s):

(1) Dr. Dariusz Jacek Jakóbczak, Technical University of Koszalin, Poland.

#### Reviewers:

(1) Isra Natheer Alkallak, University of Mosul, Iraq.

(2) Hossam Kamal, Egypt.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here: <https://www.sdiarticle5.com/review-history/73297>

*Received: 16 July 2021*

*Accepted: 07 September 2021*

*Published: 23 December 2021*

**Original Research Article**

## Abstract

**Aims/Objectives:** We propose a dual method of Carroll's generalized canonical correlation analysis and we prove by means of the proposed criterion that the duality is formulated by exchanging the operators. It is an extension of Carroll's generalized canonical correlation analysis. The approach of analysis is illustrated on the basis of case study.

*Keywords:* Dual generalized canonical correlation analysis; data set partitioned in groups of individuals; principal component analysis.

**2010 Mathematics Subject Classification:** 53C25; 83C05; 57N16.

\*Corresponding author: E-mail: onesimero@gmail.com;

## 1 Introduction

Hotelling [1] proposed the first and oldest method called canonical correlation analysis. This method studies the relationships between two data matrices. Canonical correlation analysis focuses on the correlation between a linear combination of the variables in one set and a linear combination of the variables in another set. The idea is first to determine the pair of linear combinations having the largest correlation. Next, we determine the pair of linear combinations having the largest correlation among all pairs uncorrelated with the initially selected pair, and so on. Other developments of the canonical correlation analysis are in the research of Gittins [2] and Saporta [3].

A survey of the plethora of generalizations of canonical correlation analysis have been proposed for the analysis of the relationships with more than two sets of variables (e.g., Horst [4], Carroll [5], Kettenring [6]; see also Takane and Hwang [7], Wold [8-9], Gifi [10], Takane and Oshima-Takane [11], Dahl and Næs [12], Hanafi and Kiers [13]). However, we will restrict ourselves to the Carroll's [5] generalization.

Currently, in the context where the set of variables is observed on different groups of individuals, this problem has been studied in many scientific articles. If one expects the structure of each of the data blocks to be different, standard principal component analysis (PCA) (Jolliffe [14], Pearson [15]) can be performed on each data block. Moreover, one can carry out the simultaneous processing of all groups of variables. First, methods that are similar to PCA. Several methods combine the covariance matrices or the correlation matrices related to the various groups. The goal of these methods is to redistribute the explained variance by rotation. Krzanowski [16] proposed to carry out a multigroup PCA by diagonalizing the within group covariance matrix. This problem has been reworked and developed into a succession of PCA by Eslami, Qannari, Kohler and Bougeard [17]. In this context, we can cite the common principal component analysis (Flury [18]), simultaneous component analysis methods (Kiers and Ten Berge [19,20], Timmerman and Kiers [21], De Roover, Ceulemans and Timmerman [22]).

In practice, when we have a group of variables, the cloud of the individuals and the cloud of the variables are two representations of the same group: one across the rows and the other across the columns. Very strong relations called duality link these two clouds. Thus, there is an equivalence between multigroup data analysis methods and multiblock data analysis methods. In these conditions, several multigroup data analysis methods, such as the dual STATIS (Lavit [23]; Lavit, Escoufier, Sabatier and Traissac [24]), the dual multiple factor analysis (L, Husson and Pags [25]), can be seen as dual methods of multiblock data analyses.

Moreover, there are similar methods to the canonical correlation analysis among multigroup data analyses. We can cite: Between groups comparison of principal components (Krzanowski [26]) and Regularized generalized canonical correlation analysis for multiblock or multigroup data analysis (Takane, Hwang and Abdi [27], Tenenhaus and Tenenhaus [28]).

However, in the context where the set of variables is observed on different groups of individuals, the generalization of canonical correlation analysis (GCCA) according to Carroll [5] has never been discussed. In this paper, a version of GCCA called DGCCA (Dual generalized canonical correlation analysis), which can be applied to multiblock and multigroup, is described. We prove by means of the criterion that the duality is formulated by exchanging the covariance operator into scalar product. This criterion can be seen as a PCA in which the influence of the variables is stable.

This paper is organized as follows: in Section 2, we will describe the canonical correlation analysis proposed by Hotelling [1] and extended by Carroll [5]. In section 3, by using the criterion proposed by Carroll, we will propose the dual generalized canonical correlation analysis (section 3.1). Next, we also show that the dual generalized canonical correlation analysis can be reduced to the dual canonical correlation analysis (section 3.2) in the case of two matrices. Finally, in section 4, the experimental design is described and the obtained results are analyzed.

## 1.1 Sub-heading

Provide a factual background, clearly defined problem, proposed solution, a brief literature survey and the scope and justification of the work done, objective, etc.

# 2 Materials and Methods

## 2.1 Generalized canonical correlation analysis

We shall be interested in measures of association between two matrices. The first matrix  $X_1$ , of  $p$  columns, is summarized by the  $(n \times 1)$  canonical variate  $X_1 u_1$ . The second matrix  $X_2$ , of  $q$  columns, is summarized by the  $(n \times 1)$  canonical variate  $X_2 u_2$ . Specifically, we define canonical correlation analysis as the following optimization problem:

$$\text{Maximize } \text{corr}(X_1 u_1, X_2 u_2) \tag{2.1}$$

$$\text{subject to the constraints } \text{var}(X_1 u_1) = \text{var}(X_2 u_2) = 1$$

The first pair of canonical variates is the pair of linear combinations  $(X_1 u_1^1, X_2 u_2^1)$  having unit variances, which maximize the correlation (1); The second pair of canonical variates is the pair of linear combinations  $(X_1 u_1^2, X_2 u_2^2)$  having unit variances, which maximize the correlation (1) among all choices that are uncorrelated with the first pair of canonical variates. One may continue the search for canonical variates until all solutions are found.

To generalize the canonical correlation analysis of two sets of variables to  $K$  sets of variables, let  $X$  be a matrix  $X = [X_1, \dots, X_K]$  having  $n$  rows and  $p = \sum_{i=1}^K p_i$  columns, partitioned in  $K$  submatrices  $X_i$ . Each  $n \times p_i$  ( $n > p_i$ ) data matrix  $X_i$  represents a set of  $p_i$  centered and standardized variables observed on a set of  $n$  individuals. Let  $E_i$  ( $i = 1, \dots, K$ ) denote column spaces of  $X_i$ . Let  $P_i$  be the orthogonal projector onto the column space  $E_i$  of  $X_i$ ,  $P_i = X_i(X_i' X_i)^{-1} X_i'$ .

We only consider in this paper the generalization developed by Carroll [5] which consists of the following optimization problem:

$$\text{Maximize } \frac{1}{N} \sum_{i=1}^K \text{corr}^2(z, X_i u_i) \tag{2.2}$$

$$\text{subject to the constraint } z' z = 1$$

The principle of the generalized canonical correlation analysis is first to investigate the variables related to the set of the groups. These variables which summarize the general trends of the groups are called general variables. Furthermore, a general variable being obtained, we seek in each group a linear combination of variables related to this general variable. These linear combinations called canonical variates are the representations of the general variables in groups.

One of the advantages of this approach is to obtain the solution noniteratively (Kroonenberg [29]). Another advantage of this approach is that it is not necessary to define a link measurement

between two groups of variables but between a variable and a group (Escofier and Pags [30]). The measurement used by Carroll [5] is the square of the multiple correlation coefficient. Finally, this approach has the advantage of being probably the most simple and the richest of interpretations, because it easily connects to all the other methods of data analysis.

By definition, the multiple correlation coefficient between a variable and a group of variables  $X_i$  is the correlation coefficient between  $z$  and the linear combination of the variables of the group  $X_i$  most correlated with  $z$ . Geometrically, this linear combination is the orthogonal projection  $P_i z$  of  $z$  onto the column space  $E_i$  of  $X_i$ .

The relationship between the correlation coefficients and the variable  $z$  follows from the special structure of the matrix  $\sum_{i=1}^K P_i$ . In other words, the generalized canonical correlation analysis verifies the following stationary equation:

$$\frac{1}{N} \sum_{i=1}^K P_i z = \rho z \quad (2.3)$$

The solution of the generalized canonical correlation analysis shows that it can be obtained from the principal component analysis (Gifi [10], Saporta [3]). Thus, by setting  $\Sigma_i = X_i' X_i$ , the solution of GCCA verifies the following stationary equation:

$$XDX'z = \rho z \quad (2.4)$$

where  $D$  is a block diagonal matrix formed from  $\Sigma_i^{-1}$  as the  $i$ th diagonal block.

The GCCA is similar to a PCA in which each column space  $E_i$  of  $X_i$  is associated with the matrix  $\Sigma_i^{-1}$ .

In this section we will give some details of a solution to the dual generalized canonical correlation analysis.

## 2.2 Dual generalized canonical correlation analysis

We now consider a matrix  $X = [X_1' \cdots X_K']'$  having  $n = \sum_{i=1}^K n_i$  rows and  $p$  columns, partitioned in  $K$  submatrices  $X_i$  of order  $(n_i \times p)$  and row spaces  $E_i$  ( $i = 1, \dots, K$ ) of  $X_i$ . Each data matrix  $X_i$  represents a set of  $p$  centered and standardized variables observed on a set of  $n_i$  individuals. Let  $J_i$  be the orthogonal projector onto the row space  $E_i$  of  $X_i$ ,  $J_i = X_i'(X_i X_i')^{-1} X_i$ , where  $A^{-}$  is a  $g$ -inverse of  $A$ .

**Definition 2.1.** In the first step, the solution of the dual generalized canonical correlation analysis is a loading vector  $u$  and specific coefficients  $t_i = X_i' a_i$  ( $a_i$  being a vector to be determined of order  $(n_i \times 1)$ ) associated with data matrices  $X_i$  ( $i = 1, \dots, K$ ).

In our case, as in many similar situations, we define a mean-squared loss function. The core optimization problem considered in this paper is defined as follows:

$$\text{Minimize} \quad g(u) = \frac{1}{N} \sum_{i=1}^K \|u - J_i u\|^2 \quad (2.5)$$

subject to the constraints  $\|u\| = \|t_i\| = 1$ .

The loading vector is common to all matrices and its projection onto the subspace  $E_i$  is the specific coefficient  $t_i = J_i u$ . To convert the minimization problem into a maximization problem we rewrite (5) and manipulate the various terms. We obtain the following maximization problem:

$$\begin{aligned} \text{Maximize} \quad h(u) &= \frac{1}{N} \sum_{i=1}^K \langle u | J_i u \rangle = \frac{1}{N} \sum_{i=1}^K u' J_i u & (2.6) \\ \text{subject to the constraints} \quad & \|u\| = \|t_i\| = 1. \end{aligned}$$

From this problem, we can incorporate the constraint in the maximization problem by using Lagrange multiplier  $\lambda$  and obtain  $\tilde{h}$ ,

$$\tilde{h}(u, \lambda) = h(u) - \lambda(u'u - 1) \quad (2.7)$$

The maximum of  $h$  follows from the requirement that the first order partial derivatives of  $\tilde{h}$  are simultaneously zero at the maximum of  $h$  and that the Hessian is negative. We will state here the exact nature of the solution as proposition.

**Proposition 2.1.** *For  $J_i u = t_i$ , we obtain the following result:*

$$\frac{1}{N} \sum_{i=1}^K J_i u = \lambda u \quad (2.8)$$

*Proof.* Differentiating with respect to  $u$  and  $\lambda$ , and setting all the derivatives equal to zero, we obtain the following set of equations which have to be solved simultaneously for  $u$  and  $\lambda$ . Thus,

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^K J_i u &= \lambda u \\ u'u &= 1 \end{aligned}$$

□

It can be easily shown that for  $J_i u = t_i$ , the solution  $u$  verifies the stationary equation where  $u$  is a eigenvector of the matrix  $\frac{1}{N} \sum_{i=1}^K J_i$  related to the largest eigenvalue  $\lambda$ . By replacing  $J_i$  in  $J_i u = t_i$  and taking into account of  $t_i = X_i' a_i$ , we obtain  $a_i = (X_i X_i')^{-1} X_i u$ . The partial components are obtained by setting  $\xi_i = X_i u$ .

Moreover, as the projectors  $J_i$  are symmetric and idempotent, it follows that  $h(u) = \frac{1}{N} \sum_{i=1}^K \|J_i u\|^2$ . It seems that the dual generalized canonical correlation analysis (DGCCA) is similar to principal component analysis (PCA)(see proposition 3.2). The components are determined using the same loading vector  $u$  and the explained variance can be redistributed by rotation. It can be easily shown that the vector  $u$  which maximizes this average inertia is obtained by the eigenvector of  $\frac{1}{N} \sum_{i=1}^K J_i$  related to the eigenvalue  $\lambda = \frac{1}{N} \sum_{i=1}^K \|J_i u\|^2$ .

That is, after finding a first axis, this column is fixed and the second column is found in the residual space, by replacing  $X_i$  by  $X_i(I_p - uu')$  and proceeding in the same way as for the first step. Other loading vectors can be obtained by repeating the procedure. This procedure allows to construct the orthonormal basis. We can remark that the global components  $\xi^s = X u^s$  ( $s = 1 \dots, A$ , where  $A$  is the rank of the matrix  $X$ ) are orthogonal. The following proposition establishes that the dual generalized canonical correlation analysis is a PCA.

**Proposition 2.2.** For  $W_i = X_i X_i'$ , the dual generalized canonical correlation analysis is a PCA and verifies the following stationary equations:

$$X' D^- X u = \lambda u \quad (2.9)$$

$$X X' D^- \xi = \lambda \xi \quad (2.10)$$

where  $D^-$  is a block diagonal matrix formed from  $W_i^-$  as the  $i$ th diagonal block.

*Proof.* Indeed, as we have indicated, the same components may be determined by the PCA of the matrix  $X$ . One only has to consider  $X$  as a matrix having  $n = \sum_{i=1}^K n_i$  rows and  $p$  columns and  $J_i = X_i'(X_i X_i')^- X_i$ , we then set  $W_i = X_i X_i'$  as the matrix of scalar products between observations in the row space of  $X_i$  and  $\Lambda^-$  the block-diagonal matrix formed from  $W_i^-$ . We obtain  $\sum_{i=1}^K J_i = \sum_{i=1}^K X_i' W_i^- X_i = X' \Lambda^- X$ . The stationary equation (8) becomes (11)

$$\frac{1}{N} X' \Lambda^- X u = \lambda u. \quad (2.11)$$

By setting  $D^- = \frac{1}{N} \Lambda^-$  and by premultiplying (11) by  $X$ , it follows

$$X X' D^- \xi = \lambda \xi$$

$$X' D^- X u = \lambda u.$$

□

We then recognize the eigen equations yielding principal components and factor axes in PCA of  $X$ .

From the above, we conclude that the criterion of DGCCA is a PCA in which the duality is formulated by exchanging the covariance matrix  $\Sigma_i$  into scalar product matrix  $W_i$ . So, It is about substituting the block diagonal matrix  $D$  formed from  $\Sigma_i^{-1}$  by the block-diagonal matrix  $\Lambda^-$  formed from  $W_i^-$ .

Geometrically, each specific coefficient  $t_i$  associated with different matrices is the orthogonal projection  $J_i u$  of  $u$  onto the subspaces  $E_i$  spanned by the rows of  $X_i$ . Thus, the multiple correlation coefficient is the cosine of the angle  $\theta_i$  between the loading vector  $u$  and its projection  $J_i u$  onto  $E_i$ . Since  $u$  and  $J_i u$  are normalized, we then have:

$$\cos^2 \theta_i = \langle u | J_i u \rangle.$$

As  $u^{s'}(u^{s-1}, \dots, u^1) = 0$ , we obtain:

$$\sum_{i=1}^K \cos^2 \theta_i = \sum_{i=1}^K \langle u^s | J_i u^s \rangle = \langle u^s | \sum_{i=1}^K J_i u^s \rangle.$$

We achieve the same conclusion: the operator  $\sum_{i=1}^K J_i$  being a sum of the orthogonal projection operators, It is symmetric, diagonalizable and the eigenvectors are orthonormal.

### 2.2.1 Dual canonical correlation analysis

We shall be interested in the measures of association between two data matrices. Let  $E_1$  and  $E_2$  be the row spaces of  $X_1$  and  $X_2$ . We also introduce for each matrix, unit vectors  $t_1 = X_1' a_1$  and  $t_2 = X_2' a_2$  onto  $E_1$  and  $E_2$  respectively. The two vectors are as close as possible. Let  $J_1$  and  $J_2$  be orthogonal projections onto  $E_1$  and  $E_2$  defined by:  $J_1 = X_1'(X_1 X_1')^- X_1$  and  $J_2 = X_2'(X_2 X_2')^- X_2$  where  $A^-$  is a  $g$ -inverse of  $A$ . Let  $W_1 = X_1 X_1'$  and  $W_2 = X_2 X_2'$  be scalar product

matrices. Let  $W_{12}$  be the scalar product matrix of the individuals inter-data matrices and  $W_{21}$  its transpose.

The dual canonical correlation analysis (DCCA) problem consists of optimizing the following squared distance from  $t_1 = J_1 u$  and  $t_2 = J_2 u$ :

$$\text{Minimize} \quad k(a_1, a_2) = \|t_1 - t_2\|^2 \quad (2.12)$$

subject to the constraints  $\|t_1\| = \|t_2\| = 1$ .

This problem is equivalent to optimize the following criterion:

$$\text{Maximize} \quad k_1(a_1, a_2) = a_1 W_{12} a_2 \quad (2.13)$$

subject to the constraints  $a_1 W_1 a_1 = a_2 W_2 a_2 = 1$

We define dual canonical correlation (DCCA) as the optimization problem which consists of measuring the proximity of the individuals between the groups  $X_1$  and  $X_2$ . The Lagrangian function of optimization problem (8) is then considered:

$$\tilde{k}_1(a_1, a_2, \lambda_1, \lambda_2) = a_1 W_{12} a_2 - \frac{\lambda_1}{2}(a_1 W_1 a_1 - 1) - \frac{\lambda_2}{2}(a_2 W_2 a_2 - 1) \quad (2.14)$$

where  $\lambda_1$  and  $\lambda_2$  are the Lagrange multipliers associated with the constraints. Canceling the derivatives of the Lagrangian function with respect to  $a_1$ ,  $a_2$ ,  $\lambda_1$  and  $\lambda_2$ , we obtain using the stationary equations:

- $a_1$  is the eigenvector of the matrix  $W_1^- W_{12} W_2^- W_{21}$  associated with the largest eigenvalue  $\lambda_1^2$ ,
- $a_2$  is the eigenvector of the matrix  $W_2^- W_{21} W_1^- W_{12}$  associated with the largest eigenvalue  $\lambda_1^2$ .

The eigenvectors  $t_1$  and  $t_2$  are associated with the same eigenvalues. Thus, in DCCA, we also get:

- $t_1$  is the eigenvector of the matrix  $J_1 J_2$  associated with the largest eigenvalue  $\lambda_1^2$ ,
- $t_2$  is the eigenvector of the matrix  $J_2 J_1$  associated with the largest eigenvalue  $\lambda_1^2$ .

*Remarks 2.1* (See proof in the Appendix).

1. The dual generalized canonical correlation analysis generalizes the dual canonical correlation analysis.
2. The loading vector  $u$  is the principal factor according to normalized PCA of the vectors  $t_i$ .
3. The mean (sum) of the specific coefficients  $t_i$  and  $u$  are proportional.

### 3 Results and Discussion

We present the results of the dual generalized canonical correlation analysis and we repeat part of the multigroup PCA of Eslami, Qannari, Kohler and Bougeard [17]. The data set is the one used by Cortez, Cerdeira, Almeida and Reis [31] and by Eslami, Qannari, Kohler and Bougeard [17]. The study focuses on Portuguese wines. Specifically, we reanalyze matrices of eleven physicochemical variables observed on 4898 individuals for the white wines and 1599 individuals for the red wines. To eliminate the differences in variable means and variances, the data are centered and standardized per data matrix.

Regarding the graphical representation, the physicochemical variables are as follows:

Fixac=fixed acidity, Volac=volatile acidity, Citac=citric acid, Resug=residual sugar, Chlor=chlorides, Fredi=free sulfur dioxide, Totdi=total sulfur dioxide, Densi=density, pH, Sulph=sulphates, Alcoh=alcohol.

The goal of the dual generalized canonical correlation analysis is to help discover how the relationships vary in the experimental setup. In Table 1 below, we report the percentages of explained variances for first four principal components and global components obtained from the dual generalized canonical correlation analysis. Table 2 give the analogous results for the multigroup PCA.


**Table 1. The percentages of the explained variances of DGCCA**

Group	Dim1	Dim2	Dim3	Dim4
Red wines	28.61	13.74	11.69	9.30
White wines	21.96	21.51	10.44	10.41
Global component	44.02	19.17	11.94	8.29

**Table 2. The percentages of the explained variances of multigroup PCA**

Group	Dim1	Dim2	Dim3	Dim4
Red wines	29.19	14.08	11.21	9.28
White wines	23.36	21.58	13.46	11.12
Global component	27.25	15.91	11.61	9.75

Fig. 1 shows the representation of the variables on the first two loading vectors. The proportion of total variance accounted for the first principal component is 44.02%. The first principal component which explains 44.02% of the total variance, has an interesting subject-matter interpretation. The first principal component opposes the alcohol concentration to the sugar concentration and the density. Moreover, this component is correlated with sulphates and chlorides. This corresponds to the fermentation where some variables can be increased or decreased in the production process.



**Fig. 1. Map of the columns on the first two loading vectors of DGCCA**



The second axis explains 19.17% of the total variance. The first two principal components, collectively, explain 63.19% of the total variance, which is a very well result. The second axis represents 19.17% of the total variance and shows an opposition between the acid measurements and the pH.

For each of the two analyses, the eigenvalues of DGCCA are superior to the eigenvalues of multigroup PCA. The results of the DGCCA and the multigroup PCA are very little different.

## 4 Conclusions

In this paper, the duality between generalized canonical correlation analysis and its dual method is clearly defined. Generalized canonical correlation analysis uses the covariance operator, while dual generalized canonical correlation analysis uses the scalar product operator. The main aim of generalized canonical correlation analysis is to investigate the relationships between blocks, while the main aim of dual generalized canonical correlation analysis is to investigate the relationships among variables within the various groups. the PCA can be considered for both multiblock and multigroup data. The dual generalized canonical correlation analysis benefits the same advantages that the generalized canonical correlation analysis according to Carroll. We have shown in this paper that due to their higher flexibility and properties, it can be considered a straightforward extension of the PCA to multigroup PCA which facilitates the interpretation of results. The DGCCA criterion opens several ways of research in particular to find variables that discriminate the groups of variables. Moreover, if the number of individuals is much smaller than the number of variables, this method may exhibit instability due to ill-conditioned scalar product matrix. Under these conditions, a regularization of the method may be useful as in the case of generalized canonical correlation analysis (Tenenhaus et al. [32], Takane, Hwang and Abdi [27]).

## Competing Interests

Authors have declared that no competing interests exist.

## References

- [1] Hotelling H. Relations between two sets of variates. *Biometrika*. 1936;28:321-377.
- [2] Gittins R. *Canonical Analysis. A review with Applications in Ecology*, Springer-Verlag, Berlin; 1985.
- [3] Saporta G. *Probabilits, analyse des donnes et statistique*. Editions Technip, Paris; 2006.
- [4] Horst P. Relations among  $m$  sets of variables. *Psychometrika*. 1961;26:126-149.
- [5] Carroll JD. A generalization of canonical correlation analysis to three or more sets of variables. *Proceedings of the 76th Annual Convention of the American Psychological Association*. 1968;227-228.
- [6] Kettenring JR. Canonical analysis of several sets of variables. *Biometrika*. 1971;58:433-451.
- [7] Takane Y, Hwang H. Generalized constrained canonical correlation analysis. *Multivariate Behavioral Research*. 2002;37:163-195.
- [8] Wold H. Soft modeling: The basic design and some extensions. In K. G. Jreskog & H. Wold (Eds.), *Systems under Indirect Observation*. 1982;Part 2:1-54. Amsterdam: North-Holland.
- [9] Wold H. Partial least squares. In S. Kotz & N. L. Johnson (Eds.). *Encyclopedia of statistical sciences*. New York: John Wiley & Sons. 1985;6:581-591.
- [10] Gifi A. *Nonlinear multivariate analysis*. Chichester: Wiley; 1990.

- [11] Takane Y, Oshima-Takane Y. Nonlinear generalized canonical correlation analysis by neural network models. In S. Nishisato, Y. Baba, H. Bozdogan, & K. Kanefuji (Eds.), *Measurement and multivariate analysis*. Tokyo: Springer. 2002;183-190.
- [12] Dahl T, Ns T. A bridge between Tucker-1 and Carroll's generalized canonical analysis. *Computational Statistics and Data Analysis*. 2006;50:3086-3098.
- [13] Hanafi M, Kiers HAL. Analysis of K sets of data, with differential emphasis on agreement between and within sets. *Computational Statistics & Data Analysis*. 2006;51:1491-1508.
- [14] Jolliffe IT. *Principal component analysis* (2nd ed.). New York: Springer; 2002.
- [15] Pearson K. On lines and planes of closest fit to systems of points in space. *Philosophical Magazine*. 1901;2:559-572.
- [16] Krzanowski WJ. Principal component analysis in the presence of group structure. *Applied Statistics*. 1984;33:164-168.
- [17] Eslami E, Qannari EM, Kohler A, Bougeard S. Analyses factorielles de donnees structures en groupes d'individus. *Journal de la Societe Francaise de Statistique*. 2013;154:44-57.
- [18] Flury BN. Common principal components in  $K$  groups. *Journal of the American Statistical Association*. 1984;79:892-898.
- [19] Kiers HAL, Ten Berge JMF. Alternating least squares algorithms for simultaneous components analysis with equal component weight matrices in two or more populations. *Psychometrika*. 1989;54:467-473.
- [20] Kiers HAL, Ten Berge JMF. Hierarchical relations between methods for simultaneous component analysis and a technique for rotation to a simple simultaneous structure. *British Journal of Mathematical and Statistical Psychology*. 1994;47:109-126.
- [21] Timmerman ME, Kiers HAL. Four simultaneous component models for the analysis of multivariate time series from more than one subject to model intra-individual and inter-individual differences. *Psychometrika*. 2003;68:105-121.
- [22] De Roover K, Ceulemans E, Timmerman ME, Vansteelandt K, Stouten J, Onghena P. Clusterwise simultaneous component analysis for analyzing structural differences in multivariate multiblock data. *Psychological Methods*. 2012;17:100-119.
- [23] Lavit C. *Analyse conjointe de tableaux quantitatifs*. Masson; 1988.
- [24] Lavit C, Escouer Y, Sabatier R, Traissac P. The act (statis method). *Computational Statistics & Data Analysis*. 1994;18:97-117.
- [25] L S, Husson F, Pags J. Dmfa: Dual multiple factor analysis. *Communication in Statistics-Theory and Methods*. 2010;39(3):483-492.
- [26] Krzanowski WJ. Between-groups comparison of principal components. *Journal of the American Statistical Association*. 1979;33:164-168.
- [27] Takane Y, Hwang H, Abdi H. Regularized multiple-set canonical correlation analysis. *Psychometrika*. 2008;73(4):753-775.
- [28] Tenenhaus A, Tenenhaus M. (In press). Regularized generalized canonical correlation analysis for multiblock or multigroup data analysis. *European Journal of Operational Research*; 2014. Available:<http://dx.doi.org/10.1016/j.ejor.2014.01.008>.
- [29] Kroonenberg PM. *Applied multiway data analysis*. New York: Wiley; 2008.
- [30] Escofier B, Pags J. *Analyses factorielles simples et multiples. Objectifs, methodes et interpretation*. Dunod, Paris; 2008.
- [31] Cortez P, Cerdeira A, Almeida F, Matos T, Reis J. Modeling wine preferences by data mining from physicochemical properties. *Decision Support Systems*, Elsevier. 2009;47(4):547-553.
- [32] Tenenhaus A, Tenenhaus M. Regularized generalized canonical correlation analysis. *Psychometrika*. 2011;76:257-284.

## Appendix A. Proofs of the remarks

**Proof of 1.** The dual generalized canonical correlation analysis generalizes the dual canonical correlation analysis.

Indeed, the normalization constraints are the same as those of the dual canonical correlation analysis. To minimize  $\|u - J_1 u\|^2 + \|u - J_2 u\|^2$  subject to the normalization constraints over  $u$ ,  $t_1$  and  $t_2$  is equivalent to minimize  $\|J_1 u - J_2 u\|^2$ .

$$\begin{aligned} \|u - J_1 u\|^2 + \|u - J_2 u\|^2 &= \langle J_1 u | u \rangle + \langle u | J_2 u \rangle \\ &= \langle J_1 u | J_2 u \rangle \\ &= \|J_1 u - J_2 u\|^2 \end{aligned} \quad (4.1)$$

**Proof of 2.** The loading vector  $u$  is the principal factor according to normalized PCA of the vectors  $t_i$ .

In fact the specific coefficients being given,  $u$  is the loading vector that yields maximum  $\frac{1}{N} \sum_{i=1}^K \|u - J_i u\|^2$ , then by definition the principal factor of the specific coefficients  $t_i$ .

**Proof of 3.** The mean (sum) of the  $t_i$  and  $u$  are proportional.

Indeed, The stationary equation (8) shows that the only possible proportionality constant is  $\lambda$ . Since

$$\frac{1}{N} \sum_{i=1}^K t_i = \frac{1}{N} \sum_{i=1}^K J_i u = \lambda u \quad (4.2)$$

$\lambda$  being the largest eigenvalue of the mean (sum) of  $J_i$ .

© 2021 Malouata et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits un-restricted use, distribution and reproduction in any medium, provided the original work is properly cited.

**Peer-review history:**

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/73297>