



Influence of Organic Manures on Growth, Flowering, Quality and Yield of Carnation (*Dianthus caryophyllus*) Cv. Master under Naturally Ventilated Polyhouse Conditions of Prayagraj, India

Dikshit^{a++*} and Urfi Fatmi^{a#}

^a Department of Horticulture, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj- 211007 (U.P), India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i193518

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104885>

Original Research Article

Received: 07/06/2023

Accepted: 11/08/2023

Published: 14/08/2023

ABSTRACT

An experiment entitled Influence of organic manures on growth, flowering, quality and yield of carnation (*Dianthus caryophyllus*) cv. Master under naturally ventilated polyhouse conditions of Prayagraj was conducted in naturally ventilated polyhouse of Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj, during November, 2022 to April, 2023 with an aim to identify the most suitable organic manure combination under the agroclimatic conditions of

⁺⁺ M.Sc Research Scholar;

[#] Assistant Professor;

*Corresponding author: E-mail: dikshitarora5@gmail.com;

Prayagraj. There were thirteen treatments replicated thrice in Randomized Block Design (RBD). It was observed that treatment T₁₁ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m²) was significantly better in all the parameters studied like plant height (48.5 cm), number of primary branches (7.6), days taken to first bud initiation (93.5), flower diameter (6.3 cm), flower stalk length (77.2 cm), vase life (6.2 days), number of cut flowers per plant (5.6), number of cut flowers per 200 square meter (27,833.3).

Keywords: *Carnation; organic manures; FYM; vermicompost; neem cake; trichoderma.*

1. INTRODUCTION

Carnation (*Dianthus caryophyllus* L.) belongs to the Caryophyllaceae family and holds a significant position as one of the world's most popular cut flower crops, consistently ranking among the top ten cut flowers in terms of demand and commercial value. It is a half-hardy perennial cultivated in various regions across the globe, believed to be native to the Mediterranean. It is characterized by its diploid chromosome number of $2n = 30$. Carnations can be used in the garden for bedding, edging, borders, pots, and rock gardens in addition to producing cut flowers [1].

Excessive use of inorganic fertilizers for increasing the cut flower production has led to the leaching of nutrients causing soil hazards, altering the soil fertility and leading to pollution of soil and water bodies. They imposed disturbance in the ecosystem. Chemical fertilizers are commonly employed in carnation cultivation due to their ability to rapidly release essential elements to the crop [2,3]. However, their usage has been associated with certain negative effects on flower quality, as well as adverse impacts on soil health, water, and the environment. Organic manure is characterized as the product resulting from the controlled biological decomposition of organic matter. Organic matter has significant effect on the physico-chemical and biological characteristics of soil. Organic manures enrich the soil with organic matter, facilitating the gradual release of plant nutrients in a readily accessible form for crop utilization. Application of the organic manures and microbial agents make easy uptake of nutrients when required by the crop contrasting with chemical fertilizers [4].

Organic manures like FYM, Neem cake, Vermicompost etc. has great potential to boost yield. They play a vital role in enhancing flower yield, maintain soil health and also to sustain productivity [5,6]. Neem Cake is highly valued for its diverse composition of micro and macro nutrients. These nutrients play a crucial role in

supporting plant growth and development. Its organic nature makes it an environmentally friendly choice for sustainable agriculture practices. One of the significant benefits of neem cake is its ability to control soil-borne pathogens and nematodes.

Vermicompost is the end product of the decomposition process involving various species of worms, such as red wigglers, white worms, and earthworms. These worms work to break down a combination of decomposing vegetable or food waste, bedding material, and vermicast [6,7]. The process, known as vermicomposting, results in nutrient-rich organic matter suitable for enhancing soil fertility and plant growth. It has more nutrient contents and the availability of growth stimulating substances. In addition, there are many helpful microorganisms which assist in activities like nitrogen fixation, phosphorus solubilization and reduction of harmful microorganisms [8].

FYM is an important component of traditional farming systems and organic agriculture. It has been used for centuries as a natural fertilizer and soil conditioner. The primary objective of using FYM is to improve soil health, foster plant growth, and ensure the sustainable management of agricultural land.

Trichoderma has been widely studied and utilized as a biocontrol agent against plant pathogens, including fungi, in various agricultural systems. In case of carnation cultivation, *Trichoderma* species have shown promising results as organic fungicides. It can be applied as a preventative measure to protect carnations from fungal diseases, such as Fusarium wilt (caused by *Fusarium oxysporum*) and Botrytis blight (caused by *Botrytis cinerea*).

2. MATERIALS AND METHODS

2.1 Geographical Location and Climatic Conditions

Geographically, Prayagraj is situated in the South-Eastern part of Uttar Pradesh. It lies

between the parallels of 24°77' and 25°47' north latitudes and 81°19' and 82°21' east longitudes. The area of Prayagraj district comes under agro climatic zone V (Upper Gangetic Plain region) and sub-zone of Central Plains. The climate ranges from dry sub-humid to semi-arid and the soil is alluvium calcareous sandy loam. This District experiences average maximum temperature range between 43°– 47°C which may go as high as 48°C during peak summers (May-June). The minimum average temperature is 2-4°C, which may fall as low as 1°C during peak winter months (December-January). The average rainfall of the district is 1042 mm and the monsoon season is spread between July-September.

2.2 Experimental Details

The experiment was conducted in Randomized Block Design (RBD) with 13 treatments of organic manures with three replications in the Departmental Research Field of Department of Horticulture, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj during November, 2022 to April, 2023. Total number of treatments were thirteen viz. T₀-Control RDN (30:20:10 g/m² NPK), T₁ (RDN + FYM 5kg/m²), T₂ (RDN + Vermicompost 2.5 kg/m²), T₃ (RDN + Trichoderma 10g/m²), T₄ (RDN + Neem cake 1kg/m²), T₅ (RDN + FYM 2.5 kg/ m² + Vermicompost 1.25 kg/m²), T₆ (RDN + FYM 5 kg/m² + Trichoderma 10g/m²), T₇ (RDN + FYM 5kg/m² + Neem cake 1kg/m²), T₈ (RDN + Vermicompost 2.5kg/m² + Trichoderma 10g/m²), T₉ (RDN + Vermicompost 2.5kg/m² + Neem cake 1kg/m²), T₁₀ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Trichoderma 10g/m²), T₁₁ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m²), T₁₂ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Trichoderma 10g/m² + Neem cake 1kg/m²). Carnation cultivar Master was planted on 16th November, 2022 at a spacing of 20 cm x 20 cm.

3. RESULTS AND DISCUSSION

3.1 Plant Height and Number of Primary Branches

Maximum plant height (48.5 cm) and number of primary branches was observed from the plants grown in treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m² significantly superior to rest of the treatments. The treatment combination of RDN

(Recommended Dose of Nutrients) + FYM (Farm Yard Manure) + Vermicompost + Neem cake has shown significantly better plant height in carnations compared to other treatment combinations. This can be attributed to the synergistic effects of the various components. RDN provides essential nutrients in optimal proportions, promoting overall plant growth. FYM enhances soil structure and nutrient retention, leading to improved root development. Vermicompost enriches the soil with organic matter and beneficial microorganisms, facilitating nutrient uptake. Neem cake acts as a natural pesticide and also provides supplementary nutrients. Together, these components create a favorable environment for root expansion and nutrient absorption, resulting in taller, healthier carnation plants. The combination's balanced nutrient supply and disease-resistant properties contribute to the superior plant height observed. Similar findings were reported by Pooja et al. [9]; Bohra et al. [10]; Pandey et al. [11].

3.2 Days Taken to First Flower Bud Initiation

In terms of days taken to first bud initiation treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m² recorded minimum days taken to first bud initiation (93.5) whereas the maximum days were recorded in the treatment T₀ – Control (116.4). Organic manures likely promote early flowering by improving source-sink dynamics. This increases cytokinin synthesis in roots, aiding its swift transport to buds. This accelerates assimilate mobilization, facilitating the shift from vegetative to reproductive phase. Similar results were observed by Ranjan et al. [12]; Pandey et al. [11].

3.3 Flower Diameter (cm)

Data revealed that higher flower diameter was recorded in treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m² (5.3cm) whereas the minimum flower diameter was recorded in the treatment T₀- Control (4.2cm). This increase in flower diameter might be due to better nutrient uptake, higher photosynthesis and excellent physiological, biological activities due to presence of organic manures which have resulted in rapid synthesis and translocation of photosynthetic from the source to developing flower bud and finally increase in flower diameter Ranjan et al. [12]; Sindhu et al. [13].

Table 1. Effect of organic manures on plant height, number of primary branches, days taken to first flower bud initiation and flower diameter of carnation

Treatment Symbols	Treatment Combinations	Plant height (cm)	No. of primary branches	Days taken to first flower bud initiation	Flower diameter (cm)
		60 Days*	60 Days*		
T ₀	Control RDN (30:20:10 g/m ² NPK)	31.9	4.4	116.4	4.2
T ₁	RDN + FYM 5kg/m ²	45.1	6.1	114.0	5.1
T ₂	RDN + Vermicompost 2.5 kg/m ²	34.1	6.2	100.9	4.9
T ₃	RDN + Trichoderma 10g/m ²	40.5	5.9	102.9	4.8
T ₄	RDN + Neem cake 1kg/m ²	34.6	6.2	102.4	4.3
T ₅	RDN + FYM 2.5 kg/ m ² + Vermicompost 1.25 kg/m ²	39.5	4.5	111.4	5.3
T ₆	RDN + FYM 5 kg/m ² + Trichoderma 10g/m ²	36.4	6.4	101.5	4.9
T ₇	RDN + FYM 5kg/m ² + Neem cake 1kg/m ²	37.5	4.8	101.3	5.0
T ₈	RDN + Vermicompost 2.5kg/m ² + Trichoderma 10g/m ²	35.9	4.7	107.4	5.3
T ₉	RDN + Vermicompost 2.5kg/m ² + Neem cake 1kg/m ²	46.5	5.8	99	6.0
T ₁₀	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² +Trichoderma 10g/m ²	35.5	5.1	115.7	5.5
T ₁₁	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Neem cake 1kg/m ²	48.5	7.6	93.5	6.3
T ₁₂	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Trichoderma 10g/m ² + Neem cake 1kg/m ²	46.1	6.3	110.0	4.7
F-Test		S	S	S	S
SEd (±)		0.75	0.46	0.34	0.11
CV		2.33	9.80	0.55	2.64
CD_{0.05}		1.54	0.94	0.98	0.23

*Days after pinching

Table 2. Effects of organic manures on stalk length, vase life of flowers, no. of flowers per plant and number of flowers per 200m² of carnation

Treatment Symbols	Treatment Combinations	Stalk Length (cm)	Vase Life (days)	Number of flower stalks per plant	Number of flower stalks per 200m ²
T ₀	Control RDN (30:20:10 g/m ² NPK)	61.0	3.7	4.2	21000.0
T ₁	RDN + FYM 5kg/m ²	68.0	4.8	5.0	25166.7
T ₂	RDN + Vermicompost 2.5 kg/m ²	65.7	5.2	5.1	25333.3
T ₃	RDN + Trichoderma 10g/m ²	68.4	4.9	4.9	24500.0
T ₄	RDN + Neem cake 1kg/m ²	67.9	5.1	4.5	22666.7
T ₅	RDN + FYM 2.5 kg/ m ² + Vermicompost 1.25 kg/m ²	68.1	4.9	4.9	24666.7
T ₆	RDN + FYM 5 kg/m ² + Trichoderma 10g/m ²	64.9	4.8	4.8	23833.3
T ₇	RDN + FYM 5kg/m ² + Neem cake 1kg/m ²	66.8	4.8	4.8	23833.3
T ₈	RDN + Vermicompost 2.5kg/m ² + Trichoderma 10g/m ²	70.6	4.5	4.4	22166.7
T ₉	RDN + Vermicompost 2.5kg/m ² + Neem cake 1kg/m ²	73.9	5.3	5.3	26333.3
T ₁₀	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² +Trichoderma 10g/m ²	66.3	4.9	4.9	24500.0
T ₁₁	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Neem cake 1kg/m ²	77.2	6.2	5.6	27833.3
T ₁₂	RDN + FYM 2.5kg/m ² + Vermicompost 1.25kg/m ² + Trichoderma 10g/m ² + Neem cake 1kg/m ²	64.5	5.1	5.1	25500.0
F-Test		S	S	S	S
SEd (±)		0.69	0.24	0.13	660.76
CV		1.24	5.94	3.32	3.32
CD_{0.05}		1.42	0.49	0.27	1363.16

3.4 Stalk Length (cm)

Maximum flower stalk length was recorded in treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m² significantly superior to rest of the treatments. All treatments that included combinations with varying phosphorus levels exhibited significant effectiveness compared to the control. This can be attributed to the improved nutrient absorption in plants treated with organic manures, resulting in greater availability of assimilates required for enhancing flower stalk length. Pooja et al. [9]; Sindhu et al. [13] reported similar results.

3.5 Vase Life (Number of days)

In terms of vase life treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m² recorded maximum vase life (6.2 days) whereas the minimum vase life was recorded in the treatment T₀- Control (3.7 days). The use of organic fertilizer led to an increase in the vase life of carnations. This extension in vase life could be attributed to the stimulation of metabolic activity and the narrowing of the C:N ratio, which might have been facilitated by the substantial accumulation of carbohydrates. Furthermore, the use of the treatment combination RDN (Recommended Dose of Nutrients) + FYM (Farm Yard Manure) + Vermicompost + Neem cake has resulted in an improved vase life of carnation flowers compared to other treatment combinations. This improvement can be attributed to the combined effects of these components on flower quality and longevity. RDN supplies essential nutrients in optimal proportions, ensuring better flower health and resilience. FYM enhances soil structure and nutrient retention, allowing the flowers to access nutrients even after being cut. Vermicompost enriches the soil with organic matter and beneficial microorganisms, contributing to the flowers' overall vitality and vase life extension. Neem cake's presence helps in reducing post-harvest pests, minimizing petal damage and decay. The synergy of these components creates a favorable environment for cut flowers, resulting in longer-lasting and aesthetically pleasing carnations. These findings were supported by Pooja et al. [9]; Bohra et al. [10]; Sindhu et al. [13].

3.6 Number of Cut Flowers Per Plant/200m²

Yield is an important parameter to decide the efficacy of a treatment. Data recorded on

maximum number of flowers per plant (5.6) and maximum number of flowers per 200m² (27,833) in treatment T₁₁ containing RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m². The treatment combination of RDN (Recommended Dose of Nutrients) + FYM (Farm Yard Manure) + Vermicompost + Neem cake has led to a higher number of cut flowers per plant/200m² in carnations compared to other treatment combinations. This increase can be attributed to the combined effects of these components on plant growth and reproductive development. RDN supplies essential nutrients in optimal proportions, promoting robust flower bud initiation and development. FYM enhances soil fertility and water retention, ensuring continuous nutrient availability to support flower production. Vermicompost enriches the soil with organic matter and beneficial microorganisms, promoting healthy flowering and increased flower bud formation. Neem cake's presence provides natural pest resistance, preventing potential flower damage and loss. The synergy of these components creates an ideal growth environment, resulting in more abundant flower production in carnations, leading to higher yields of cut flowers per plant and per 200 square meter. Findings were in accordance with findings of Sindhu et al. [13]; Bohra et al. [10].

4. CONCLUSION

Based on the present investigation it is concluded that treatment T₁₁ (RDN + FYM 2.5kg/m² + Vermicompost 1.25kg/m² + Neem cake 1kg/m²) was found to be best in all the parameters like plant height, number of primary branches, days taken to first bud initiation, flower diameter, flower stalk length, vase life, number of cut flowers per plant, number of cut flowers per 200 square meter followed by T₉ (RDN + Vermicompost 2.5kg/m² + Neem cake 1kg/m²), whereas minimum is recorded in treatment T₀ (Control).

ACKNOWLEDGEMENT

I express my gratitude to my advisor Dr. Urfi Fatmi and all the faculty members of the Department of Horticulture for the constant supply to carry out the whole experimental research study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Dole JM, Wilkins HF. Floriculture principles and species, 2nd edition. Pearson Prentice Hall, Upper Saddle River, New Jersey. 2005;347.
2. Gupta P, Rajwal N, Dhaka VK, Rajwal D. Effect of different levels of vermicompost, NPK and FYM on performance of Gladiolus (*Gladiolus grandiflorus* L) cv. Happy End. The Asian Journal of Horticulture. 2008;3(1):142-143.
3. Kumar P, Sheoran S, Beniwal BS. Growth and yield parameters of rose as influenced by different organic manures and their levels. The Pharma Innovation Journal. 2022;SP-11(6):394-398.
4. Vanilarasu K, Balakrishnamurthy G. Influences of organic manures and amendments in soil physiochemical properties and their impact on growth, yield and nutrient uptake of banana. The Bioscan. 2014;9(2):525-529.
5. Gupta R, Yadav A, Garg VK. Influence of vermicompost application in potting media on growth and flowering of marigold crop. International Journal of Recycling of Organic Waste in Agriculture. 2014;3:47.
6. Swaroop K, Janakiram T. Response of bio fertilizers, FYM and their combined application with different levels of inorganic fertilizers in gladiolus (*Gladiolus hybridus* L.). International Journal of Tropical Agriculture. 2010;28(3-4):435-438.
7. Mogal SA, Khiratkar SD, Chopde NK, Dalvi AM, Kuchanwar OD, Khobragade YR. Effect of organic manures and biofertilizers with reduced doses of nitrogen on growth, yield and quality of China aster. Journal of Soils and Crops. 2006;16(1):180-185.
8. Barik T, Barik KC. Vermicomposting practices for its quality maintenance. Indian Farming. 2009;58(12):22-25.
9. Pooja G, Kumari S, Dikshit SN. Response of African marigold (*Tagetes erecta* L.) to integrated nutrient management. Annals of Biology. 2012;28(1):66-67.
10. Bohra M, Rana A, Punetha P, Upadhyay S, Nautiyal BP. Effect of organic manures and biofertilizers on growth and of Karnini China aster. Indian Journal of Horticulture. 2019;76(2):329-333.
11. Pandey SK, Kumari S, Singh D, Singh VK, Prasad VM. Effect of biofertilizers and organic manures on plant growth, flowering and tuber production of dahlia (*Dahlia variabilis* L.) Cv. SP. Kamala. Indian Journal of Pure & Applied Biosciences. 2017;5(2):549-555.
12. Ranjan S, Preetham SP, Satish C. Effect of organic manures and biofertilizers on vegetative, floral and post harvest attributes in tuberose (*Polianthes tuberosa*) var. Shringar. Asian Journal of Biological and Life Sciences. 2014;3(1): 6-9.
13. Sindhu SS, Gholap DB, Singh MC, Dhiman MR. Effect of medium amendments on growth and flowering in gerbera. Indian Journal of Horticulture. 2010;67(Special Issue):391-394.

© 2023 Dikshit and Fatmi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/104885>