



Plants Used in the Traditional Treatment of Female Infertility in Kita, Mali

Issiaka Togola ^{a*}, Mamadou A. Konaré ^a,
Haraba Dembele ^a, Savio Samake ^b and Nouhoum Diarra ^a

^a *Laboratoire de Biochimie Alimentaire et des Substances Naturelles (LBASNa), Faculté des Sciences et Techniques (FST), Université des Sciences, des Techniques et des Technologies de Bamako (USTTB), BP: 3206 Bamako, Mali.*

^b *Laboratoire de Botanique et d'Ecotoxicologie (LaBoTec), Faculté des Sciences et Techniques (FST), Université des Sciences, des Techniques et des Technologies de Bamako (USTTB), BP: 3206 Bamako, Mali.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2023/v38i1230619

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112400>

Original Research Article

Received: 25/10/2023
Accepted: 29/12/2023
Published: 30/12/2023

ABSTRACT

Aims: The aim of this study was to identify the plants used in the traditional treatment of female infertility in Kita, Mali.

Place and Duration of Study: The ethnobotanical investigation has been conducted in the town of Kita (Mali) from April to July, 2021. The data management and analysis were carried out at Faculty of Sciences and Techniques (FST), University of Sciences, Techniques and Technologies of Bamako (USTTB), Mali, between August 2021 and January 2023.

Methodology: Ethnobotanical survey was carried out using questionnaire addressed to the available people who gave their free consent.

*Corresponding author: E-mail: togola3@gmail.com, issika.togola@usttb.edu.ml;

Results: An ethnobotanical survey was carried out among fifty-two (52) people, including twenty (20) traditional practitioners, ten (10) herbalists, six (6) people who are both herbalists and traditional practitioners, and sixteen (16) housewives inside the town of Kita. The survey revealed 38 plant species belonged to 25 botanical families. The most represented families were Moraceae (28.94%), Sapotaceae (26.31%), Fabaceae (26.30%), Apocynaceae (21.05%). The most coveted plant species by the local populations to treat female infertility were *Ficus capensis* with 26.31%, *Vitellaria paradoxa* with 23.68%, *Stereospermum kunthinum* with 21.05%, *Leptadenia hastata* with 21.05%. The leaves have been the most frequently used organs (43%). These plants were mostly used as decoctions (80%) and administered by oral route (89%). The commonly associated ailments with female infertility included pelvic pains, dysmenorrhea and *Toxoplasma gondii* infections.

Conclusion: This survey has enabled us to gain a better understanding of the anti-fertility plants used in the town of Kita.

Keywords: Ethnobotanical survey; female infertility; Kita; Mali.

1. INTRODUCTION

Procreation has always been a vital objective for human beings and particularly for housewives. Today, women's ability to have children can be significantly impacted by environmental and social factors (Ménézo et al., 2012). These pressures, if poorly managed, could lead to infertility in women. This concept of infertility can be defined as a woman's inability to conceive a pregnancy after 12 months of regular unprotected sex. The infertility is estimated to affect between 8% and 12% of couples trough the world [1]. Very high rates of female infertility have been recorded in in sub-Saharan Africa [2] 11.7% in Kayes, Mali [3].

It is well known that the procreation always remains the fundamental reason for marriage in Africa As a result, the problems related to procreation negatively affect the life and well-being of individuals, couples and families until they become a heavy burden on their socio-economic development [4]. In Mali, marital infertility is a real social scourge, considered to be a specifically female concern in order to avoid to highlight the man's virility [5]. Accused of infidelity and prostitution, infertile women are ridiculed in their homes. Faced to this calvary, they become an easy prey for fetishists and charlatans. [6].

However, female infertility needs to be handled delicately, because it is often a divorce reason for couples in Africa, and generate psychological disorders in the Western countries [7]. Its treatment in modern medicine is costly and sometimes presents many side effects [2]. That's why the use of medicinal plants appears to be an appropriate alternative for resolving this infertility

problem in the worldwide and singularly in Africa [8]. Indeed, studies have reported that the African flora in general, and Malian flora in particular, is made up of a large reserve of medicinal plants, occupying an important place in African pharmacopoeia [9] Traoré et al., [10]. According to [11], traditional medicine offers effective and accessible options for the predominated pathologies in the local communities. It is also a national cultural heritage and a better way to connect people with their own history and culture. But, this common traditional practice for managing pathologies, like infertility, deserves to be framed by scientific research. That's why this study was initiated, with the aim of contributing to a better understanding and rational use of local plants used in the treatment of female infertility by the local populations of Kita in Mali.

2. MATERIALS AND METHODS

2.1 Study Sites

The ethnobotanical survey was carried out in the region of Kita, located at 187 km from Bamako, Mali (Fig. 1).

2.2 Data Collection

The survey took place from April 03 to July 28, 2021 in the town of Kita. It concerned anyone with some knowledge of medicinal plants who was willing to answer the semi-structured questionnaires. The random sampling and the snowball method were used to select the 52 respondents in the that study. After obtaining the free written consent from each participant, a questionnaire form (containing the opened and closed questions) was addressed to them.

Questions focused on the plant species used to treat female infertility, the organs/parts used, and methods of preparation and administration. In addition, the sociodemographic characteristics of the respondents were registered. Interviews were

conducted in the local language “Bamanankan”. The species cited were identified through search engines and resource persons. The genera and specific diversities were estimated based on the formula used by [9].




Fig. 1. Map of Mali with survey site location

2.3 Data Analysis

Excel version 2017® and SPSS Statistics 17 were used to analyze the data.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Socio-demographic characteristics of respondents

3.1.1.1 Distribution of respondents by gender

The Fig. 2 shows the repartition of respondents by gender. Of the 52 people surveyed, 38 were female (73%) and 14 male (27%).




Fig. 2. Distribution of respondents by gender

3.1.1.2 Distribution of respondents by age and profession

The Table 1 shows the distribution of 52 respondents by age and profession. The survey participants came from diverse backgrounds, encompassing ages from 20 to 80. Traditional practitioners and herbalists formed the majority, representing nearly 50% of those interviewed. Housewives (30.77%) also constituted a significant group, reflecting the community's involvement in traditional knowledge. While younger individuals participated, the most frequent respondents fell within the (50-65) age range, highlighting the value placed on experience in managing female infertility locally.

The Shapiro-wilk normality test reveals an asymmetrical distribution ($P= 0.000 < 0.05$) of respondents by gender, profession and age.

3.1.2 Distribution of species by botanical family and morphological types

The Fig. 3 shows the list of botanical families registered and their frequencies. Twenty-five (25) families were identified. The most represented families were Moraceae, Fabaceae, Sapotaceae, Apocynaceae and Bigoniaceae with more than 21.05% as citation frequencies. These species are belonged to four (4) morphological groups as indicated in the Table 3. The shrubs (42.10%) followed by the trees (31.59%) were the most important morphological types used by the respondents.

While 25 plant families were identified, Fabaceae and Rubiaceae showed the highest diversity in number of genera ($I_{dg} = 0.36$ and 0.12 , respectively). However, Moraceae had the highest diversity in number of species ($I_{ds} = 2$).

Table 1. Distribution of respondents by age and profession

Age groups (years)	Number & Percentages		Professions	Number & Percentages	
[20-35]	9 (17.31%)		Traditional healers	20 (38.46%)	
[35-50[17 (32.69%)		Herbalists	10 (19.23%)	
[50-65]	18 (34.62%)		Traditional herbalists	6 (11.54%)	
[65-80]	8 (15.38%)		Housewives	16 (30.77%)	
Total	52 (100%)		Total	52 (100%)	

Table 2. Normality test Shapiro-Wilk

Characteristics	Statistic	degree of liberty	P-value
Gender of respondent	0.554	52	0.000
Profession of respondent	0.786	52	0.000
Age of respondent	0.882	52	0.000

**The test is considered to be significant if P-value < 0.05*




Fig. 3. Distribution of species by botanical family

Table 3. Distribution of species according to morphological groups

Morphological types	Tree	Shrub	herbaceous	Liana	Total
Effectives	12	16	08	02	38
Frequencies (%)	31.59	42.10	21.05	5.26	100

Table 4. Index of genera and specific diversity

Botanical families	Number of genera	Number of species	Idg	Ids
Amaryllidaceae	1	1	0.04	1.00
Anacardiaceae	1	1	0.04	1.00
Annonaceae	1	1	0.04	1.00
Apocynaceae	1	1	0.04	1.00
Asteraceae	1	1	0.04	1.00
Bignoniaceae	1	1	0.04	1.00
Borraginaceae	1	1	0.04	1.00
Combretaceae	3	3	0.08	1.00
Euphorbiaceae	1	1	0.04	1.00
Fabaceae	7	7	0.36	1.00
Loganiaceae	1	1	0.04	1.00
Lythraceae	1	1	0.04	1.00
Meliaceae	1	1	0.04	1.00
Moraceae	1	2	0.04	2.00
Nymphaeaceae	1	1	0.04	1.00
Opiliaceae	1	1	0.04	1.00
Phyllanthaceae	1	1	0.04	1.00
Poaceae	2	2	0.08	1.00
Rhamnaceae	1	1	0.04	1.00
Rubiaceae	3	3	0.12	1.00
Rutaceae	1	1	0.04	1.00
Sapindaceae	1	1	0.04	1.00
Sapotaceae	2	2	0.08	1.00
Sterculiaceae	1	1	0.04	1.00
Vitaceae	1	1	0.04	1.00

*Idg: index of genera diversity; Ids: index of specific diversity

3.1.3 Plant organs used

The Fig. 4 shows the plant parts used and their frequencies. Several parts were cited, including leaves, bark, fruit, roots, stems, sheaths, the whole plant and the young plant. The leaves are the most frequently used, with 43% of mentions.

3.1.4 Preparation and administration modes

The Fig. 5 illustrates the different preparation methods and administration routes for the plant parts used.

Three main preparation methods were mentioned by respondents: decoction (80%), maceration (14%), spraying (5%) and other methods (1%). These different recipes were

administered by several routes, mainly by oral one (89%). Vaginal steaming was also noted.

3.1.5 Conditions favoring infertility

The survey results shed light on the multifaceted understanding of female infertility within the local context. Respondents attributed infertility to a spectrum of factors, with pelvic pain (54%) topping the list. Reproductive health concerns, including dysmenorrhea (17%) and abnormal menstrual cycles, were also prominent. Notably, respondents recognized the potential impact of infectious diseases like toxoplasmosis (17%) and other infections (4%). Additionally, a range of gynecological conditions, including uterine fibroids (3%) and candidiasis (3%), were identified as potential contributing factors.




Fig. 4. Frequencies of use of plant organs or parts




Fig. 5. Preparation and Administration modes of plant organs/parts

Table 5. List of plants: Families, organs used, preparation and administration methods and frequencies of use

N°	Families	Scientific names	Local names	Treated conditions	Parts used	Preparation mode	Mode of administration	Frequencies of citation (%)
1	Amaryllidaceae	<i>Allium ursinum</i> L.	Bakani-sabali	Fibroma	Whole plant	Decoction	Exposing the vagina to steam	2.63
2	Anacardiaceae	<i>Sclerocarya birrea</i> (A.Rich.) Hochst	Gouna	Pelvic pains	Barks	Decoction	Oral	2.63
3	Annonaceae	<i>Annona senegalensis</i> L.	Sounsoundiè	Pelvic pains	Leaves	Spraying	Oral	2.63
4	Apocynaceae	<i>Leptadenia hastata</i> (Pers.) Decne	Sarafaté / zoyé	Pelvic pains / dysmenorrhea / incessant menstruation	Leaves	Decoction	Oral	21.05
5	Asteraceae	<i>Eclipta prostrata</i> (L.) L.	Mousofing	Pelvic pains /Infection/candidose/ toxoplasmosis	Leaves	Decoction	Oral/bath	15.79
6	Bignoniaceae	<i>Stereospermum kunthinum</i> Cham.	Mogoyiri	Pelvic pains / toxoplasmosis	Leaves / Barks	Decoction / Maceration / Spraying	Oral	21.05
7	Borraginaceae	<i>Heliotropium indicum</i> L.	Nonsikou	Pelvic pains/ toxoplasmosis	Leaves/whole plant	Decoction	Oral	15.79
8	Combretaceae	<i>Combretum nigricans</i> Lepr. ex Guill. & Perr	Dianbagatawoulè	Pelvic pains	Roots	Decoction	Oral	2.63
9		<i>Guiera senegalensis</i> J.F.Gmel.	Koundiè	dysmenorrhea	Leaves	Decoction	Oral/Bath	2.63
10		<i>Pteleopsis suberosa</i> Engl et Diels	Tèrèni	Pelvic pains /candidosis	Barks	Decoction	Oral/Intimate bath	10.53
11	Euphorbiaceae	<i>Euphorbia hirta</i> L.	Denbasidji	Infection	Leaves	Decoction	Oral	2.63
12	Fabaceae	<i>Acacia nilotica</i> L.	Bouana	Infection/Candidosis	Fruits	Decoction	Oral / Intimate bath	7.89

N°	Families	Scientific names	Local names	Treated conditions	Parts used	Preparation mode	Mode of administration	Frequencies of citation (%)
13		<i>Cassia occidentalis</i> L.	Balan-balan	Pelvic pains	Roots	Maceration	Oral	2.63
14		<i>Dichrostachys glomerata</i> (Forssk.) Chiov	Triki	Pelvic pains	Leaves	Decoction	Oral	2.63
15		<i>Erythrina senegalensis</i> DC.	Ntébiléni	Pelvic pains	Leaves/Bark	Decoction	Oral	5.26
16		<i>Parkia biglobosa</i> (Jacq.) R.Br. ex G.Don	Néré	Pelvic pains	Barks	Maceration	Oral	2.63
17		<i>Pilostigma tonningii</i> (Schum.) Milne-Redh	Niama	Toxoplasmosis	Leaf	Decoction	Oral	2.63
18		<i>Tamarindus indica</i> L.	N'tomi	Pelvic pains	Leaf	Decoction	Oral	2.63
19	Loganiaceae	<i>Strychnos spinosa</i> Lam.	Kera / gangoro	Pelvic pains	Leaves	Decoction	Oral	2.63
20	Lythraceae	<i>Lawsonia inermis</i> L.	Diabi	Pelvic pains	Leaves	Decoction/spraying	Oral	7.89
21	Meliaceae	<i>Trichilia emetica</i> Vahl	soulafinsan	Pelvic pains	Roots/Bark	Spraying	Oral	2.63
22	Moraceae	<i>Ficus capensis</i> Thumb	Seretoro	Pelvic pains / toxoplasmosis	Leaves/Fruit/Bark	Decoction	Oral	26.31
23		<i>Ficus lecardii</i> Engl. & Diejs	Céré-diatikifaga	Pelvic pains	Leaves	Decoction	Oral	2.63
24	Nymphaeaceae	<i>Nymphaea lotus</i> L.	Djila gokou	dysmenorrhea / Pelvic pains	Leaves	Decoction	Oral	7.89
25	Opiliaceae	<i>Opilia celtidifolia</i> (Guill. & Perr.) Endl.	Koronkoi	Pelvic pains	Leaves	Decoction	Oral/ bath	2.63
26	Phyllanthaceae	<i>Hymenocardia</i>	Kalakari	Pelvic pains	Leaves	Decoction	Oral	2.63

N°	Families	Scientific names	Local names	Treated conditions	Parts used	Preparation mode	Mode of administration	Frequencies of citation (%)
		<i>acida</i> Tul.						
27	Poaceae	<i>Pennisetum clandestinum</i> Hochst. ex chov.	Sinkiri	Pelvic pains	Whole plant	Decoction	Oral	2.63
28		<i>Vetiveria nigriflora</i> (Benth.) Stapf	komourouni	Pelvic pains	Leaves	Decoction	Oral	2.63
29	Rhamnaceae	<i>Ziziphus mauritania</i> Lam.	N'tomono-missin	dysmenorrhea	Barks	Decoction	Oral	10.52
30	Rubiaceae	<i>Gardenia erubescens</i> Stapf et Hutch	M'bouré muso	Pelvic pains / dysmenorrhea	Mistletoe	Spraying	Oral	2.63
31		<i>Mitragyna inermis</i> (Willd.) Korth.	Djoun	dysmenorrhea	Fruits	Decoction	Oral	2.63
32		<i>Sarcocephalus latifolius</i> (Sm.) Bruce	Bati / Baro	Pelvic pains	Leaves	Decoction	Oral	2.63
33	Rutaceae	<i>Citrus lemon</i> (L.) Burm.f.	Lemouroucoumouni	Pelvic pains	Leaf	Decoction	Oral	2.63
34	Sapindaceae	<i>Paulinia pinnata</i> L.	Sokoni-nonfon	Toxoplasmosis	Leaves	Decoction	Oral	2.63
35	Sapotaceae	<i>Manilkara multinervis</i> (Baker) Dubard	Chi-sina	Pelvic pains	Leaves/Bark	Spraying	Oral	2.63
36		<i>Vitellaria paradoxa</i> C.F.Gaertn.	Chi	Toxo/ Pelvic pains	Bark / Mistletoe / Young plant	Decoction / Maceration / Spraying	Oral	23.68
37	Sterculiaceae	<i>Waltheria indica</i> L.	Dabada	Amenorrhea	Leaves	Decoction	Oral	2.63
38	Vitaceae	<i>Cissus populnea</i> Guill & Perr	Goumban / Ngaro	dysmenorrhea / Pelvic pains	Bark	Decoction	Oral	2.63




Fig. 6. Conditions leading to infertility

3.1.6 Plant species surveyed

Analysis of Table 2 shows that a total of 38 different species belonging to 25 families used in the treatment of female infertility at Kita were recorded. *Ficus capensis* (26.31%), *Vitellaria paradoxa* (23.68%), *Leptadenia hastata* (21.05%) and *Stereospermum kunthinum* (21.05%) were the most coveted species by the local populations of Kita. Respondents have also mentioned incessant menstruations (1%) or heavy menstrual bleeding or metrorrhagia.

3.2 Discussion

3.2.1 Socio-demographic data

The analysis of sociodemographic characteristics of 52 people surveyed show that only 14 were male gender (27%) against 38 females (73%) (Fig. 2). The target population was made up of three occupational categories, of which 38% were traditional practitioners, 19% herbalists, 12% traditional practitioner-herbalists and 31% housewives. Respondents ranged in age from 20 to 80, with the majority between 35-65 age group (Table 1). Otherwise, the Khi^2 test shows that the traditional treatment are strongly dominant by the female among the respondents ($P = 0.000 < 0.05$). This activity is also dependent of the profession and age ($P = \text{value} < 0.05$). These results are contrary to those of several authors [8, 12] who have reported that the majority of respondents are male gender.

3.2.2 Medicinal plants used

Examination of the survey forms revealed 38 plant species belonged to 25 botanical families

(Fig. 3). The most frequently cited species was *Ficus capensis* with 26.31% followed by *Vitellaria paradoxa* (23.68%). The survey conducted by Diawara [13] in two Bamako markets showed that the various *Ficus* species are frequently used in the treatment of dysmenorrhea in Mali, which is one of the best-known factors that can cause female infertility. Among these 38 plant species used to treat the female infertility, 28.94% of them were Moraceae, 26.30% were Fabaceae and Sapotaceae and 21.05% were Apocynaceae and Bigoniaceae. Furthermore, these species are belonged to four (4) morphological. The shrubs (42.10%) followed by the trees (31.59%) were the most important morphological forms used by the local populations of Kita. The same families and morphological groups have been reported by [2] in Dabakala's department in Ivory coast. In opposite, 48% of these found plant species were herbaceous against only 16% for shrubs. These differences could be linked to the geographical and climatic conditions of investigated areas.

The most frequently used organ was the leaf, with 43% of citations, followed by bark (26%), stems, and mistletoe (6%) each (Fig. 4). Ethnobotanical surveys conducted by Kemadji et al. [14] and Koman et al. [2] on plants used in female infertility have also reported high rates of leaf utilization; 35% and 46% respectively. This high use of leaves could be due to the availability and the ease of collection [15]. Furthermore, the literature have attributed this high use of leaves to their richness in bioactive compounds which are endowed with many pharmacological properties [16, 10].

Thanks to their pharmacological properties, these bioactive compounds are responsible for their use in the treatment of female infertility [14]. The use of leaves must be encouraged as it represents no danger to plant regeneration and ensures the conservation of floristic richness [17].

These organs were prepared mainly as decoctions by the majority of local populations (80%) and generally administered orally (89%). These results corroborate those of many authors [4,2,17]. According to Savio et al. [17] this could be justified by the fact that decoction allows the most active principle to be collected and cancels out the toxic effect of certain recipes. The preference of oral route could be explained by the fact that the disease is linked to bacterial and fungal infections most often located in the deep organs [2]; so to reach them, the biocompounds must transit through the digestive tract [18,19].

As the treatment of female infertility is traditionally complex, local populations provided better management of factors that contribute to it. Among these factors favoring female infertility, they are pelvic pains (54%), dysmenorrhea (17%) and toxoplasmosis (17%).

These collected data, could be affected by people's perceptions and methods of respondents' recruitment (snowball, random sampling). To address these limitations, future investigations must be undertaken to gain deeper insights into local understandings and practices related to female infertility and to establish causal relationships between identified factors and infertility outcomes.

4. CONCLUSION

This study explored local knowledge and practices in managing female infertility in Kita. Participants revealed a diverse arsenal of plant species, favoring leaves and barks prepared as decoctions and administered orally. Notably, pelvic pain, dysmenorrhea, and toxoplasmosis were recognized as potential contributors. These findings hold significant value. Further research should delve into the scientific potential of these plants (phytochemical composition, toxicity and anti-fertility activities), while recognizing the importance of understanding their cultural significance, traditional knowledge systems, and ecological sustainability. This holistic approach could benefit numerous stakeholders – from traditional healers who can enhance their

practices to healthcare professionals who can better communicate with patients seeking traditional remedies. Ultimately, such collaboration bridges the gap between traditional knowledge and scientific exploration, paving the way for improved health outcomes in Kita and beyond.

CONSENT

As per international standards or university standards, Participants' written consent has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mélodie VB, Christine W. fertility and infertility: Definition and epidemiology, Clinical Biochemistry. Clinical Biochemistry. 2018;62:2–10. Available: <https://doi.org/10.1016/j.clinbiochem.2018.03.012>
2. Koman Silvère R, Blaise KW, Konan YAO, Ouattara D. Palms used in the traditional treatment of female infertility in the Dabakala department (Ivory Coast). 2019;42(1):7086–7099.
3. Diassana M, Dembele S, Macalou B, Sidibe A, Hamidou A, Doumbia F, Haidara M, Kane F, Sylla C, Bocoum A, Traore S. Impact of the couple's infertility on the psychological state of women at the Fousseyni Daou hospital in Kayes. 2023;7(2):323–329.
4. Houmenou V, Adjatin A, Tossou MG, Yedomonhan H, Dansi A, Gbenou J, Akoegninou A. Ethnobotanical study of plants used in female sterility in the departments of Ouémé and the plateau in southern Benin. International Journal of Biological and Chemical Sciences. 2017;11(4):1851–1871.
5. Traore FB. Study of marital infertility in the Gyneco-obstetrics departments of the POINTG hospital. 2010;123.
6. Tembely F. Sterility of the couple in the reference health center of commune V-Bamako in relation to 145 cases. 2008;101.
7. Kouamé N, N'goan-Domoua AM, Konan N, Sétchéou A, OlivierTra-Bi, N'gbesso R,

- Kéita AK, Kéita A. Contribution of transvaginal ultrasound associated with hysterosalpingography in the etiological research of female infertility in Abidjan (Ivory Coast). 2012;16(4):43–49.
8. Françoise AA, Koffi K, William D, Emmanuel B, Atèhèzi T, Kosi NM, Kudzo GA, Jacques D, Amégnona A, Sodokè TK, Messanvi G. Ethnobotanical investigation into the traditional management of female infertility in the Savanes health region of Togo. *European Scientific Journal, ESJ*. 2018;14(3):358. Available: <https://doi.org/10.19044/esj.2018.v14n3p358>
 9. Togola I, Konaré MA, Diarra N, Karembé M. Ethnobotanical and floristic investigations of *Datura* species in Mali. *European Journal of Medicinal Plants*. 2022;33(12):73–87. Available: <https://doi.org/10.9734/EJMP/2022/v33i121114>
 10. Togola I, N Diaye M, Traoré N, Konaré AM, Diarra N. Ethnobotanical and comparative study of the antioxidant and anti-inflammatory potential of three organs of *Zanthoxylum zanthoxyloides*, a plant used in the traditional treatment of sickle-cell disease in Bamako. *GSC Biological and Pharmaceutical Sciences*. 2023;25(01):019–030.
 11. Sanogo B. Study of the antioxidant and anticonvulsant activities of two (02) medicinal plants from Mali. 2010;116.
 12. Houmènou V, Adjatin A, Assogba F, Gbénou J, Akoègninou A. Phytochemical and cytotoxicity study of some plants used in the treatment of female sterility in South Benin. *European Scientific Journal*. 2018;14:6. Available:<https://doi.org/10.19044/esj.2018.v14n6p156>
 13. Diawara B. Study of the phytochemistry and biological activities of two ficus used in the traditional treatment of dysmenorrhea. Faculty of Medicine, Pharmacy and Odontostomatology of the University of Bamako. 2010;88.
 14. Kemadji JN, Bakarnga-VIA I, Nguinambaye MM, Otchom BB. Ethnobotanical and phytochemical study of medicinal plants used in the treatment of female infertility in N'Djamena, Chad. 2022;21(2):131–140.
 15. Tuttolomondo T, Licata M, Leto C, Savo, Bonsangue G, Letizia Gargano M, Venturella G, La Bella S. Ethnobotanical Investigation on Wild Medicinal Plants in the Monti Sicani Regional Park (Sicily, Italy). *Journal of Ethnopharmacology*. 2014;153(3):568–586.
 16. Telefo P, Lemfack M, Bayala B, Lienou LL, Goka M, Mouokeu C, Tagne S, Moundipa F. Ethnopharmacological investigation of plants used in the treatment of female infertility in the localities of Fossong-Wentcheng and Foto, Cameroon. Researchgate; 2012. Available: <https://doi.org/10.1007/s10298-011-0678-6>
 17. Savio S, Issiaka T, Nouhoum D, Samake M, Rokia S. Ethnobotanical study and phytochemical screening of ornamental plants for therapeutic use in the district of Bamako, Mali. *International Journal of Advanced Research*. 2020;8(8):251–262. Available: <https://doi.org/10.21474/ijar01/11487>
 18. Tra Bi F, Irie G, N'Gaman K, Mahou C. Studies of some therapeutic plants used in the treatment of high blood pressure and diabetes: two emerging diseases in Ivory Coast. *Science & Nature*. 2008;5(1):39–48. <https://doi.org/10.4314/scinat.v5i1.42150>
 19. Traoré K, Haidara M, Denou A, Dembélé SM, Sanogo R. Medicinal plants used in the treatment of childhood illnesses in four ecological and cultural zones of Mali. *RAMReS Review – Pharmacopoeia and Traditional African Medicine Series*. 2023;22(1):91–102.

© 2023 Togola et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112400>