



# Mean Performance of Various Quantitative Characters in Bottle Gourd [*Lagenaria siceraria* (Molina) Standl.] Genotypes Under Temperate Conditions of Kashmir

Khansa Bashir <sup>a</sup>, Ummyiah H. Masoodi <sup>a\*</sup>,  
Baseerat Afroza <sup>a</sup>, Gazala Nazir <sup>a</sup>, Gowhar Ali <sup>b</sup>,  
Nageena Nazir <sup>c</sup>, Syed Berjes Zehra <sup>a</sup> and Oneza Aftab <sup>a</sup>

<sup>a</sup> Division of Vegetable Science, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar, Srinagar-190 025, Jammu and Kashmir, India.

<sup>b</sup> Division of Genetics and Plant Breeding, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar, Srinagar-190 025, Jammu and Kashmir, India.

<sup>c</sup> Division of Agricultural Statistics, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar, Srinagar-190 025, Jammu and Kashmir, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JABB/2024/v27i1689

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112442>

Original Research Article

Received: 26/11/2023

Accepted: 31/01/2024

Published: 09/02/2024

## ABSTRACT

Seventy-seven phenotypically diverse genotypes, including two checks (Pusa Naveen and Pusa Santushti) of bottle gourd, were evaluated for various quantitative traits at the Experimental field of Urban Technological Park (Habbak), Srinagar, Jammu and Kashmir during Kharif-2022. A single-

\*Corresponding author: E-mail: masoodi.ummyiah@gmail.com;

factor experiment was laid out in an Augmented Randomized Block Design with three blocks. Seventy-seven genotypes were randomly distributed among the three blocks, with each block receiving thirty-one lines, including two checks (Pusa Naveen and Pusa Santushti), replicated thrice. Analysis of variance revealed significant differences among genotypes for all the traits. The maximum fruit yield per hectare (ha<sup>-1</sup>) was recorded in IC-371747-1 (1132.21 q), followed by IC-382258 (848.199 q) and IC-398541 (845.33 q). On the other hand, the lowest fruit yield per hectare (ha<sup>-1</sup>) was recorded in IC-522878 (211.83 q), followed by Pusa Naveen (220.30 q) and SKUA-BG-7 (241.78 q). The average fruit yield per hectare (ha<sup>-1</sup>) for all the genotypes was 430.78 q.

**Keywords:** Bottle gourd; fruit yield; genotypes; mean performance.

## 1. INTRODUCTION

Bottle gourd [*Lagenaria siceraria* (Molina) Standl.] (2n=2x=22) belongs to family Cucurbitaceae and is one of the most ancient crops cultivated during summer throughout the world. The genus *Lagenaria* is derived from the latin word *lagena*, meaning the bottle. It is also known as Calabash, Doodhi and Lauki in different parts of India [1]. Its primary centre of origin is Africa [2]. The fossil records indicate its culture in India even before 200 B.C. It has been found wild in India, the Moluccas and Ethiopia. It has spread to western countries from India and Africa. The genus *Lagenaria* includes six species that are distributed in Africa, Madagascar, Indo-Malaysia and the neotropics. There is only one cultivated species, *Lagenaria siceraria*, which is annual and monoecious. The five other species are wild, perennial and dioecious, occurring in East Africa and Madagascar. The tender fruits are also used to prepare sweets, raita and pickles. The dried shells of mature fruits are extremely hard and are used as containers, utensils, musical instruments, floats of fishnets or ornamental items. The leaves are also used to prepare vegetable and they have higher nutritive value than fruits, in respect of protein, fat, minerals, fibre, carbohydrate, energy, calcium and phosphorus content. Different plant parts of bottle gourd have several putative medicinal properties. It is suggested for consumption during convalescence because it is easily digestible. The bottle gourd is a very beneficial vegetable in avoiding digestive issues like constipation due to the dietary fiber it contains. Consuming fiber has been linked positively to a decline in the incidence of diabetes and coronary heart disease [3]. Along with being a potent thirst quencher, bottle gourd juice has been used traditionally as a medicine to cure ulcers, indigestion and acid reflux. The natural remedy is thought to be effective against certain poisons and scorpion stings. It also has laxative and

cooling properties. The fruit is used in the treatment of asthma and other bronchial illnesses because it is thought to have the potential to reduce pain and is efficient against fever. It is a good source of organic antioxidants as well. Due to its high dietary fiber and low fat and cholesterol content, it aids in weight loss quickly [4]. Jaundice can be treated with a decoction produced from leaves. To decrease the effects of heat during the hotter months, people massage the cut surface of tiny fruit on their hands and feet. Bottle gourd is recommended for the treatment of diabetes, hypertension, flatulence, liver illnesses, excessive weight problems and other conditions in Ayurveda. The edible portion of bottle gourd contains about 95.54% moisture and is rich in vitamins, minerals, antioxidants and dietary fibres. Its fruit contains Vitamin C (10.1 mg), Vitamin A (0.0048 mg), Thiamine (29 mg), Riboflavin (22 mg), Niacin (320 mg), Carbohydrates (3390 mg), Fats (2 mg), Potassium (150 mg), Calcium (26 mg) and Iron (0.20 mg)/100g [5].

In India the total area under its cultivation is 187 thousand hectares with an annual production of 3165 thousand MT [6]. Nonetheless, in Jammu and Kashmir it is grown over an area of 1.60 thousand hectares with a production 36.17 thousand MT [7]. Major bottle gourd growing states in India are Bihar, Uttar Pradesh, Madhya Pradesh, Haryana, Chhattisgarh and West Bengal [8].

## 2. MATERIALS AND METHODS

The present research study was carried out at the experimental field of Urban Technological Park, Habbak, Srinagar, Jammu and Kashmir during *khariif*-2022. It is situated at an altitude of 1608 meters above mean sea level lying between 34.16° North latitude and 74.83° East longitude. The

climate is temperate characterized by mild summers. The mean minimum and maximum temperatures are 2.42°C and 30°C recorded in the months of October and August-September (respectively). The maximum rain fall is received during June (14.94 mm). The material used for research work consisted seventy-seven genotypes of bottle gourd which were procured from different sources. The experiment was laid out in accordance with Augmented Randomized Block Design comprising of seventy-seven treatments and three blocks. Recommended agronomic practices were followed to raise a good crop. Observations recorded were node number at which 1<sup>st</sup> male flower appeared, node number at which 1<sup>st</sup> female flower appeared, days to appearance of 1<sup>st</sup> male flower, days to appearance of 1<sup>st</sup> female flower, days to anthesis of 1<sup>st</sup> male flower, days to anthesis of 1<sup>st</sup> female flower, number of male flowers plant<sup>-1</sup>, number of female flowers plant<sup>-1</sup>, days to 1<sup>st</sup> fruit harvest, days to last fruit harvest, vine length (m), number of primary branches, number of fruits plant<sup>-1</sup>, fruit weight (kg), fruit length (cm), fruit diameter (cm), fruit yield plant<sup>-1</sup> (kg), fruit yield ha<sup>-1</sup> (q).

### 3. RESULTS AND DISCUSSION

The analysis of variance revealed that all the characters under study exhibited highly significant differences among the genotypes thus, suggesting existence of sufficient variability in the germplasm studied [Table 1 (a-b)]. The estimates of mean values revealed that no genotype was superior for all the characters under consideration (Tables 2, 3 and 4). However, genotypes like SKAU-BG-2 (6.14) followed by IC-276153 (6.21) and SKAU-BG-5 (6.25) were found superior for node number at which 1<sup>st</sup> male flower appeared; IC-536894-1 (7.24) followed by IC-331981 (7.25) and IC-310206-1 (7.27) were superior for node number at which 1<sup>st</sup> female flower appeared; Pusa Naveen (41.62 days) followed by IC-294891-1 (41.68 days) and SKUA-BG-4, IC-418491-A (41.91 days) for days to appearance of 1<sup>st</sup> male flower; IC-385816-1 (46.08 days) followed by IC-276153 (46.40 days) and Pusa Naveen (46.42 days)

for days to appearance of 1<sup>st</sup> female flower; Pusa Naveen (45.88 days) followed by IC-385816-1 (46.04 days) and SKUA-BG-3 (46.09 days) for days to anthesis of 1<sup>st</sup> male flower; IC-276153 (48.99 days) followed by SKUA-BG-4 (49.93 days) and Pusa Naveen (49.95 days) for days to anthesis of 1<sup>st</sup> female flower. Similar results have also been reported by Harika et al. [9], Jain et al. [10], Rambabu et al. [11] and Rashid et al. [12].

Genotypes like IC-316017-1 (159.20) followed by IC-307077 (155.90) and IC-276552 (150.10) were found superior for number of male flowers plant<sup>-1</sup>; IC-310206 (18.64) followed by IC-306422 (18.24) and IC-382258 (16.82) for number of female flowers plant<sup>-1</sup>; IC-294891-1 (61.06 days) followed by SKUA-BG-4 (61.57 days) and Pusa Naveen (61.67 days) for days to 1<sup>st</sup> fruit harvest; IC-424502 (147.78 days) followed by IC-385816 (147.57 days) and IC-321410 (147.33 days) for days to last fruit harvest; SKUA-BG-8 (8.98 m) followed by IC-321121 (8.53 m) and SKUA-BG-3 (8.19 m) for vine length; IC-371747-1 (21.33) followed by IC-398541 (21.31) and IC-536594 (21.22) for number of primary branches. This is in conformity to the findings of Jain et al. [10], Rambabu et al. [11] and Kandasamy et al. [13], Khan et al. [14] and Rashid et al. [12].

IC-310206 (8.6) followed by SKUA-BG-4 (8.2) and SKUA-BG-3 (7.4) were found superior for number of fruits plant<sup>-1</sup>; IC-382258 (4.40 kg) followed by IC-371747-1 (4.36 kg) and IC-398541 (4.17 kg) for fruit weight; IC-382258 (63.54 cm) followed by SKUA-BG-1 (61.46 cm) and IC-316017-1 (56.16 cm) for fruit length; IC-421962 (16.43 cm) followed by IC-319460 (16.42 cm) and Pusa Santushti (16.00 cm) for fruit diameter; IC-371747-1 (22.64 kg) followed by IC-382258 (16.96 kg) and IC-398541 (16.93 kg) for fruit yield plant<sup>-1</sup>; IC-371747-1 (1132.21 q) followed by IC-382258 (848.199 q) and IC-398541 (845.33 q) for fruit yield ha<sup>-1</sup>. Similar findings were observed by Jain et al. [10], Rambabu et al. [11], Kandasamy et al. [13], Rashid et al. [12], Sohi et al. [15] and Dubey et al. [16].

**Table 1(a). Analysis of variance with respect to MSS for various characters in bottle gourd [*Lagenaria siceraria* (Molina) Standl.]**

Source of variation	d.f	Mean sum of squares (MSS)											
		Node No. at which 1 <sup>st</sup> male flower appears	Node No. at which 1 <sup>st</sup> female flower appears	Days to appearance of 1 <sup>st</sup> male flower	Days to appearance of 1 <sup>st</sup> female flower	Days to anthesis of 1 <sup>st</sup> male flower	Days to anthesis of 1 <sup>st</sup> female flower	No. of male flowers plant <sup>-1</sup>	No. of female flowers plant <sup>-1</sup>	Days to 1 <sup>st</sup> fruit harvest	Days to last fruit harvest	Vine length (m)	No. of primary branches
Blocks	2	0.008	0.171	32333	0.361	0.122	0.665	7.41	0.210	0.513	0.081	0.382	3.028*
Treatments	76	1.162**	6.870**	31.130**	34.213**	35.031**	31.143**	704.17**	4.010**	81.576**	35.837**	4.651**	23.822**
Error	15 2	0.148	0.139	0.595	0.439	0.369	0.305	6.29	0.211	0.263	0.304	0.132	0.974

\*, \*\*= Significant at 5% and 1% respectively

**Table 1(b). Analysis of variance with respect to MSS for various characters in bottle gourd [*Lagenaria siceraria* (Molina) Standl.]**

Source of variation	d.f	Mean sum of squares (MSS)					
		No. of fruits plant <sup>-1</sup>	Fruit weight (kg)	Fruit length (cm)	Fruit diameter (cm)	Fruit yield plant <sup>-1</sup> (kg)	Fruit yield ha <sup>-1</sup> (q)
Blocks	2	0.570	0.055	3.035	0.284	1.180	2183
Treatments	76	3.765**	1.604**	203.635**	16.298**	34.637**	86146**
Error	152	0.18	0.031	1.780	0.341	0.613	1504

\*, \*\*= Significant at 5% and 1% respectively

**Table 2. Mean performance of Bottle gourd [*Lagenaria siceraria* (Molina) Standl.] genotypes for various quantitative characters**

S. No.	Genotypes	Node No. at which 1 <sup>st</sup> male flower appears	Node No. at which 1 <sup>st</sup> female flower appears	Days to appearance of 1 <sup>st</sup> male flower	Days to appearance of 1 <sup>st</sup> female flower	Days to anthesis of 1 <sup>st</sup> male flower	Days to anthesis of 1 <sup>st</sup> female flower
1.	Pusa Naveen	6.80	9.76	41.62	46.42	45.88	49.95
2.	Pusa Santushti	6.61	8.33	42.15	47.03	47.08	50.14
3.	SKUA-BG-1	7.64	9.16	44.87	49.75	49.33	52.85
4.	SKUA-BG-2	6.14	7.92	48.84	54.27	54.26	57.42
5.	SKUA-BG-3	8.13	10.28	42.23	47.04	46.09	49.99
6.	SKUA-BG-4	6.99	8.34	41.91	47.34	46.73	49.93
7.	SKUA-BG-5	6.25	8.70	43.25	47.89	48.08	51.04
8.	SKUA-BG-6	6.66	9.79	45.39	49.33	50.25	52.82
9.	SKUA-BG-7	6.90	11.55	42.82	47.42	47.05	50.94
10.	SKUA-BG-8	8.33	11.92	50.08	55.20	55.82	58.39
11.	IC-047045	7.55	9.26	48.89	54.05	53.67	57.17
12.	IC-262868	6.84	10.45	44.17	49.09	49.42	52.29

S. No.	Genotypes	Node No. at which 1 <sup>st</sup> male flower appears	Node No. at which 1 <sup>st</sup> female flower appears	Days to appearance of 1 <sup>st</sup> male flower	Days to appearance of 1 <sup>st</sup> female flower	Days to anthesis of 1 <sup>st</sup> male flower	Days to anthesis of 1 <sup>st</sup> female flower
13.	IC-256053	6.33	10.92	51.14	56.17	56.04	58.04
14.	IC-256052	7.33	8.88	51.06	56.05	55.82	58.56
15.	IC-256051	8.26	12.05	50.00	55.08	54.96	58.07
16.	IC-256043	6.42	9.13	48.73	53.77	53.81	56.64
17.	IC-284953	7.25	10.28	44.34	49.42	49.98	52.46
18.	IC-284895	7.01	7.56	50.94	55.87	55.79	58.90
19.	IC-284874-1	6.54	8.02	48.95	54.09	53.82	56.39
20.	IC-284874	8.12	12.10	47.91	52.99	52.28	55.93
21.	IC-284816	7.31	7.69	50.62	55.94	56.16	58.83
22.	IC-276552	7.72	11.37	48.85	53.78	54.05	56.64
23.	IC-276153	6.21	8.99	41.92	46.40	46.33	48.99
24.	IC-294891	6.85	9.40	45.68	50.79	50.93	53.52
25.	IC-294891-1	7.64	7.78	41.68	46.76	46.46	50.15
26.	IC-319460	7.07	11.88	43.75	48.78	48.88	51.72
27.	IC-316017-1	8.16	8.44	49.66	54.58	55.17	58.04
28.	IC-318883-1	8.31	11.04	42.33	47.77	46.83	50.45
29.	IC-310206-1	6.33	7.27	44.64	49.37	49.24	52.52
30.	IC-310206	6.33	9.23	51.48	56.03	56.30	59.10
31.	IC-307077	7.19	8.15	42.45	46.74	47.15	49.98
32.	IC-310188	7.83	11.99	45.20	51.03	50.37	54.11
33.	IC-306422	7.34	11.11	42.99	47.94	47.58	51.07
34.	IC-297846	7.04	8.77	45.37	50.54	50.56	53.64
35.	IC-306128-A	6.62	10.25	50.03	54.94	55.23	57.99
36.	IC-418491	6.84	9.25	43.81	48.42	48.86	51.73
37.	IC-385816	6.71	7.85	48.82	53.42	53.66	57.02
38.	IC-382258	7.54	8.47	47.14	52.34	52.33	55.89
39.	IC-371747-1	8.32	8.79	42.03	47.52	46.11	50.73
40.	IC-371747	8.04	11.05	50.11	55.31	55.04	58.86
41.	IC-392392	7.72	7.75	48.90	54.48	54.44	58.86
42.	IC-522868-1	7.45	9.65	44.73	48.63	49.31	52.62
43.	IC-371697	7.29	8.77	44.24	49.18	47.80	52.93
44.	IC-522866	7.32	10.53	50.21	55.09	54.99	58.65
45.	IC-522856	6.92	11.14	45.27	51.01	50.72	54.97
46.	IC-426990	6.95	8.74	48.10	53.48	53.03	58.20
47.	IC-424502	6.63	8.17	50.02	55.50	54.84	58.59
48.	IC-411915	6.78	7.84	49.58	54.62	54.48	57.77
49.	IC-522868	7.15	8.21	48.46	53.38	52.74	57.47
50.	IC-522876	7.22	10.32	46.74	51.72	50.95	54.60
51.	IC-522878	6.74	7.44	48.38	53.35	53.44	56.96

S. No.	Genotypes	Node No. at which 1 <sup>st</sup> male flower appears	Node No. at which 1 <sup>st</sup> female flower appears	Days to appearance of 1 <sup>st</sup> male flower	Days to appearance of 1 <sup>st</sup> female flower	Days to anthesis of 1 <sup>st</sup> male flower	Days to anthesis of 1 <sup>st</sup> female flower
52.	IC-385816-1	6.88	11.24	42.44	46.08	46.04	50.56
53.	IC-394736	7.98	11.95	43.62	47.57	47.44	51.64
54.	IC-394857	8.16	9.85	45.78	51.08	50.98	54.70
55.	IC-398541	8.32	7.66	44.67	49.52	49.65	53.53
56.	IC-417705	7.75	8.55	49.98	54.86	54.99	58.37
57.	IC-421962	7.66	10.35	42.49	47.66	47.56	51.69
58.	IC-418491-A	6.82	7.95	41.91	46.68	46.96	50.05
59.	IC-342078	6.62	11.99	43.77	48.34	48.66	54.30
60.	IC-321121	8.15	11.11	44.60	49.69	49.68	53.76
61.	IC-341390	7.62	9.08	50.06	55.98	55.99	58.50
62.	IC-339209	7.08	8.54	47.19	51.97	52.20	56.00
63.	IC-536894-1	7.27	7.24	42.38	47.50	47.30	51.57
64.	IC-331981	6.79	7.25	41.95	46.60	46.92	51.06
65.	IC-330999	8.26	11.61	42.70	47.34	47.82	50.12
66.	IC-321460	7.44	10.12	46.38	51.52	51.27	54.07
67.	IC-325973	7.53	11.98	43.50	48.68	47.98	51.30
68.	IC-321559	7.97	9.69	42.99	48.09	47.06	51.40
69.	IC-321412	6.76	7.54	44.51	49.21	49.61	52.63
70.	IC-321414-1	8.55	8.32	50.49	55.26	55.12	58.48
71.	IC-312410-1	6.73	8.74	49.96	55.06	54.98	58.08
72.	IC-321410	8.51	10.35	50.10	55.03	55.23	58.22
73.	IC-330987	7.56	7.75	51.28	55.06	56.01	58.46
74.	IC-331121	7.88	11.86	50.45	55.26	55.52	58.33
75.	IC-342080	6.77	9.15	47.69	52.84	52.64	55.76
76.	IC-546151	7.14	12.54	50.02	55.37	55.21	58.14
77.	IC-536594	6.84	9.15	48.43	53.76	52.63	56.24
<b>Mean</b>		<b>7.24</b>	<b>9.52</b>	<b>46.38</b>	<b>51.37</b>	<b>51.24</b>	<b>54.64</b>
<b>C.V</b>		<b>5.32</b>	<b>3.91</b>	<b>1.66</b>	<b>1.28</b>	<b>1.18</b>	<b>1.01</b>
<b>C.D. 5%</b>		<b>0.82</b>	<b>0.60</b>	<b>1.24</b>	<b>1.06</b>	<b>0.97</b>	<b>0.89</b>

**Table 3. Mean performance of bottle gourd [*Lagenaria siceraria* (Molina) Standl.] genotypes for various quantitative characters**

S.No.	Genotypes	No. of male flowers plant <sup>-1</sup>	No. of female flowers plant <sup>-1</sup>	Days to 1 <sup>st</sup> fruit harvest	Days to last fruit harvest	Vine Length (m)	No. of Primary Branches
1	Pusa Naveen	98.73	14.09	61.67	135.89	4.45	10.50
2	Pusa Santushti	99.17	14.12	65.03	136.90	7.92	17.33
3	SKUA-BG-1	148.69	15.95	62.47	140.36	7.49	18.24
4	SKUA-BG-2	134.55	16.02	75.60	145.57	5.42	15.74
5	SKUA-BG-3	149.75	16.63	62.64	139.55	8.19	19.88
6	SKUA-BG-4	143.21	17.88	61.57	134.77	8.09	18.37
7	SKUA-BG-5	97.79	13.94	70.78	140.68	4.84	12.38
8	SKUA-BG-6	122.28	15.26	62.75	137.11	5.13	14.35
9	SKUA-BG-7	137.87	15.33	62.57	137.83	4.01	10.40
10	SKUA-BG-8	148.03	14.92	68.56	139.99	8.98	19.10
11	IC-047045	141.17	15.66	74.74	142.66	7.79	20.16
12	IC-262868	100.92	14.37	71.62	144.12	4.96	14.79
13	IC-256053	135.74	15.08	77.54	146.31	6.16	13.70
14	IC-256052	122.70	15.32	73.66	143.04	4.41	13.41
15	IC-256051	100.45	14.33	75.50	145.33	3.92	14.69
16	IC-256043	124.24	13.79	75.27	147.00	3.46	12.09
17	IC-284953	149.60	14.96	62.32	139.53	5.39	14.87
18	IC-284895	139.55	13.94	75.66	142.89	4.90	15.65
19	IC-284874-1	131.04	14.60	74.21	144.94	5.32	16.42
20	IC-284874	147.60	15.12	74.78	140.98	5.59	16.61
21	IC-284816	139.00	14.27	74.52	143.74	5.46	14.63
22	IC-276552	150.10	15.98	72.06	145.44	4.93	15.54
23	IC-276153	114.72	14.89	62.23	137.89	4.44	12.72
24	IC-294891	134.46	15.00	72.31	143.77	5.23	13.73
25	IC-294891-1	134.70	15.13	61.06	138.44	4.95	14.91
26	IC-319460	143.01	15.71	75.26	143.74	6.13	18.59
27	IC-316017-1	159.20	15.95	72.27	141.77	6.36	17.72
28	IC-318883-1	136.20	16.44	62.13	135.83	6.73	20.33
29	IC-310206-1	124.29	13.96	75.75	144.80	6.66	19.99
30	IC-310206	150.08	18.64	77.45	146.82	6.75	20.35
31	IC-307077	155.90	15.65	65.64	138.94	5.49	17.00
32	IC-310188	130.05	15.38	75.18	145.20	6.71	18.73
33	IC-306422	140.88	18.24	74.13	143.90	6.17	17.11
34	IC-297846	146.50	15.76	75.62	145.47	5.02	13.58
35	IC-306128-A	134.10	14.85	75.78	146.14	6.68	18.62
36	IC-418491	121.36	15.72	67.32	140.09	5.42	17.70

S.No.	Genotypes	No. of male flowers plant <sup>-1</sup>	No. of female flowers plant <sup>-1</sup>	Days to 1 <sup>st</sup> fruit harvest	Days to last fruit harvest	Vine Length (m)	No. of Primary Branches
37	IC-385816	101.99	15.78	77.10	147.57	5.45	18.64
38	IC-382258	120.96	16.82	77.19	147.16	6.74	18.98
39	IC-371747-1	131.58	14.89	77.04	146.95	6.76	21.33
40	IC-371747	136.70	14.04	77.16	146.61	5.27	20.36
41	IC-392392	113.89	16.66	72.24	140.99	5.47	19.75
42	IC-522868-1	120.75	17.51	62.42	136.58	5.41	18.44
43	IC-371697	127.60	16.08	63.16	135.71	6.18	20.59
44	IC-522866	119.28	15.12	72.86	142.73	5.33	18.63
45	IC-522856	140.76	16.56	76.20	145.63	5.26	17.88
46	IC-426990	134.90	13.67	76.43	146.19	3.71	11.16
47	IC-424502	120.69	13.68	77.18	147.78	4.80	14.48
48	IC-411915	132.57	14.73	71.85	142.18	4.94	15.71
49	IC-522868	140.04	15.56	76.64	144.67	5.25	17.42
50	IC-522876	144.00	14.40	77.92	146.74	4.72	15.16
51	IC-522878	100.59	14.37	75.97	145.35	3.42	11.80
52	IC-385816-1	139.50	13.95	76.07	144.79	5.08	15.32
53	IC-394736	142.70	14.27	76.25	145.99	5.23	16.59
54	IC-394857	124.47	13.83	76.35	146.06	4.87	14.75
55	IC-398541	114.88	14.36	73.91	141.82	6.75	21.31
56	IC-417705	125.73	13.97	76.66	145.53	5.09	14.07
57	IC-421962	111.76	13.97	74.93	143.47	5.28	16.80
58	IC-418491-A	149.90	14.99	75.56	146.18	5.20	17.43
59	IC-342078	105.04	13.13	76.34	147.10	5.44	17.16
60	IC-321121	125.64	13.96	75.24	145.95	8.53	19.44
61	IC-341390	110.80	13.85	77.81	146.59	7.85	21.12
62	IC-339209	127.44	14.16	74.78	144.62	6.76	17.49
63	IC-536894-1	143.00	14.30	73.85	145.31	5.53	16.59
64	IC-331981	136.00	13.60	74.87	146.56	5.76	16.39
65	IC-330999	136.56	15.14	75.19	146.06	7.70	17.22
66	IC-321460	118.86	14.77	75.78	145.80	5.15	18.04
67	IC-325973	133.91	14.86	75.11	145.95	6.00	20.25
68	IC-321559	125.89	13.99	76.18	144.73	6.59	19.95
69	IC-321412	109.45	13.64	75.22	143.64	6.41	20.05
70	IC-321414-1	136.33	13.64	76.33	145.56	5.74	18.91
71	IC-312410-1	122.29	15.45	76.19	146.91	7.22	17.79
72	IC-321410	116.83	14.62	76.96	147.33	7.23	16.94
73	IC-330987	145.64	16.17	76.78	147.21	6.79	20.84
74	IC-331121	124.51	13.76	77.82	147.00	8.10	21.17



S.No.	Genotypes	No. of male flowers plant <sup>-1</sup>	No. of female flowers plant <sup>-1</sup>	Days to 1 <sup>st</sup> fruit harvest	Days to last fruit harvest	Vine Length (m)	No. of Primary Branches
75	IC-342080	106.01	13.18	71.37	142.92	8.05	20.36
76	IC-546151	135.99	13.58	76.97	146.78	6.68	20.85
77	IC-536594	148.13	14.81	75.73	145.59	7.62	21.22
<b>Mean</b>		<b>129.78</b>	<b>14.97</b>	<b>72.82</b>	<b>143.58</b>	<b>5.90</b>	<b>17.12</b>
<b>C.V</b>		<b>1.93</b>	<b>3.07</b>	<b>0.70</b>	<b>0.38</b>	<b>6.16</b>	<b>5.76</b>
<b>C.D. 5%</b>		<b>4.04</b>	<b>0.74</b>	<b>0.82</b>	<b>0.88</b>	<b>0.58</b>	<b>1.59</b>

**Table 4. Mean performance of Bottle gourd [*Lagenaria siceraria* (Molina) Standl.] genotypes for various quantitative characters**

S. No.	Genotypes	No. of Fruits plant <sup>-1</sup>	Fruit Weight (kg)	Fruit Length (cm)	Fruit Diameter (cm)	Yield plant <sup>-1</sup> (kg)	Yield ha <sup>-1</sup> (q)
1.	Pusa Naveen	4.9	0.89	31.83	6.35	4.363	220.307
2.	Pusa Santushti	3.5	3.48	42.81	16.00	12.368	618.638
3.	SKUA-BG-1	5.4	2.24	61.46	5.00	12.262	612.829
4.	SKUA-BG-2	5.2	1.38	32.80	7.23	7.189	361.071
5.	SKUA-BG-3	7.4	1.65	44.77	5.72	12.482	630.362
6.	SKUA-BG-4	8.2	1.48	37.13	5.64	12.315	616.282
7.	SKUA-BG-5	4.3	1.23	43.50	6.15	5.286	265.729
8.	SKUA-BG-6	5.1	1.17	38.19	5.04	5.970	301.489
9.	SKUA-BG-7	6.0	0.79	41.16	6.14	4.763	241.783
10.	SKUA-BG-8	5.3	2.57	44.43	7.13	13.889	694.981
11.	IC-047045	5.9	1.88	48.23	5.22	11.461	573.328
12.	IC-262868	3.5	1.72	48.46	6.15	6.027	303.511
13.	IC-256053	4.1	1.99	26.22	8.01	8.160	409.141
14.	IC-256052	5.2	1.43	53.13	6.18	7.429	372.610
15.	IC-256051	3.4	1.59	38.70	6.00	5.461	275.802
16.	IC-256043	3.5	1.41	43.39	5.50	4.989	251.782
17.	IC-284953	4.3	1.50	54.76	5.40	6.446	323.767
18.	IC-284895	3.5	1.62	42.71	5.31	5.667	284.353
19.	IC-284874-1	3.7	1.69	53.05	5.88	6.248	313.895
20.	IC-284874	4.5	2.01	51.76	5.49	9.043	453.235
21.	IC-284816	3.6	1.92	49.81	5.37	6.972	352.276
22.	IC-276552	4.4	1.50	51.74	5.72	6.603	331.097
23.	IC-276153	3.7	1.59	53.84	5.90	5.884	295.424
24.	IC-294891	3.6	2.25	45.67	7.99	8.083	404.612
25.	IC-294891-1	3.6	1.70	36.52	8.70	6.112	306.699
26.	IC-319460	5.3	1.87	12.67	16.42	10.161	508.480

S. No.	Genotypes	No. of Fruits plant <sup>-1</sup>	Fruit Weight (kg)	Fruit Length (cm)	Fruit Diameter (cm)	Yield plant <sup>-1</sup> (kg)	Yield ha <sup>-1</sup> (q)
27.	IC-316017-1	5.4	2.16	56.16	5.61	12.162	607.856
28.	IC-318883-1	6.8	2.05	51.14	7.52	13.861	693.116
29.	IC-310206-1	3.7	3.70	53.49	8.29	13.786	690.066
30.	IC-310206	8.6	1.50	42.04	5.81	13.000	651.134
31.	IC-307077	5.2	1.51	36.70	6.23	7.903	397.826
32.	IC-310188	3.8	4.26	43.90	8.28	16.428	822.145
33.	IC-306422	6.7	1.61	38.00	12.64	10.849	544.290
34.	IC-297846	4.4	1.41	39.00	6.89	6.213	313.190
35.	IC-306128-A	5.3	2.52	34.93	8.13	13.224	661.778
36.	IC-418491	4.1	1.69	33.80	6.63	6.988	351.761
37.	IC-385816	4.9	1.76	40.90	6.07	8.623	432.624
38.	IC-382258	3.8	4.40	63.54	8.97	16.964	848.199
39.	IC-371747-1	5.2	4.36	47.00	9.51	22.648	1132.212
40.	IC-371747	3.4	2.13	46.20	9.14	7.241	362.883
41.	IC-392392	6.9	1.25	40.90	6.86	8.626	432.728
42.	IC-522868-1	6.2	1.26	51.50	6.04	7.815	392.025
43.	IC-371697	5.5	2.10	37.90	8.81	11.627	582.166
44.	IC-522866	5.4	1.63	35.70	8.36	8.857	445.228
45.	IC-522856	4.6	1.44	34.10	7.80	6.627	332.308
46.	IC-426990	3.6	1.52	36.90	7.83	5.471	274.403
47.	IC-424502	3.5	1.78	43.14	6.19	6.291	317.501
48.	IC-411915	3.7	2.03	32.20	10.64	7.513	376.660
49.	IC-522868	4.4	1.49	44.10	5.70	6.557	328.624
50.	IC-522876	3.5	1.52	41.30	5.58	5.319	267.444
51.	IC-522878	3.6	1.17	48.90	5.25	4.211	211.839
52.	IC-385816-1	4.1	1.94	47.20	6.27	8.021	404.137
53.	IC-394736	3.7	1.75	42.70	6.86	6.477	324.993
54.	IC-394857	3.5	1.49	45.95	6.92	5.216	261.952
55.	IC-398541	4.1	4.17	51.00	7.95	16.938	845.332
56.	IC-417705	3.8	1.90	51.00	7.52	7.221	362.053
57.	IC-421962	3.9	1.99	22.50	16.43	7.760	375.605
58.	IC-418491-A	4.1	1.79	52.40	6.45	7.338	368.234
59.	IC-342078	3.7	1.56	48.30	5.11	5.837	294.502
60.	IC-321121	3.8	3.16	44.20	9.07	11.913	598.087
61.	IC-341390	3.7	2.43	46.10	5.90	9.023	456.043
62.	IC-339209	3.9	1.99	47.20	6.57	7.839	394.227
63.	IC-536894-1	4.0	1.56	51.70	5.60	6.251	314.105
64.	IC-331981	3.4	1.50	34.43	5.62	5.150	259.388
65.	IC-330999	4.3	1.64	39.90	5.32	7.109	357.914

S. No.	Genotypes	No. of Fruits plant <sup>-1</sup>	Fruit Weight (kg)	Fruit Length (cm)	Fruit Diameter (cm)	Yield plant <sup>-1</sup> (kg)	Yield ha <sup>-1</sup> (q)
66.	IC-321460	4.2	1.58	38.40	5.11	6.700	338.102
67.	IC-325973	4.5	1.79	41.50	6.11	8.116	408.479
68.	IC-321559	4.2	1.86	45.00	6.32	7.811	391.790
69.	IC-321412	3.7	1.82	51.93	6.31	6.793	342.531
70.	IC-321414-1	3.7	1.76	41.00	5.79	6.509	326.426
71.	IC-312410-1	4.4	1.91	40.50	6.48	8.483	427.269
72.	IC-321410	4.3	1.77	40.60	5.79	7.668	385.156
73.	IC-330987	5.0	1.65	40.00	5.59	8.322	418.657
74.	IC-331121	4.6	1.97	41.00	6.18	9.147	459.926
75.	IC-342080	3.9	2.03	44.60	6.26	7.984	402.047
76.	IC-546151	3.8	2.22	49.50	7.46	8.450	424.293
77.	IC-536594	4.4	1.95	48.90	7.10	8.658	435.659
<b>Mean</b>		<b>4.52</b>	<b>1.92</b>	<b>43.50</b>	<b>7.07</b>	<b>8.58</b>	<b>430.78</b>
<b>C.V</b>		<b>2.98</b>	<b>9.26</b>	<b>3.06</b>	<b>8.26</b>	<b>9.11</b>	<b>9.00</b>
<b>C.D. 5%</b>		<b>0.21</b>	<b>0.28</b>	<b>2.15</b>	<b>0.94</b>	<b>1.26</b>	<b>62.56</b>

#### 4. CONCLUSION

In the present study to evaluate the germplasm collection the mean of seventy-seven genotypes with two checks for 18 characters is presented in Tables 2, 3 and 4. Wide range of variation in mean performance of genotypes was observed for all the characters under study. Since no genotype could be identified to have superior performance for all the characters, the genotypes with diverse characteristics could be used in a well-planned hybridization programme to select superior performing lines in the successive segregating lines. Based on mean performance of the genotypes, it was observed that highest yield ha<sup>-1</sup> was recorded in IC-371747-1 (1132.21 q) followed by IC-382258 (848.19 q) and IC-398541 (845.33 q) whereas, the lowest yield ha<sup>-1</sup> was recorded in IC-522878 (211.83 q) followed by Pusa Naveen (220.30 q) and SKUA-BG-7 (241.78 q).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Deore SL, Khadabadi SS, Patel QR. In vitro antioxidant activity and quantitative estimation of phenolic content of *Lagenaria siceraria*. Rasayan Journal of Chemistry. 2009;2(1):129-132.
2. Singh AK. Cytogenetics and evolution in the cucurbitaceae. Cornell University, London. 1990;10-28.
3. Hemeda HM, Ibrahim HS. and Khattab HAH. Bakery fortification and biological evaluation of bottle gourd seeds (*Lagenaria siceraria*). Egyptian Journal of Food Science. 2008;36(1):93-113.
4. Parle M, Kaur S. Is bottle gourd a natural gourd. International Research Journal of Pharmacy. 2011;2(6):13-17.
5. USDA. National nutrient database for standard reference, release 1. The National Agricultural Library; 2018. Available:https://ndb.nal.usda.gov/ndb/
6. NHB, 2020-21. Indian Horticulture Database. Ministry of Agriculture. Government of India. Available:www.nhb.gov.in
7. Department of agriculture. J&K government. Area and production of bottle gourd in Jammu and Kashmir; 2020.
8. NHB, 2021-22. Indian Horticulture Database. Ministry of Agriculture. Government of India. Available:www.nhb.gov.in
9. Harika M, Gasti VD, Shantappa T, Mulge R, Shirol AM, Mastiholi AB, Kulkarni MS. Evaluation of bottle gourd genotypes [*Lagenaria siceraria* (Mol.) Standl.] for various horticultural characters. Karnataka Journal of Agricultural Sciences. 2012; 25(2): 241-244.
10. Jain Akshay, Singh SP. Evaluation on mean performance in Bottle Gourd (*Lagenaria siceraria* (Molina) Standl.) genotypes. Journal of Global Biosciences. 2016;5(8):4515-4519.
11. Rambabu E, Mandal AR, Hazra P, Senapati BK, Thapa U. Morphological characterization and genetic variability studies in bottle gourd [*Lagenaria siceraria* (Molina) Standl.]. International Journal of Current Microbiology and Applied Sciences. 2017;6(9):3585-3592.
12. Rashid M, Hussain K, Hussain SM, Farwah S, Javeed I, Sultan A, Azrah S, Akhter A, Ahmad M, Maqbool S. Comparative Performance of Various Bottle Gourd [*Lagenaria siceraria* (Molina) Standl.] Genotypes. International Journal of Current Microbiology and Applied Sciences. 2020; 9(6): 371-375. Available:https://doi.org/10.20546/ijcmas.2020.906.048
13. Kandasamy R, Arivazhagan E, Bharathi SS. Evaluation of growth and yield characters in bottle gourd (*Lagenaria siceraria* (Mol.) Standl.). Journal of Pharmacognosy and Phytochemistry. 2019;8(3):4653-4655.
14. Khan MM, Singh SP, Ram H, Singh VB, Mubeen. Genetic variability in bottle gourd. International Journal of Chemical Studies. 2020;8(3):2175-2177.
15. Sohi A, Prasad VM, Bahadur V, Topno SE. Hybrids Evaluation of Bottle Gourd [*Lagenaria siceraria* (Molina) Standl.] for Fruit Growth Yield Quality and Morphological Traits in Prayagraj Agro-Climatic Conditions. Biological Forum – An International Journal. 2021;13(2): 477-480.

16. Dubey A, Ram CN, Alam K, Shukla R, Pandey V. To study the genetic variability, heritability and genetic advance for agronomic traits of Bottle gourd [*Lagenaria siceraria* (Mol.) Standl]. The Pharma Innovation Journal. 2022;11(3):1362-1366.

© 2024 Bashir et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/112442>