



Effect of Seed Bioprimeing on Germination & Seedling Attributes in Chickpea (*Cicer arietinum L.*)

Lella Naga Swetha ^{a++*} and Abhinav Dayal ^{a#}

^a Department of Genetics and Plant Breeding, SHUATS, Prayagraj, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i23958

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111036>

Original Research Article

Received: 05/11/2023

Accepted: 10/01/2024

Published: 12/02/2024

ABSTRACT

The Study was conducted during Zaid season of 2022-2023 with IPC-05-59 genotype of chickpea (*Cicer arietinum L.*) Find out the Effect of seed bioprimeing on germination & seedling attributes in chickpea (*Cicer arietinum L.*). The design of experiment was Completely Randomized Design comprising treatments [viz.T0-uninoculated T1-Bacillus sp., T2- Mesorhizobium sp., T3- Pseudomonas sp, T4- Bacillus sp(0.5)+ Mesorhizobium sp.(0.5),T5- Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5), T6- Bacillus sp.(0.5)+Pseudomonas sp(0.5).+ Mesorhizobium sp.(0.5), T7- Bacillus sp,T8- Mesorhizobium sp, T9- Pseudomonas sp, T10- Bacillus sp.(0.5) + Mesorhizobium sp(0.5),T11-Pseudomonas sp. (0.5)+ Mesorhizobiumsp(0.5),T12- Bacillus sp.(0.5) + Pseudomonas sp. (0.5) + Mesorhizobium sp. (0.5) . The treated seeds along with control were evaluated for their morpho-physiological, growth, yield, root parameters and biochemical parameters under laboratory conditions, The study revealed that seeds bioprimeing techniques with Bacillus sp.(0.5)+ Pseudomonas sp(0.5).+ Mesorhizobium sp.(0.5) concentration @ 1.0ml for 1 hours T6 recorded significantly higher germination percent (93%), rate of germination (5.38), mean germination time

⁺⁺M. Sc Scholar;

[#]Assistant Professor;

*Corresponding author: E-mail: nagaswetharameshlella@gmail.com; nagaswethalella270699@gmail.com;

(11.16), root length (11.73cm), shoot length (13.27cm), seedling length (25.00cm), fresh weight (7.91gm), dry weight (1.62gm), vigor index-I (2324.91), vigor index-II (135.82), seed density (1.27), electrical conductivity (0.817), seed metabolic efficiency (1.018). which was followed by seed priming with T12- *Bacillus* sp (0.5) + *Pseudomonas* sp (0.5) + *Mesorhizobium* sp (0.5) @ 1.5ml for 2 hours and minimum was recorded in T0 (control).

Keywords: Chickpea; bioprime treatment; crbd (completely randomized block design); germination; speed of germination; seedling length; vigour index.

1. INTRODUCTION

Chickpea (*Cicer arietinum*.L) belongs to the family Fabaceae, subfamily Faboideae. It is commonly known by other names like gramor, Bengal gram, Egyptian pea, garbanzo bean or garbanzo. Chickpea have protein (23.3- 28.9%), carbohydrates (61.5%), fats (4.5%) and minerals (phosphorus, calcium, magnesium, iron, zinc). Seed priming is a physiological strategy that involves soaking of seeds in a solution of a specific priming agent followed by drying of seeds that initiates germination related process. This crop was cultivated in about 99Lha. The country harvested a record production of 107 Lt at a highest productivity level of 1086kg/ha. As usual, Madhya Pradesh has contributed a significant 28% of the total gram area and 34% of total gram production in the country, thereby ranking first both in area and production followed by Maharashtra (20% and 18%), Rajasthan (19% & 18%) and Karnataka (10% & 6%). About 97 percent of gram production of the country during the period under report has been realized by 10 states of Madhya Pradesh, Maharashtra, Rajasthan, Karnataka, Uttar Pradesh, Andhra Pradesh, Gujarat, Chhattisgarh and Jharkhand (DAC & FW, GOI, 2022).

Among all the different priming methods, the recent modern techniques bioprime which is a combination of beneficial microorganisms' application on surface of seed and seed hydration Upadhyay et al., [1] that integrates both biological and physiological methods to improve growth of plant, along with controlling diseases. Beneficial microorganisms like *Pseudomonas* and *Rhizobium* improve plant growth by colonizing on root surface, aiding in enhanced growth of plant. Chickpea is mostly consumed in the form of processed whole seed (boiled, roasted, parched, fried, steamed, sprouted etc) or dal or rasdal flour(besan). It is used in preparing a variety of snacks, sweets and condiments. It is mixed with wheat flour for "chapati" making. Fresh green seeds are consumed as green vegetable. Green leaves are used as vegetable. Grains are also used as vegetable (chole). Husk

and bits of dal are used as nutritious feed for animals. Chickpea can also be used as green fodder for animals. It is mainly grown in more than 50 countries including India, Pakistan, Turkey, Iran, Myanmar, Australia, Ethiopia, Canada, Mexico and Iraq Gaur et al., (2010).

Biofertilizers may colonize the rhizosphere and promotes growth by increasing the availability and supply of nutrients and/or growth stimulus to crop. Nitrogen fixer and phosphate solubilizing microorganisms play an important role in supplementing nitrogen and phosphorus to the plant, allowing a sustainable use of nitrogen and phosphate fertilizers.

Some important strains are mentioned as plant growth promoting rhizobacteria (PGPR) and that can be used as biofertilizers Kennedy et al., (2004) i.e. *Rhizobium*, *Azospirillum*, *Azotobacter*, *Bacillus*, *Burkholderia*, *Erwinia*, *Mycobacterium*, *Flavobacterium* etc. The bio-prime techniques changes physiological behavior of the host under stress more over significantly plays vital role to improve seeds tolerance level under environmental stress conditions [2]. Priming of seeds with helpful microorganisms and biocontrol means has been testified more proficiently for the management of diseases and pests as equated to other available methodologies (Prabha et al., 2019). The seed bioprime is a recently adapted method of seed priming. Seed priming is a pre-sowing treatment which leads to a physiological state that allows seed to germinate more proficiently. The preponderance of seed treatment are based on seed imbibition allowing the seeds to go through the first reversible stage of germination but does not allow radical protrusion through the seed coat. Seeds keeping their desiccation tolerance are then dehydrated and can be stored until final sowing (Stanley et al., 2016).

2. PHYSIOLOGICAL AND BIOCHEMICAL CHANGES IN SEEDS ON PRIMING

During seed germination an initial rapid uptake of water is followed by a visible lag phase Only after

the germination is completed a further increase in water up take occurs as the embryonic axis elongates. Movement of water across cell membranes is essential for the initiation of metabolism and the water movement is mediated by aquaporins. Transmembrane proteins are aquaporins (AQPs) which plays a vital role in plant water relations. High expression of AQPs during priming increases the water transport across the plasma membrane which facilitate water supply to the expanding tissues and enhance the germination potential of primed seeds. This may be the reason for the faster imbibition of water by primed seeds in comparison with non-primed ones. Priming leads to different biochemical and physiological changes in seeds. It synchronizes germination after breaking dormancy Nascimento et al., [3], diminishes the lag time required for imbibition, hydrolyses or metabolises inhibitors, activates enzymes, mobilises reserved food and enhances embryonic tissue outgrowth Rafi et al. [4]. Starch metabolism is of great importance during seed metabolism which influences seedling vigour under stress. This metabolism is brought about by α - amylases which hydrolyse the starch reserves into metabolisable sugars providing energy to the developing embryo. Seed priming enhances α - amylase and dehydrogenase activity that could hydrolyse the starch macro molecules in to smaller and simple sugars with increased ATP production and respiration. Phytase, amylase, and protease also increase during this process. Likewise Kamithi et al., [5] have shown increased activity of α -amylase in primed barley seeds by 2.8 times, whereas in primed wheat seeds activity was increased by 2.7 times compared to unprimed seeds. This might have led to enhanced germination events in primed seeds [6-15].

2.1 Bio-priming of Seeds in Seed Production

Seed bio-priming offers significant advantages to the seed manufacturing business. In the same way that genetic traits are passed down through generations, seed bacteria and other promising bio-gents are now being employed to increase seed vigour. In many crops, the addition of these helpful microorganisms to the seed and soil interface is a crucial strategy for producing high-quality seeds. Therefore, the seed producing sectors will be the first to gain from seed bio-priming techniques. By undergoing a number of structural, physiological, and biochemical changes within the plants, seed priming with

advantageous microorganisms or biocontrol agents increases the availability of nutrients to the plants and induces systemic resistance against biotic and abiotic stresses in various ecological conditions. These microorganisms encompass a variety of bacterial and fungal agents that encourage plant development.

Keeping in the view, the importance of crop and benefits of seed bio priming with microbes, the aim of the study is to identify the most effective agents for bio priming of chickpea seeds and their effect one mergence, growth and seed quality parameters.

2.2 Objectives

- To produce the inoculums of the indigenous bacterial isolates.
- To evaluate the effect of selected indigenous potential isolates on germination and seedling parameters of chickpea.

2.3 Justification

There is to be proper study regarding impact of seed biopriming treatment for improvement of plant growth attributes in chickpea, so this study helps us to investigate effect of bio priming seed treatment, Biofertilizers may colonizes the rhizosphere and promotes growth by increasing the availability and supply of nutrient sand or growth stimulus to crop on chickpea. Selecting appropriate seeds and treatments is key in crop production. Biofertilizers has shown better improvement of physiological and morphological growth attributes of crop growth.

3. MATERIALS AND METHODS

The materials, methodology and techniques adopted in the present experiment entitled, "Effect of seed biopriming on germination & seedling attributes in chickpea (*Cicer arietinum* L.)" with a brief description regarding the site of the experiment, temperature conditions, sampling techniques and statistical analysis are presented in this chapter with following headings.

3.1 Experimental Design

The experiment consists of 13 treatments including control which are replicated four times and allocated in Completely Randomized Block Design in each replication.

List 1. Details of layout

Crop	Chickpea (<i>Cicer arietinum L.</i>)
Genotype	IPC-05-59
Season	Zaid2022-23
Number of Replications	4
Number of Treatments	13(12+1)
Number of seeds per replication	100
Method of germination	Between the paper method
First count	4thDAS
Final Count	8thDAS

List 2. Treatment Combinations

Treatments	Organisms	Concentrations	Duration (hours)
T0	Uninoculated	NIL	-
T1	<i>Bacillus sp.</i>	1ml	1
T2	<i>Mesorhizobium sp.</i>	1ml	1
T3	<i>Pseudomonas sp.</i>	1ml	1
T4	<i>Bacillus sp(0.5)+Mesorhizobium sp.(0.5)</i>	1ml	1
T5	<i>Pseudomonas sp(0.5)+Mesorhizobiumsp.(0.5)</i>	1ml	1
T6	<i>Bacillus sp (0.5) +Pseudomonas sp(0.5).+ Mesorhizobium sp.(0.5)</i>	1ml	1
T7	<i>Bacillussp.</i>	1.5ml	2
T8	<i>Mesorhizobium sp.</i>	1.5ml	2
T9	<i>Pseudomonas sp.</i>	1.5ml	2
T10	<i>Bacillus sp(0.5)+Mesorhizobium sp(0.5)</i>	1.5ml	2
T11	<i>Pseudomonas sp(0.5).+ Mesorhizobium sp.(0.5)</i>	1.5ml	2
T12	<i>Bacillus sp (0.5) + Pseudomonas sp (0.5) + Mesorhizobium sp (0.5)</i>	1.5ml	2

3.2 Preparation of Solutions

Bacterial cultures:

Casine hydrolysate-10gm/litre Protease, peptone-10gm/litre

K₂HPO₄ - 1.5gm/litre

Mg So₄ .7H₂O - 1.5gm/litre pH- 7

3.3 Preparation of Media

For *Bacillus sp* Nutrient broth media will be used which is a composition of Manitol agar - 10 gm/litre

K₂HPO₄ - 0.5gm/litre
MgSO₄ - 0.2gm/litre
NaCl - 0.1 gm/litre
CaCO₃ - 4gm/litre
Yeast extract - 0.1gm/litre

For *Rhizobium sp* Yama broth media will be used which is a composition of:

Peptone - 5 gm/litre
Beef extract - 3gm/litre
Nacl- 5gm/litre

For *Pseudomonas sp*: pseudomonas agaris used which is a composition of:

- Autoclavemediaat120°C for 15min.
- After media gets cooled down fresh culture of *Bacillus spp*, *Pseudomonas spp*, *Rhizobium spp*. A reseparately inoculated in specific broth.
- Incubate at 28°C for2-3days.
- Seeds were surface disinfected with ethanol
- The sterilized seeds were soaked bacterial sustention (1×10^7 cells/ml) for 1 ,2 hours respectively, in case of control, the seeds are soaked in distilled water.
- Seeds were air driedfor48hours
- Seeds were placed on each germination paper for germination tests
- Observations were recorded on first count and last count ie on 4thand 8th day after kept for germination.

4. RESULTS AND DISCUSSION

An experiment entitled “Effect of seed bioprimering on germination & seedling attributes in chickpea (*Cicer arietinum L.*)” was conducted at Seed Testing Laboratory, Department of Genetics and Plant Breeding, Naini Agricultural Institute, SHUATS, Prayagraj, (U.P). during Zaid season of 2022-2023. The results of this experiment were presented in the following pages under appropriate sub headings. Experimental findings are discussed in the light of scientific reasoning and their conformity with the previous researchers [17-20].

Analysis of Variance: The analysis of variance for Germination and Seedling Attribute sin chickpea was given in Table 1. Analysis of variance revealed that the differences among thirteen treatments (Table 1) were significant for germination and seedling attributes viz., Germination percentage (%), Rate of germination, Mean germination time, Root length (cm), Shoot length(cm), Seedling length (cm), Fresh weight(g), Dry weight (g), Vigour Index-I, Vigour Index-II, Seed density, Electrical conductivity (dSm-1), Seed metabolic efficiency (Table 2) [21-25].

4.1 Germination Percentage

A perusal of data on germination percentage was recorded during the experimental period revealing a significant difference in germination percent, which was recorded As per the data tabulated germination percent was recorded significantly higher in treatment (T6 - *Bacillus* sp.(0.5)+ *Pseudomonas* sp (0.5).+ *Mesorhizobium* sp.(0.5) @ 1ml) with 93% and treatment (T12- *Bacillus* sp.(0.5)+ *Pseudomonas* sp. (0.5)+*Mesorhizobium* sp.(0.5)@1.5ml) with 89%was statistically at par with treatment T6 and the lowest germination percent was recorded in treatment T0 (control) with 68%.

Pseudomonas strains are known to produce plant growth-promoting hormones such as auxins and cytokines. These hormones can stimulate seed germination and early seedling growth by promoting cell elongation, root development, and nutrient uptake. *Pseudomonas* treatments can induce systemic resistance in plants. This means that chickpea seeds treated with *Pseudomonas* may become more resilient to stressors and pathogens, there by improving their chances of germination and early growth.

4.2 Rate of Germination

From the data recorded it is obvious that treatment (T6 - *Bacillus* sp. (0.5) + *Pseudomonas* sp(0.5).+ *Mesorhizobium* sp.(0.5) @ 1ml) gave significantly higher rate of germination with 5.38% over all other treatments. However, treatment (T12- *Bacillus* sp (0.5) + *Pseudomonas* sp. (0.5) + *Mesorhizobium* sp. (0.5) @1.5ml) recorded rate of germination of 5.23% and was statistically at par with T3 and the lowest rate of germination was obtained in T0 (control) with 3.77%.

4.3 Mean germination time

As per the data recorded the treatment (T6 - *Bacillus* sp.(0.5)+ *Pseudomonas* sp(0.5).+ *Mesorhizobium* sp.(0.5) @ 1ml) had significantly highest mean germination time of 11.16 over all other treatments and treatment T12-*Bacillus* sp.(0.5)+ *Pseudomonas* sp.(0.5)+ *Mesorhizobium* sp.(0.5) @ 1.5ml with 10.68 of mean germination time was statistically at par with T3. The lowest mean germination time was recorded in T0 (control) with 8.16.

4.4 Root Length

From the data recorded it is obvious that treatment (T6 - *Bacillus* sp. (0.5) + *Pseudomonas* sp (0.5) + *Mesorhizobium* sp. (0.5) @ 1ml) gave significantly higher root length with 11.73cm over all other treatments considered. Treatment (T12-*Bacillus* sp. (0.5) + *Pseudomonas* sp. (0.5) + *Mesorhizobium* sp. (0.5) @1.5ml) recorded a root length of 11.60cm and was statistically at par with T6 and the lowest root length was obtained in T0 (control) with 6.64cm [26-30].

Pseudomonas treatments can stimulate the development of root hairs, which are extensions of root epidermal cells. Root hairs increase the surface area for nutrient and water absorption, enhancing Overall root growth. *Pseudomonas* bacteria can trigger the production of antioxidants in chickpea seedlings. Antioxidants protect plant tissues from oxidative stress, allowing roots to grow without hindrance.

4.5 Shoot Length

As per the data recorded the treatment (T6 - *Bacillus* sp. (0.5) + *Pseudomonas* sp (0.5) + *Mesorhizobium* sp (0.5) @ 1ml) had significantly highest shoot length of 13.27cm over all other treatments and treatment T12- *Bacillus* sp. (0.5) + *Pseudomonas* sp. (0.5) + *Mesorhizobium* sp.(0.5) @ 1.5 ml with 13.27cm of shoot length was statistically at par with T6. The lowest shoot length was recorded in T0 (control) with 7.67cm.

Table 1. Analysis of variance for seed germination and seedling parameters of bio-priming by *Rhizobium*, *Bacillus*, *Pseudomonas* in chickpea (*Cicer arietinum L.*)

S.no	Characters	Mean sum of squares Treatments (df=12)	Error (df=39)
1	Germination percentage (%)	125.4231*	4.0000
2	Rate of germination	0.5409*	0.4288
3	Mean germination time	1.8061*	0.0576
4	Root length(cm)	7.8748*	0.0112
5	Shoot length(cm)	7.4281*	0.0617
6	Seedling length(cm)	29.8297*	0.0921
7	Fresh weight(g)	4.8217*	0.0999
8	Dry weight(g)	0.0296*	0.0051
9	Vigour index-I	418188.5837*	283.6725
10	Vigour index-II	527.8566*	42.1658
11	Seed density	0.0032*	0.0013
12	Electrical conductivity(dSm-1)	0.5077*	0.0338
13	Seed metabolic efficiency	0.0545*	0.0192

*Indicates 5% Level of Significance

Table 2. Mean Performance of germination and seedling parameters of bio-primed seeds by *Rhizobium*, *Bacillus*, *Pesudomonas* in Chickpea (*Cicer arietinum L.*) Genotype: IPC-05-59

Treatments	Germination percentage (%)	Rate of germination	Mean germination time	Root length (cm)	Shoot Length (cm)	Seedling length (cm)	Fresh Weight (g)	Dry weight (g)	Vigour index I	Vigour index II	Seed density	Electrical conductivity	Seed metabolic efficiency
T0	68	3.77	8.16	6.64	7.67	14.32	3.07	1.25	973.18	84.71	1.15	1.905	0.556
T1	78	4.66	9.36	9.31	10.95	20.27	4.54	1.35	1580.45	105.72	1.21	1.766	0.823
T2	79	4.76	9.48	10.4	10.65	21.05	3.81	1.36	1662.81	128.26	1.21	1.686	0.690
T3	81	4.95	9.72	7.49	9.15	16.65	4.21	1.39	1348.03	112.68	1.19	1.625	0.789
T4	82	4.95	9.84	10.14	10.77	20.91	4.88	1.46	1714.33	126.28	1.20	1.078	0.722
T5	86	5.10	10.32	10.62	12.41	23.04	5.46	1.51	1980.81	130.21	1.23	0.902	0.591
T6	93	5.38	11.16	11.73	13.27	25.00	7.91	1.62	2324.91	135.82	1.27	0.817	1.018
T7	73	4.23	8.76	6.85	8.59	15.45	3.26	1.34	1127.26	106.24	1.17	1.848	0.861
T8	79	5.09	9.48	9.77	10.51	20.29	3.8	1.36	1602.14	109.57	1.24	1.734	0.862
T9	80	4.95	9.60	9.21	10.77	19.99	5.53	1.39	1598.59	114.2	1.20	1.686	0.841
T10	82	4.81	9.84	8.85	10.43	19.29	4.04	1.43	1581.16	111.48	1.22	1.331	0.650
T11	85	4.81	10.20	9.41	11.19	20.61	4.59	1.46	1751.22	114.94	1.19	1.002	0.772
T12	89	5.23	10.68	11.60	12.63	24.23	5.51	1.54	2156.37	119.02	1.25	0.891	0.948
Grand mean	81	4.82	9.74	9.39	10.69	20.08	4.66	1.42	1646.25	115.66	1.21	1.406	0.779
S.Em	1.155	0.378	0.139	0.061	0.143	0.175	0.182	0.041	9.724	3.749	0.021	0.106	0.080
SE(d)	1.633	0.535	0.196	0.086	0.203	0.248	0.258	0.058	13.752	5.302	0.029	0.150	0.113
C.D(0.05)	3.357	1.099	0.403	0.178	0.417	0.509	0.530	0.120	28.267	10.898	0.060	0.309	0.233

4.6 Seedling Length

The significantly higher seedling length (25.00cm) was recorded with treatment (T6 - Bacillus sp. (0.5)+ Pseudomonas sp (0.5).+ Mesorhizobium sp.(0.5) @ 1ml) over the rest of the treatments taken. In contrast a significantly lower seedling length of 14.32cm was recorded with treatment (T0- control). However, treatment (T12- Bacillus sp. (0.5) + Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5 ml) with 24.23cm was statistically at par with T6.

Pseudomonas treatments can trigger the production of antioxidants in chickpea seedlings. Antioxidants protect plant tissues from oxidative stress, which can hinder growth. With lower oxidative stress, seedlings can focus on elongating their stems and leaves. Pseudomonas treatments can induce systemic resistance in chickpea seedlings. When the seedlings are primed for defense responses, they allocate more resources to growth, resulting in increased seedling length.

4.7 Fresh Weight

The significantly higher seedling fresh weight was obtained in treatment (T6 - Bacillus sp.(0.5)+ Pseudomonas sp (0.5).+Mesorhizobium sp.(0.5) @1ml) with 7.91gm among all the other treatments and the lowest seedling fresh weight was recorded in T0 (control) with 3.07gm respectively. The treatment (T12- Bacillus sp. (0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml) was statistically at par with T6 with 5.51gm.

Pseudomonas treatments can promote more efficient nutrient uptake by the seedlings. This increased nutrient absorption can support the development of larger and heavier seedlings. Pseudomonas can produce plant growth-promoting hormones like auxins and cytokinins. These hormones stimulate cell division and elongation, leading to increased biomass and fresh weight in seedlings. Pseudomonas treatments can induce systemic resistance in chickpea seedlings. As a result, seedlings are better protected against pathogens and environmental stresses, allowing them to allocate more energy to growth and fresh weight gain.

4.8 Dry Weight

The significantly higher seedling dry weight (1.62gm) was recorded with treatment (T6 - Bacillus sp (0.5) + Pseudomonas sp. (0.5).+ Mesorhizobium sp.(0.5) @ 1ml). In contrast a significantly lower dry weight of 1.25 was

recorded with treatment (T0-control). However, treatment (T12- Bacillus sp. (0.5) +Pseudomonas sp. (0.5) + Mesorhizobium sp. (0.5) @1.5ml) with 1.54gm was statistically at par with treatment T2.

Pseudomonas treatments often lead to the development of more extensive and well-branched root systems. A healthy root system allows for better nutrient and water absorption, ultimately contributing to increased seedling dry weight. Pseudomonas treatments may enhance the efficiency of photosynthesis in chickpea seedlings, leading to increased production of carbohydrates and dry biomass.

4.9 Vigour Index I

Significantly higher vigour index I was recorded in treatment T6 - Bacillus sp. (0.5)+ Pseudomonas sp (0.5).+ Mesorhizobium sp.(0.5) @ 1ml with 2324.91 and the treatment T12-Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml with vigour index I of 2156.37 was statistically at par with T6. The lowest vigour index I was recorded in treatment T0-control with vigour index I of 973.18.

Pseudomonas treatments often lead to the development of a more extensive and well-branched root system. A healthy root system enhances nutrient and water uptake, supporting overall seedling vigor. Pseudomonas treatment promotes germination, seedling length, and dry weight through better cell division increasing the seedling vigour.

4.10 Vigour Index II

As per the data recorded during laboratory study the vigour index II was recorded significantly higher in treatment T6 - Bacillus sp. (0.5)+ Pseudomonas sp(0.5).+ Mesorhizobium sp.(0.5) @ 1ml with vigour index II of 135.82 and statistically at par value was recorded in treatment T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @1.5ml with 130.21. The lowest data on vigour index II was recorded in treatment (T0-control) with 84.71.

4.11 Seed Density

The treatments varied significantly and the highest seed density was recorded in treatment T6 - Bacillus sp.(0.5)+ Pseudomonas sp(0.5).+ Mesorhizobium sp.(0.5) @ 1ml with 1.27 dsm^{-1} . The treatment (T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5)

@ 1.5ml) with seed density of 1.25 dsm^{-1} was statistically at par with T6 and the lowest seed density was recorded in treatment T0 - control with 1.15 dsm^{-1} .

Pseudomonas treatments can stimulate the development of a more extensive root system in chick pea seedlings. These well-developed root systems enable seedlings to explore a larger soil volume for resources, indirectly contributing to their overall vigor and potential seedling density.

4.12 Electrical Conductivity

As per the data tabulated electrical conductivity was recorded significantly lower in treatment (T6 – Bacillus sp. (0.5) + Pseudomonas sp (0.5). + Mesorhizobium sp. (0.5) @ 1ml) with 0.817 and treatment (T12-Bacillussp. (0.5)+ Pseudomonas sp. (0.5)+ Mesorhizobiumsp. (0.5) @ 1.5 ml) with 0.891 was at par with T6 and the highest electrical conductivity was recorded in treatment (T0-control) with 1.905.

Pseudomonas seed treatment may influence the ion-selective transport systems in chickpea root cells. These bacteria can modulate ion channels and transporters to favor the uptake of essential nutrients (e.g., potassium and calcium) while limiting the uptake of potentially detrimental ions (e.g., sodium and chloride). This selective ion regulation can lead to an altered ion composition within the plant and, consequently, a lower EC of the plant sap.

4.13 Seed Metabolic Efficiency

A perusal of data on seed mobilization efficiency was recorded during the experimental period on different treatments revealing a significant difference in seed mobilization efficiency, As per the data tabulated seed mobilization efficiency was recorded significantly higher in treatment (T6 - Bacillus sp.(0.5)+ Pseudomonas sp (0.5).+ Mesorhizobium sp.(0.5) @ 1ml) with 1.018 and treatment (T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml) with 0.948 was statistically at par with T6 and the lowest seed mobilization efficiency was recorded in treatment(T0- control) with 0.556.

4.14 Summary

The present lab experiment entitled ““Effect of seed biopriming on germination & seedling attributes in chickpea (*CicerarietinumL*)” was conducted at Seed Testing Laboratory, Department of Genetics and Plant Breeding, Naini Agricultural Institute, SHUATS, Prayagraj, (U.P). during Zaid season of 2022-2023.

The study conducted in the seed testing laboratory revealed that germination percent (93%), rate of germination (5.38), mean germination time(11.16),root length (11.73cm), shoot length(13.27cm), seedling length (25.00 cm), fresh weight (7.91 gm), dry weight (1.62 gm), vigor index-I (2324.91), vigor index – II (135.82), seed density(1.27), electrical conductivity (0.817), seed metabolic efficiency (1.018) were recorded significantly higher in T6 - Bacillus sp.(0.5)+ Pseudomonas sp (0.5).+ Mesorhizobium sp.(0.5) @ 1ml over other treatments and treatment T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml was at par with T6.

- Analysis of variance showed significant difference among the treatments for the characters studied of the
- Chickpea which indicates significant effect of the treatment son seedling parameters of the crop.
- The salient finding of the experiment is summarized and conclusion is drawn. The study was undertaken to know the effect of seed biopriming treatments on Chick pea seeds have been presented and discussed in the previous chapter. The data was recorded for seed ling parameters of Chick pea.

5. CONCLUSIONS AND RECOMMENDATIONS

Studying indigenous bacteria can improve plant health and plant growth and Inoculating seed with beneficial indigenous bacteria can enhance nutrient availability and protect plants from harmful pathogens. Indigenous bacteria can provide insights into the local microbial community and its role in various ecosystems. They contain unique enzymes or metabolic pathways that can be harnessed for bio technological applications in crop.

It is concluded from the present study that (T6) treatment in which seeds treated with Bacillus sp.(0.5)+ Pseudomonas sp(0.5).+ Mesorhizobium sp.(0.5) @ 1ml had out performed all treatments in laboratory conditions with highest germination percentage, seedling length, fresh weight, vigor indices, electrical conductivity followed by (T12) treatment in which seeds treated with Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5)

@ 1.5ml and the lowest were recorded in (T0-control).

6. FUTURE PROSPECTS

The bio priming technique can be recommended for priming seeds prior to field planting as an environmentally friendly strategy to improve seed germination and initial seedling growth. Extensive field trials under different environmental conditions, using several seed lots of cultivars, will be necessary in order to establish the efficiency of these strains as potential bio-priming seed treatments for improving Crop productivity

ACKNOWLEDGEMENT

The authors are thankful to Dr. Abhinav Dayal, Assistant Professor, Department of Genetics, SHUATS, Prayagraj, U.P. for providing the support to make this study success.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh V, Upadhyay RS, Sarma BK, Singh HB. Seed bio-priming with trichoderma a sperellum effectively modulate plant growth promotion in pea. International Journal of Agriculture, Environment and Biotechnology. 2016;9(3):361- 365.
2. Entesari M, Sharifzadeh F, Ahmadzadeh M, Farhangfar M. Seed biopriming with trichoderma species and Pseudomonas fluorescent on growth parameters, enzymes activity and nutritional status of soybean International Journal of Agronomy and Plant Production. 2013;4(4):610-619.
3. Nascimento WM, Cantliffe DJ, Huber DJ. Seed aging affects ethylene production and endo- β -mannanase activity during lettuce seed germination at high temperature. Seed Science and Technology. 2005;33(1):11-17.
4. Rafi H, Dawar S, Zaki MJ. Seed priming with extracts of *Acacia ni lotica* (L.) willd. ex delile and *Sapindus mukorossi* (L.) plant parts in the control of root rot fungi and growth of plants. Pak. J. Bot. 2015;47(3):1129-1135.
5. Kamithi KD, Wachira F, Kibe AM. Effects of different priming methods and priming durations on enzyme activities in germinating chickpea (*Cicer arietinum* L.). American Journal of Natural and Applied Sciences. 2016;1(1):1-9.
6. Abdul-Baki A, Anderson JD. Vigor determination in soybean seed by multiple criteria. Crop Science. 1973;13(6):630-633.
7. Baliyan N, Dheeman S, Maheshwari DK, Dubey RC, Vishnoi VK. Rhizobacteria isolated under field first strategy improved chickpea growth and productivity. Environmental Sustainability. 2018;1:461-469.
8. Chaurasia AK, Naik BP, Bara BM. Influence of priming on seed yield parameters of chickpea (*Cicer arietinum* L.). Journal of Pharmacognosy and Phytochemistry. 2018;7(2):753-755.
9. Ditta A, Imtiaz M, Mehmood S, Rizwan MS, Mubeen F, Aziz O, Qian Z, Ijaz R, Tu S. Rock phosphate-enriched organic fertilizer with phosphate-solubilizing microorganisms improves nodulation, growth, and yield of legumes. Commun. Soil Sci. Plant Anal. 2018;49(21):2715–2725.
10. Fahsi N, Mahdi I, Mesfioui A, Biskri L, Allaoui A. Plant growth promoting rhizobacteria isolated from the jujube (*Ziziphus lotus*) plant enhance wheat growth, Zn uptake, and heavy metal tolerance. Agriculture. 2021;11:316.
11. Gupta SC. Effect of combined inoculation on nodulation, nutrient uptake and yield of chickpea in Vertisol. Journal of the Indian Society of Soil Science. 2006;54:251-254.
12. Hayat R, Ali S, Amara U, Khalid R, Ahmed I. Soil beneficial bacteria and their role in plant growth promotion: A review. Annals of Microbiology. 2010;60:579-598.
13. Jeffries P, Gianinazzi S, Perotto S, Turnau K, Barea JM. The contribution of arbuscular mycorrhizal fungi in sustainable maintenance of plant health and soil fertility. Biology and Fertility of Soils. 2003;37:1-16.
14. Khan MS, Zaidi A, Wani PA. Role of phosphate solubilizing microorganisms in sustainable agriculture. A Review. Agronomy for sustainable Development. 2007;27:29- 43.
15. Lohitha B, Dutta A. Bioprimer in chickpea (*Cicer arietinum* L.): An ecofriendly

- approach for quality seed production. *Pharma Innovation.* 2022;11(2):375-379.
16. Moeinzadeh A, Sharif-Zadeh F, Ahmadzadeh M, Tajabadi FH. Bioprimering of sunflower (*Helianthus annuus* L.) seed with *Pseudomonas fluorescens* for improvement of seed invigoration and seedling growth. *Australian Journal of Crop Science.* 2010;4(7):564-570.
17. Negi S, Bharat NK, Kaushal R, Rohiwala P. Screening of bioagents for seed bioprimering in French bean (*Phaseolus vulgaris* L.) under Laboratory condition. *International Journal of Communication Systems.* 2020;8(3):790-793.
18. Pandey S, Gupta S. Evaluation of *Pseudomonas sp.* for its multifarious plant growth promoting potential and its ability to alleviate biotic and abiotic stress in tomato (*Solanum lycopersicum*) plants. *Scientific Reports.* 2020;10:20951.
19. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. Indian council of agricultural research, New Delhi. 1967;381.
20. Pramanik K, Bera AK. Response of biofertilizer and phytohormone on growth and yield of chickpea (*Cicer arietinum* L.). *Journal of Crop and Weed.* 2012;8(2):45-49.
21. Rao DG, Sinha SK. Efficiency of mobilization of seed reserves in sorghum hybrids and their parents as influenced by temperature regimes. *Seed Research.* 1993;2(2):97-100.
22. Reddy ASR, Madhavi GB, Reddy KG, Yella Reddy Gari SK, Reddy MS. Effect of seed bioprimering with *Trichoderma viride* and *Pseudomonas fluorescens* in chickpea (*Cicer arietinum*) in Andhra Pradesh, India. In *Plant growth-promoting rhizobacteria (PGPR) for sustainable agriculture, Proceedings of the 2nd Asian PGPR conference, Beijing, China.* 2011;324-429.
23. Tambekar DH, Gulhane SR, Somkuwar DO, Ingle KB, Kanchalwar SP, Upadhye MA, Bidwai UA. Potential Rhizobium and phosphate solubilizers as a biofertilizers from saline belt of Akola and Buldhana District, India. *Res. J. Agric. Biol. Sci.* 2009;5(4):578-582.
24. Tanwar A, Aggarwal A, Kaushish S, Chauhan S. Interactive effect of AM fungi with *Trichoderma viride* and *Pseudomonas fluorescens* on growth and yield of broccoli. *Plant Protection Science.* 2013;49(3):137-145.
25. Valverde A, Burgos A, Fiscella T, Rivas R, Igual JM. Differential effects of coinoculations with *Pseudomonas jessenii* PS06 (aphosphate-solubilizing bacterium) and *Mesorhizobium ciceri* C-2/2 strains on the growth and seed yield of chickpea under green house and field conditions. *Plant and Soil.* 2006;287:43-50.
26. Van der Maesen LJG. Origin, history and taxonomy of chickpea. In *The Chickpea.* 1987;11- 34.
27. Veena Shri Jainapur, Shalini N. Huilgol SM. Vastrad and Jolli RB. Bioprimering of chickpea seeds with biocontrol agents for enhanced seedling vigour and reduced seed borne diseases. *International Journal of Current Microbiology and Applied Sciences.* 7(2): 2746-2750.
28. Vishwas S, Chaurasia AK, Bara BM, Debnath A, Parihar NN, Brunda K, Saxena R. Effect of priming on germination and seedling establishment of chickpea (*Cicer arietinum* L.) seeds. *Journal of Pharmacognosy and Phytochemistry.* 2017;6(4):72-74.
29. Vishwas S, Chaurasia AK, Bara BM, Debnath A, Parihar NN, Brunda K, Saxena R. Effect of priming on germination and seedling establishment of chickpea (*Cicer arietinum* L.) seeds. *Journal of Pharmacognosy and Phytochemistry.* 2017;6(4):72- 74.
30. Yadav SK, Dave A, Sarkar A, Singh HB, Salma BK. Co-inoculated bioprimering with *Trichoderma*, *Pseudomonas* and *Rhizobium* improves crop growth in *Cicer arietinum* and *Phaseolus vulgaris*. *International Journal of Agriculture, Environment and Biotechnology.* 2013;6: 255-259.

© 2024 Swetha et al; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111036>