



Assessment of Genetic Variability, Heritability and Genetic Advance for Seed Yield and Its Contributing Traits in Greengram [*Vigna radiata* (L.) Wilczek]

Niyati Jain ^{a*}, R. S. Sikarwar ^a, M. K. Tripathi ^a,
Sushma Tiwari ^a, Shivangi Rahangdale ^b and Akash Barela ^b

^a Department of Genetics and Plant Breeding, College of Agriculture, Rajmata Vijayaraje Scindia Agricultural University, Gwalior, Madhya Pradesh, India.

^b Department of Genetics and Plant Breeding, College of Agriculture, Jawaharlal Nehru Agricultural University, Jabalpur, Madhya Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i34037

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112965>

Original Research Article

Received: 26/12/2023

Accepted: 01/03/2024

Published: 06/03/2024

ABSTRACT

Vigna radiata L. Wilczek, sometimes known as greengram or mungbean, is a popular pulse crop in India. Most of its cultivation takes place in India's subsistence farming systems. It is cultivated in three distinct seasons in India, despite the extremely low average productivity. Due to the crop's extensive growth in a variety of environmental circumstances, the local germplasm exhibits a high degree of genetic variation. In order to determine the genetic variability, heritability, and genetic

*Corresponding author: E-mail: j.niyati21@gmail.com;

advancement of 40 green gram genotypes, a study utilizing twelve quantitative variables was conducted. For every traits, significant variance was discovered using basic descriptive statistics. High Phenotypic Coefficients of Variation (PCV) and high Genotypic Coefficients of Variation (GCV) were identified for seed yield per plant, number of branches per plant, number of clusters per plant, biological yield per plant, number of pods per plant and 100 seed weight. Moderate PCV and GCV values observed for number of seeds per pod, plant height, harvest index, pod length and days to 50% flowering, whereas low PCV and GCV values were observed for days to maturity. Most of the characters studied in the current study had high estimates of heritability coupled with high to moderate genetic advance as a percentage of mean. This may be attributed to the preponderance of additive gene action and their high selective value, meaning that selection pressure can be applied profitably to these characters to improve them rationally.

Keywords: Green gram; genetic advance; genetic variability; heritability; genetic advance mungbean.

1. INTRODUCTION

Native to India and Southeast Asia, the green gram, or *Vigna radiata* (L.) Wilczek, is a legume crop that is widely grown. Usually, it is grown in the rainy season. Because of its brief growing season, it is also cultivated in the wheat-rice cropping system in the spring and summer to increase pulse production and crop intensity. Despite having the capacity to be cultivated in three seasons, the crop's average productivity worldwide is a pitiful 0.5 t ha⁻¹, far less than the 2.5 to 3.0 tonnes per hectare estimated yield potential [1]. Due to ongoing limitations on pulse availability, pulse costs have increased, rendering them unaffordable for individuals who are depressed. Germplasm accessions have accumulated several advantageous characteristics, like tolerance to a variety of weather conditions such as a preponderance of infections, temperature stress, or drought, among others. Thus, the evolution of the crop is still greatly influenced by natural selection. The crop has developed a diversity of agronomic features, such as yield and its determinants, pod number, biological yield, maturity time, and so forth. Assessing germplasm accessions is the most important step towards utilizing the variety that is available. Within this research, the significant yield-related traits of the green gram germplasm were evaluated. The results, which include the spectrum of genetic variability and genetic advancement appropriate for specific qualities, could support initiatives for agricultural improvement.

2. MATERIALS AND METHODS

The current study was conducted in the summer of 2022 at the College of Agriculture's Department of Plant Breeding and Genetics. 30 x

10 cm seed spacing was used to grow 40 germplasm units in a randomized block design with three replications. Plant protection measures that were based on need and recommended agronomic procedures were implemented. Data were gathered on twelve yield and yield contributing characters, which included days to 50% flowering, days to maturity, plant height(cm), number of branches per plant, number of clusters per plant, number of pods per cluster, pod length(cm), number of seeds per pod, 100 seed weight(g), harvest index, biological yield per plant and seed yield per plant (g). Five plants of various genotypes were chosen at random, tagged, and observed. The mean value was then computed for analysis over three replications. Burton's recommendations for doing an analysis of variance were followed [2]. The heritability formula provided by Hanson et al. [3] and Johnson et al. [4] was used to estimate heritability and genetic advance estimates, respectively. According to Johnson et al. [4] the percentages of genetic advance ranged from lower (<10%) to high (>20%) and moderate (10–20%).

3. RESULTS AND DISCUSSION

3.1 Analysis of Variance

For every variable under investigation, the analysis of variance revealed substantial variances (Table 1). To begin the selection process, these differences alone are adequate.

3.2 Per se Performance and Range

With a mean of 36 days, the days to 50% flowering ranged from 26 to 41 days. The range of days to maturity was 57 to 73 days, with an average of 65 days. With an average performance of 33.20 cm, the plant height's

degree of dispersion varied from 21.13 to 44.57 cm. The mean number of branches was 2.02, with a range of 1.25 to 3.95. The average number of clusters per plant was 5.24, with a range of 2.67 to 8.93. Plants had an average of 15.55 pods, ranging from 6.43 to 21.93. Between 5.8 and 9.67 inches, the average pod length was 7.82. Number of seeds per pod ranged from 4.82 to 12.80 with a mean of 9.69. With a minimum range of 1.84g and a maximum range of 5.61g, the average weight of 100 seeds was 3.57g. Plants produced an average of 17.64g of biological output, ranging from 6.08g to 27.8g respectively. Between 17.41% and 38.87%, with a mean of 30%, was the harvest index. The range of 1.58g to 8.55g was observed in the average seed yield per plant, 5.28g. The mean performance serves as a direct criterion for selecting the best genotypes from the diverse population. All it takes to raise the population's yield is to choose for traits associated with yield. The mean performance according to attributes is shown in Table 2.

3.3 Coefficient of Variation

The reliable measures that will enable us to evaluate the amount of variability and comprehend its heritable and non-heritable components are the genotypic and phenotypic coefficient of variations. The difference between the phenotypic and genotypic coefficient of variations indicates the percentage of environmental effect on the variability presented in the genotypes. The environment affects all biometrical aspects, as evidenced by the fact that PCV levels in this study are higher than GCV values. Table 2 presented the coefficients of variation. The high PCV values (Table 2) were number of branches per plant (32.11%), seed yield per plant (30.99%), biological yield per plant (27.25%), number of clusters per plant (26.75%), number of pods per plant (26.22%) and 100 seed weight (24.85%). Moderate PCV values were observed for number of seeds per pod (17.74%), harvest index (16.74%), plant height (14.83%), pod length (12.19%) and days to 50% flowering (10.22%), whereas low PCV values were observed for days to maturity (6.13%). In general, phenotypic coefficients of variation (PCV) values were higher than the genotypic coefficients of variation (GCV) for the respective characters. The high GCV values (Table 2) were reported for seed yield per plant (29.8%), number of branches per plant (29.78%), number of clusters per plant (25.96%), biological yield per

plant (25.6%), number of pods per plant (24.8%) and 100 seed weight (24.66%). Moderate GCV values were observed for number of seeds per pod (16.66%), plant height (13.96%), harvest index (13.96%), pod length (11.58%) and days to 50% flowering (10%), whereas low GCV values were observed for days to maturity (6%). These observations are in agreement with the findings of Wesly et al. [5], Sineka et al. [6], Parsaniya et al. [7] and Rahangdale et al. [8].

High coefficients of variation were identified for seed yield per plant, number of branches per plant, number of clusters per plant, biological yield per plant, number of pods per plant and 100 seed weight. This shows that these parameters may be enhanced by efficient selection. Therefore, these characters may be considered throughout the selection process to increase yield.

3.4 Heritability (h^2) and Genetic Advance as Per Cent of Mean (GAM)

Heritability estimates tell us how similar parents and children are to one another and explain the traits that are inherited. Any selection program must be heritable for the features to manifest in a comparable phenotypic way in the progeny of the selected individuals. Heritability alone, however, is not enough to genetically alter characteristics. It should also have a high level of genetic advancement, which is a gauge of how effectively a certain trait has evolved by suitable selection. Thus, for selection to be effective, there must be both high heritability and a significant genetic progress as a percentage of mean for the genetic development of the traits. Genetic advance and heritability values were shown in Table 2.

Highest heritability (bs) (Table 2) was recorded for 100 seed weight (98.48%) followed by days to maturity (96.03%), days to 50% flowering (95.72%), number of clusters per plant (94.14%), seed yield per plant (92.47%), pod length (90.29%), number of pods per plant (89.43%), plant height (88.71%), biological yield per plant (88.25%), number of seeds per pod (88.23%) and number of branches per plant (86.04%), while harvest index (69.56%) recorded medium heritability respectively.

The similar kind of results was also reported by Wesly et al. [5], Parsaniya et al. [7] and Rahangdale et al. [8].

Table 1. ANOVA for various quantitative traits of green gram.

Source of variations	df	Mean sum of square											
		Days to 50% flowering	Days to maturity	Plant height	No. of branches per plant	No. of clusters per plant	No. of pods per plant	Pod length	No. of seeds per pod	100 seed weight	Biological yield per plant	Harvest index	Seed yield per plant
Replication	2	3.475	1.308	6.476	0.017	0.299	1.058	0.059	0.295	0.03	18.109	3.859	2.633
Treatments	39	39.86**	46.53**	67.20**	1.14**	5.65**	46.38**	2.55**	8.17**	2.33**	63.90**	60.27**	7.62**
Error	78	0.586	0.633	2.734	0.059	0.115	1.758	0.088	0.348	0.012	2.715	7.672	0.201

Table 2. Genetic variability parameters for yield and its constituent traits

S. No.	Traits	Range			Coefficient of variation		h ² (bs) %	Genetic Advance as % of mean
		Mean	Min	Max	GCV %	PCV %		
1	Days to 50% flowering	36	26	41	10.00	10.22	95.72	20.16
2	Days to maturity	65	57	73	6.00	6.13	96.03	12.12
3	Plant height	33.20	21.13	44.57	13.96	14.83	88.71	27.09
4	Number of branches per plant	2.02	1.25	3.95	29.78	32.11	86.04	56.91
5	Number of clusters per plant	5.24	2.67	8.93	25.96	26.75	94.14	51.88
6	Number of pods per plant	15.55	6.43	21.93	24.80	26.22	89.43	48.30
7	Pod length	7.82	5.80	9.67	11.58	12.19	90.29	22.67
8	Number of seeds per pod	9.69	4.82	12.80	16.66	17.74	88.23	32.24
9	100 seed weight	3.57	1.84	5.61	24.66	24.85	98.48	50.42
10	Biological yield per plant	17.64	6.08	27.80	25.60	27.25	88.25	49.54
11	Harvest index	30	17.41	38.87	13.96	16.74	69.56	23.98
12	Seed yield per plant	5.28	1.58	8.55	29.80	30.99	92.47	59.03




Fig. 1. Genetic variability parameters of different quantitative traits

For the traits namely, seed yield per plant (59.03%), number of branches per plant (56.91%), number of clusters per plant (51.88%), 100 seed weight (50.42%), biological yield per plant (49.54%), number of pods per plant (48.3%), number of seeds per plant (32.24%), plant height (27.09%), harvest index (23.98%), pod length (22.67%), days to 50% flowering (20.16%), and harvest index (23.98%), there was a high expected genetic advance as a percentage of mean while, days to maturity (12.12%). recorded medium estimates of genetic advance [9].

A transmissibility index is produced for each character based on direct selection criteria, namely broad heritability and genetic advancement as a percentage of mean. This index shows how well the selection process has enhanced each particular character. The majority of the features examined in this study demonstrated high to moderate genetic progress as a percentage of mean together with high estimates of heritability. [10,11] This may be explained by the high selective value and preponderance of additive gene action, which means that selection pressure can be profitably applied to certain features to improve them logically. The similar kinds of results were also reported by Muthuswamy et al. (2022), Parsaniya et al. [7] and Rahangdale et al. [8].

4. CONCLUSION

There is sufficient genetic diversity for each characteristic, as evidenced by the substantial results of the analysis of variance for every trait included in this inquiry. High heritability (bs) was shown to be connected with the highest genetic advancement for most of the characteristics. High yielding breeding traits may be selected based on features that have been determined to have higher heritability and high genetic advance as a percentage of mean in order to optimize the crop's genetic yield potential. In order to choose transgressive segregants in the segregating generations, a methodical hybridization program may also be built, accounting for features with higher heritability and high genetic advancement as a percentage of mean. The segregants will significantly increase the greengram's future yield, productivity, and higher breeding value.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Thakur S, Sharma S, Pachori S, Nagar S, Kumar JA, Tiwari P et al. Understanding genetic variability parameters of greengram (*Vigna radiata* L. Wilczek) Germplasm for agro-morphological traits. International Journal of Plant & Soil Science. 2022;1411-1417. DOI: 10.9734/ijpss/2022/v34i232557
2. Burton GW. Quantitative inheritance in grasses. Proceedings of 6th International Grassland Congress. 1952;1:277-283.
3. Hanson GH, Robinson HF, Comstock RE. Biometrical studies on yield in segregating population. Agron. J. 1956;268-272.
4. Johnson HW, Robinson HF and Comstock HS. Estimation of Genetic and Environmental Variability in Soybean. Agron. J. 1955;47:314-318.
5. Wesley KC, Nagaraju M, Lavanya GR. Estimation of genetic variability and divergence in Greengram *Vigna radiata* (L.) Germplasm Journal of Pharmacognosy and Phytochemistry. 2020;9(2):1890-1893.
6. Sineka T, Murugan E, Sheeba A, Hemalatha G, Vanniarajan C. Genetic relatedness and variability studies in greengram (*Vigna radiata* (L.) Wilczek). Electronic Journal of Plant Breeding. 2021; 12(4):1157-1162.
7. Parsaniya TA, Patel SR, Dinisha A, Mistry HH, Patel HN, Gadhiya CJ. Genetic variability analysis for yield and yield components in mungbean [*Vigna radiata* (L.) Wilczek]. The Pharma Innovation Journal. 2022;11(8):1728-1732.
8. Rahangdale S, Lakhani J, Singh S, Mishra S, Prajapati S, Barela A et al. Study of selection criteria and genetic variability in Mungbean (*Vigna radiata* L. Wilczek). Genotypes for Seed Yield and its Contributing Traits. 2023;41:145-151.
9. Dharajiya DT, Ravindrababu Y, Pagi NK. Screening of Mungbean [*Vigna radiata* (L.) Wilczek] Genotypes for resistance against Mungbean Yellow Mosaic Virus (MYMV) under Field Condition. Int. J. Curr. Microbiol. App. Sci. 2018;7(5):3472-3483.
10. Mathivathana MK, Murukarthick J, Adhimoolam Karthikeyan A, Jang, W, Dhasarathan M, Jagadeeshselvam N et al. Detection of QTLs associated with mungbean yellow mosaic virus (MYMV) resistance using the interspecific cross of

- Vigna radiata* × *Vigna umbellata*. Journal of Applied Genetics; 2019.
11. Prajapati SS, Singh, SK, Shrivastava MK, Singh Y, Kumar P, Rahangdale S, Behera K. Assessment of genetic parameters for yield and its associated traits in greengram [*Vigna radiata* (L.) Wilczek]. International Journal of Environment and Climate Change. 2022; 12(12):840–848.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112965>