



Impact of Weather Parameters on Population Dynamics of Cabbage Aphid (*Brevicoryne brassicae* L.) on Kale (*Brassica oleracea* var. *acephala*)

Mohammad Saleem Bhat ^a, Shafeeq Ahmad Hakeem ^a,
Sameer Ahmad Malik ^b, Ishfaq Majeed Shah ^{a*},
Suhail Qayoom Wani ^b, Shoaib Momin Jan ^c
and Aadil Akbar Wani ^d

^a Division of Entomology, Faculty of Agriculture, Wadura, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 190025, India.

^b Division of Plant Pathology, Faculty of Agriculture, Wadura, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 190025, India.

^c Division of Entomology, Faculty of Horticulture, Shalimar, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 190025, India.

^d Division of Agronomy, Faculty of Agriculture, Wadura, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 190025, India

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i34038

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114051>

Original Research Article

Received: 29/12/2023

Accepted: 01/03/2024

Published: 06/03/2024

ABSTRACT

Studies on "population dynamics of Cabbage Aphid (*Brevicoryne brassicae* L.) on Kale (*Brassica oleracea* var. *acephala*)" in relation to weather parameters revealed cabbage aphid (*Brevicoryne brassicae* L.) as a major pest on Kale. The incidence of cabbage aphid commenced from 16th

*Corresponding author: E-mail: ishfaqskuast@gmail.com;

standard meteorological week (SMW) with a population of 2.02 (aphids/plant). The population of cabbage aphid started increasing and reached its peak in 25th SMW with a population of 137.16 (aphids/plant) after that the population started declining gradually in 26th SMW with a population of 83.14 (aphids/plant). However, the mean population of nymphs during 16th SMW was (2.02 ± 0.46) which gradually started increasing and was highest in 25th SMW (32.68 ± 0.74). The mean population of adults during 17th SMW was (8.24 ± 0.38) and was highest during 25th SMW (62.12 ± 0.50) and the mean population of winged aphids during 18th SMW was (7.10 ± 0.42) which gradually started increasing and attained its peak during 25th SMW (42.36 ± 0.41) respectively. Correlation between maximum and minimum temperature showed positive correlation with cabbage aphid population. Relative humidity of morning showed non-significant negative correlation while relative humidity of evening exhibited positive non-significant correlation. Rainfall were found non-significant positive correlation.

Keywords: Cabbage aphid; *Brevicoryne brassicae*; kale; *Brassica oleracea* var. *acephala*; weather parameters.

1. INTRODUCTION

Vegetables play a vital role in human diets and are essential for feeding the world's population. In the Kashmir Valley, a variety of vegetable crops are cultivated, categorized into groups based on the parts utilized, such as cucurbits, solanaceous fruits, cole crops, root crops, and bulb crops. Among these groups, cruciferous vegetables or cole crops constitute an important category, encompassing a wide variety of leafy vegetables such as cabbage, knol khol, cauliflower, broccoli, kale, mustard, and collard greens [1]. Kale (*Brassica oleracea* var. *acephala*), originating from the eastern Mediterranean region, is one of the oldest members of the cabbage family. It is cultivated across all agro-climatic conditions in the Union Territory of Jammu and Kashmir, ranging from subtropical to temperate to cold arid zones.

Kale (*B.oleracea* var. *acephala*) belongs to the Brassicaceae family, renowned for its green leafy vegetables [2]. Beyond its nutritional value, kale provides notable health benefits. Its leaves offer a plethora of medicinal advantages: it aids in reducing cholesterol levels, thus mitigating the risk of heart disease; it contains various disease-fighting compounds, is abundant in beta-carotene, and serves as an excellent source of ascorbic acid and phyloquinones [3].

Normally kale is cultivated both during *Rabi* and *Kharif* season vegetable and also for seed production [4]. In India, knol-khol is extensively cultivated in regions such as Jammu & Kashmir and West Bengal, and to a lesser extent, it is considered a rare exotic vegetable grown in

select areas of Maharashtra, Assam, Uttar Pradesh, and Punjab [5].

In Jammu and Kashmir, kale enjoys popularity as a vegetable among people from all walks of life, whether rich or poor. It is grown in nearly every kitchen garden and serves as a commercial crop in the vicinity of cities and towns. In Kashmir alone, it is cultivated across an area of 3456.6 hectares, yielding a production of 103,698.6 metric tons [6]. Its demand remains high year-round due to the diverse sizes of coloured knobs and leaves it offers [7]. Some of the popular cultivars of kale grown in Kashmir include Heenz hak, Kawdari hak, Khanyar hak, G. M. Dari hak, Wantipuri hak, Koker hak, Pumb hak and Achari hak [8].

The damage inflicted by the pest complex, ranging from germination to harvest, stands as a primary constraint in the production of cole crops. In India, a total of 37 insect pests have been documented on cole crops [9]. A total of nine species of insect pests on cabbage, cauliflower and kale in Kashmir have been reported, which include cabbage semilooper (*Thysanoplusia orichalcea* Fab.), cabbage butterfly (*Pieris brassicae* L.), diamondback moth (*Plutella xylostella* L.), two species of aphids, mustard aphid (*Lipaphis erysimi* Kalt.) and cabbage aphid -*Brevicoryne brassicae* L. [10]. These insect pests are a threat to global agriculture and cause huge yield and economic losses in *Brassica* crop production. The attractiveness and susceptibility of different cruciferous crops differs for different insect pests and their intensity of damage [11]. The present investigation was initiated by taking into account the crucial role that the kale crop plays in our lives and the prevalence of

one of its major insect pests, the cabbage aphid. This pest significantly impacts both the quantitative and qualitative aspects of kale yield.

2. MATERIALS AND METHODS

To study the population dynamics of kale an experimental trial was laid during *Kharif* 2021 in which the seedlings of kale (*Brassica oleracea* var. *acephala*) cv. G M Dari were transplanted in the 1st week of March 2021 in a plot size of 2x1m with three replications. All the recommended packages of practices were followed to raise a healthy crop. The incidence of cabbage aphid was recorded throughout the growing season. The number of individuals of damaging stages of insect pest in respect of nymphs and adults were counted on five randomly selected plants of each plot on weekly basis throughout the crop season. Population assessment was based on the counts of aphid, nymphs and adults and observation of aphid was recorded from plant measured at weekly interval from initial stage of appearance to final disappearance up to harvest. The nymphal population was counted with the help of a magnifying glass. The data on weather parameters viz., maximum, minimum temperature, relative humidity and rainfall was also collected because this data was correlated against aphid numbers [12].

2.1 Observations Recorded

2.1.1 Population of Nymph (per plant)

The total number of cabbage aphid nymphs were recorded from five randomly selected plants per plot.

2.1.2 Population of Adult (per plant)

The total number of cabbage aphid adults were recorded on five randomly selected plants per plot.

2.1.3 Population of winged aphid (per plant)

The total number of cabbage winged aphids were recorded on five randomly selected plants per plot.

2.1.4 Abiotic factors (temperature, relative humidity, rainfall)

The average aphid population during different periods of the crop growth were examined, in

relation to ambient temperature condition, relative humidity and the influence of rainfall on aphid populations was collected from Division of Agronomy.

The data collected on these characters were subjected to standard statistical analysis.

3. RESULTS

3.1 Incidence of Cabbage aphid (*Brevicoryne brassicae* L.) on kale (*Brassica oleracea* var. *acephala*) during crop season *Kharif* 2021

The results revealed that cabbage aphid was found associated with the kale crop in the experimental field trial during the entire growing season (Table 1). During the study it was observed that cabbage aphid remained closely associated with the crop and its appearance started from 16th standard meteorological week (SMW) with a population of (2.02 aphids/plant) and it gradually started increasing during the growing season and attained peak in the 25th standard week (SMW) with a population of (137.16 aphids/plant). However, the mean population of nymphs during 16th SMW was (2.02 ± 0.46) which gradually started increasing and was highest in 25th SMW (32.68 ± 0.74 nymphs/plant). The mean population of adults during 17th SMW was (8.24 ± 0.38) and was highest during 25th SMW (62.12 ± 0.50) and the mean population of winged aphids during 18th SMW was (7.10 ± 0.42) which gradually started increasing and attained its peak during 25th SMW (42.36 ± 0.41), respectively (Fig. 1).

3.2 Studies on Correlation of Cabbage aphid with Weather Parameters

The population of cabbage aphid showed a great sensibility to weather fluctuations prevailed over the period under study (Fig. no 1). The mean population per plant of cabbage aphid exhibited a highly significant positive correlation with both maximum temperature ($r=0.627^*$) and minimum temperature ($r=0.845^{**}$) while significant negative correlation with morning relative humidity ($r=-0.003^{NS}$) and pest exhibited non-significant correlation with evening relative humidity ($r=0.057^{NS}$) and rainfall ($r=0.486^{NS}$) (Table 2).

Table 1. Incidence of cabbage aphid (*Brevicoryne brassicae*) on kale (*Brassica oleracea* var. *acephala*) during *Kharif* 2021

SMW	Temperature		Relative humidity		Rainfall (mm)	Nymphs/plant	Adults/plant	Winged Aphids/plant	Total population
	Max	Min	RHM	RHE					
14	28	4.5	75	23	10.3	0.00 ± 0.0*	0.00 ± 0.0	0.00 ± 0.0	0.00
15	26.5	5	80	39	9.7	0.00 ± 0.0	0.00 ± 0.0	0.00 ± 0.0	0.00
16	28	6	89	30	8.3	2.02 ± 0.46	0.00 ± 0.0	0.00 ± 0.0	2.02
17	27.5	5	80	19	11.10	6.80 ± 0.58	8.24 ± 0.38	0.00 ± 0.0	15.04
18	29	9	90	24	11.4	14.48 ± 0.75	18.66 ± 0.33	7.10 ± 0.42	40.24
19	27.5	7.5	81	29	11.1	18.40 ± 0.5	22.54 ± 0.34	13.48 ± 0.70	54.42
20	30.5	9	80	34	10.70	21.30 ± 0.71	34.02 ± 0.64	22.40 ± 0.71	77.72
21	28.5	8	90	40	10.8	16.20 ± 0.45	29.20 ± 0.36	23.86 ± 1.05	69.26
22	29.5	9.5	81	41	9.5	16.70 ± 0.87	41.60 ± 0.38	19.9 ± 0.26	78.20
23	31	10	66	22	10.1	24.42 ± 0.77	54.60 ± 0.39	33.38 ± 0.47	112.40
24	31.5	13	82	30	10.2	27.20 ± 2.25	59.08 ± 0.52	37.86 ± 1.26	124.14
25	27	10.5	94	54	10	32.68 ± 0.74	62.12 ± 0.50	42.36 ± 0.41	137.16
26	34.5	12.5	82	41	9.5	17.80 ± 0.73	42.30 ± 0.63	23.04 ± 0.36	83.14

* Mean of three replications, ± Standard Error, SMW = Standard meteorological week

Table 2. Correlation coefficient between cabbage aphid population and Abiotic factors on kale during crop season *Kharif* 2021

Weather parameters	Nymphs	Adults	Winged Aphid
Max. temperature	0.625*	0.657*	0.627*
Min. temperature	0.839**	0.896**	0.845**
RH-M	-0.019 ^{NS}	-0.061 ^{NS}	-0.003 ^{NS}
RH-E	0.119 ^{NS}	-0.004 ^{NS}	0.057 ^{NS}
Rainfall	0.393 ^{NS}	0.426 ^{NS}	0.486 ^{NS}

*Significant at $p \leq 0.05$; Max. Temp = Maximum temperature; Min. Temp = Minimum temperature; R.f = Rainfall; RH= Relative humidity




Fig. 1. Population dynamics of Cabbage aphid (*Brevicoryne brassicae* L.) on Kale (*Brassica oleracea* var. *acephala*) during crop season *Kharif* 2021

4. DISCUSSION

4.1 Incidence of Cabbage Aphid on Kale

During the studies cabbage aphid (*Brevicoryne brassicae* L.) was observed to be the major insect pest on kale crop during *Kharif* 2021. The first appearance of the cabbage aphid population was recorded during 16th SMW 2.02 aphid/plant and thereafter pest population increased and reached at its peak 137.16 aphids/plant during 26th SMW. The population started to decline onwards and was less during 27th SMW with 83.14 aphids/plant. The present findings are in agreement with the studies of Gami et al. [13] and Sinha et al. [14].

4.2 Correlation of Weather Factors with the Population Dynamics of Cabbage Aphid on Kale

The insect pest exhibited a significant positive correlation with both maximum temperature ($r = 0.627$) and minimum temperature ($r = 0.845$) while a non-significant negative correlation with relative humidity morning ($r = -0.003$ ^{NS}) and non-significant positive correlation with relative humidity evening ($r = 0.057$ ^{NS}) and rainfall ($r = 0.486$ ^{NS}). The present findings are in agreement with the studies of Sinha et al. [14], Shonga and Getu [15], Juglan et al. [16], Sahu et al. [17] and Venkateshwarlu et al. [18].

5. CONCLUSION

The incidence of cabbage aphid commenced from 16th SMW with a population of (2.02 aphids/plant) which gradually started increasing. The peak population of cabbage aphid was observed during 25th SMW with a population of (137.16 aphids/plant). The mean population of nymphs during 16th SMW was (2.02 ± 0.46) which gradually started increasing and was highest in 25th SMW (32.68 ± 0.74). The mean population of adults during 17th SMW was (8.24 ± 0.38) and was highest during 25th SMW (62.12 ± 0.50) and the mean population of winged aphids during 18th SMW was (7.10 ± 0.42) which gradually started increasing and attained its peak during 25th SMW (42.36 ± 0.41) respectively. The correlation data revealed that maximum temperature, minimum temperature showed positive correlation with cabbage aphid population while as relative humidity morning showed non-significant negative correlation and relative humidity

evening showed positive non-significant correlation whereas rainfall showed non-significant positive correlation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ahuja I, Rohloff J, Bones AM. Defence mechanisms of Brassicaceae: Implications for plant insect interactions and potential for integrated pest management. *Agronomy for Sustainable Development*. 2010;30(4):311-348.
2. Fahey JW. Brassica. In C. Benjamin (Ed.), *Encyclopedia of food science and nutrition*, 2nd edition Academic Press. 2003;606–615.
3. Tiwari V, Joshi U. Kale: An exotic and underutilized vegetable crop. *Times of Agriculture*. 2020;5(1):30.
4. Anonymous. Annual Report of Seed production. Department of Agriculture Kashmir, J&K Government, Srinagar: 2010;76.
5. Thamburaj S, Singh N. Vegetables, Tuber crops and Spices, Indian Council of Agricultural Research New Delhi: 2016; 141-142.
6. Anonymous.2022. Area and production of crops in Kashmir. Annual report, Directorate of Agriculture, Kashmir.
7. Anonymous. Area and production of crop in J&K. Annual report, Department of Agriculture, Kashmir: 2019;75.
8. Khan SH, Ahmed N, Jaben N, Chato MA, Hussain K. Biodiversity of kale (*Brassica oleracea* var. *acephala* L.) in Kashmir valley. *Asian Journal of Horticulture*. 2010; 5(1):208-210.
9. Lal OP. A compendium of insect pests of vegetables in India. *Bulletin of Entomology*. 1975;16(2):31-56.
10. Malik RA, Punjabi AA, Bhat AA. Survey and study of insect and non-insect pest. *Horticulturist*. 1972;3(3):29-44.
11. Hiiesaar K, Metspalu L, Joger K. Attractiveness and susceptibility of *Brassica rapa*, *Brassica napus* and *Sinapis alba* to the flea beetles (Coleoptera: Chrysomelidae). *Agronomy Research*. 2006;4(2):191-196.
12. Choudhury S, Pal S. Population dynamics of mustard aphid on different *Brassica*

- cultivars under terai agro-ecological conditions of West Bengal. The Journal of Plant Protection Sciences. 2009;1(1):13-16.
13. Gami JM, Bapodra JG, Rathod R. Population dynamics of mustard aphid (*Lipaphis erysimi* Kalt.) in relation to weather parameters. Indian Journal of Plant Protection. 2002;30(2): 202-204.
 14. Sinha R, Yazdani SS, Verma GD. Population dynamics of mustard aphid, *Lipaphis erysimi* (Kalt.) in relation to ecological parameters. Indian Journal of Entomology. 1989;51(6):334-339.
 15. Shonga E, Getu E. Population dynamics of cabbage aphid, *Brevicoryne brassicae* L. (Homoptera: Aphididae) in relation to weather factors on major brassica crops in central rift valley of Ethiopia: Baseline studies for the management of the pest. International Journal of Tropical Insect Science. 2020;41(1&2):455-462.
 16. Juglan RS, Singh R, Singh H. Effect of abiotic factors on the field population of Mustard aphid, *Lipaphis erysimi* (Kalt.) Indian Journal of Ecosystem. 1988;15(2): 163-167.
 17. Sahu B, Pachori R, Nagasr N, Patidar S. Population dynamics of Cabbage aphid, *Lipaphis erysimi* on cabbage crop (*Brassica oleracea* var. *capitata* L.). Journal of Pharmacognosy and Phytochemistry. 2020;9(3):950-952.
 18. Venkateswarlu V, Sharma RK, Chander S, Singh SD. Population dynamics of major insect pests and their natural enemies in cabbage. Annals of Plant Protection Sciences. 2011;19(2): 272-277.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/114051>