



# Complementary Food Formulated from Orange Flesh Sweet Potato, Soybean and Date Palm Flour Blends Improve on the Functional, Vitamins and Mineral Properties Children Formulas

Peter Abuengmoh <sup>a\*</sup>, Dick Iorwuese Gernah <sup>a,b</sup>  
and Dinnah Ahure <sup>a,c</sup>

<sup>a</sup> Department of Chemistry, Faculty of Science, Benue State University, Center for Food Technology and Research, Makurdi, Nigeria.

<sup>b</sup> Department of Food Science and Technology, Federal University of Agriculture, Makurdi, Benue State, Nigeria.

<sup>c</sup> Department of Food Science and Technology, University of Mkar, Benue State, Nigeria.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118742>

**Original Research Article**

**Received: 27/04/2024**

**Accepted: 01/07/2024**

**Published: 04/07/2024**

## **ABSTRACT**

This study evaluated the functional properties, vitamins and mineral composition of complementary foods produced from orange flesh sweet potato (OFSP), soybean and date palm flour blends. The samples were formulated by material balancing with all samples having 16% protein/ 100g. The

\*Corresponding author: E-mail: [abuengmohpeter@gmail.com](mailto:abuengmohpeter@gmail.com);

**Cite as:** Abuengmoh, P., Gernah, D. I., & Ahure, D. (2024). Complementary Food Formulated from Orange Flesh Sweet Potato, Soybean and Date Palm Flour Blends Improve on the Functional, Vitamins and Mineral Properties Children Formulas. *Asian Journal of Food Research and Nutrition*, 3(3), 609–620. Retrieved from <https://journalajfrn.com/index.php/AJFRN/article/view/159>

samples were ascribed the following codes; CON, CFA, CFB, CFC, CFD and CFE. Sample CON being a commercial product used for control while the others were experimental formulations. The samples were subjected to functional, vitamin and mineral analysis using standard method. Results of the functional properties of flour blends were as follows; bulk density (0.66-0.91 g/ml), water absorption capacity (1.82-2.63 g/ml), oil absorption capacity (1.39-1.72 g/ml), gelatinization temperature (69.98-80.04°C), foam capacity (29.61-64.31%) swelling capacity (2.75-3.60 g/ml) and there were significant differences between some of the flour blends at ( $p < 0.05$ ). Vitamin analysis obtained the following range of values; Vitamin A (110.32-685.63 µg/100g), Vitamin B1 (0.42-0.85 mg/100g), Vitamin B2 (0.41-0.49 mg/100g), Vitamin B3 (2.55-4.21 mg/100g), Vitamin B6 (0.30-1.03 mg/100g), Vitamin B9 (27.32-193.85 mg/100g), Vitamin C (11.03-65.15 mg/100g). The experimental samples were very rich in vitamin A which is an essential nutrient that helps to build up the immune system of infants against a number of infections and sustains the integrity of the epithelial linings. Mineral composition results were obtained as follows; Calcium (261.84-351.32 mg/100g), Potassium (632.85-718.23 mg/100g), Sodium (35.85-190.11 mg/100g), Phosphorus (292.87-546.72 mg/100g), Iron (10.73-19.91 mg/100g), Zinc (5.33-25.34 mg/100g) and Iodine (47.32-90.19 mg/100g). These minerals play crucial roles in osmotic regulation, various physiological functions in body processes and are necessary for maintaining overall health and well-being.

**Keywords:** OFSP; soybean; dates; complementary foods; potassium; complementary food.

## 1. INTRODUCTION

"Complementary food (CF) is any healthful and energy containing solid, semisolid or liquid meal consumed by infants besides human milk or formula" [1]. "They are essentially introduced from the period of 4-6 months when breast milk, considered being the choicest and safest meal for young babies, can no longer provide the nutrients and energy requirements needed to enable the child to grow and thrive" [2,3]. According to WHO 1998, "it is any food other than breast milk given to a breastfeeding child and the process is called complementary feeding practice". "Complementary food is mostly prepared in the form of thin porridges or gruels" [4]. "When breast milk alone is no longer sufficient, complementary feeding can meet the additional nutritional requirements of the infants. Complementary feeding begins at 6 months when transition from exclusive breastfeeding to semi-solid foods begins and continues up to the age of 24 months" [5,6] where "most infant need additional foods, to complement the breast milk and make certain nutrient to provide enough energy and nutrients for normal growth. This goal is only achieved when these foods are prepared and fed to the infants under hygienic conditions and given in adequate proportions. The main concern is making sure that there is no gap between nutrient requirements and what a child is able to consume, absorb and utilize. Most infants suffer from malnutrition not because of the economic status of the country but because of inability to utilize the available raw materials to meet their daily requirements".

"Sweet potato (*Ipomoea batatas L.*) is one of the world's most important, versatile and underexploited food crops" [7,8]. "The sweet potato tuber grows underground, ranges in flesh colour from white, cream to yellow, purple or orange, and is high in soluble carbohydrates, minerals and vitamins. Sweet potato roots have high nutritional value, energy content and other micronutrients such as vitamin A, C, potassium, iron and zinc. Notwithstanding, it is low in protein and fat contents; hence, the need to complement it with legumes which will boost the protein contents of the complementary foods, thus the use of soy bean. Soybean (*Glycine max L.*) products have been proven to be good substitutes for animal products as they offer a "complete" protein profile, containing almost all the essential amino acids (except methionine) and a range of water-soluble and fat soluble vitamins. Soybean flour is among the products increasingly consumed in sub Saharan Africa and has been reported to be good a source of protein, lipid and other essential nutrients" [9]. In as much as we want the food product to be rich nutritionally and functionally, it also has to be palatable for the concerned to enjoy while consuming. Thus the use of date fruit as a sweetener and nutrient enhancer is very important.

"Date palm fruit (*Phoenix dactylifera L.*) is a good source of sugar (around 70–80 % sugar content), depending on the species and maturity stage of the fruits likes glucose, fructose, and sucrose which are easily absorbed by the body to provide energy" [10]. "Date fruit contains a considerable

amount of moisture (12–14), crude fibre (4.5–5.0), protein (2.0–2.5), fat (1.8–2.4), and ash (1.7–1.9) which is essential for children's growth and developments" [10]. "The vitamin content in date fruit is low but interestingly, the content of B-complex vitamins is high in date fruits. Date contains a high amount of minerals (selenium, magnesium, potassium, calcium and phosphorus) which is essential for the development of strong bones and teeth" [10]. "Dates is also rich in antioxidants and phenolic compounds and also contains antimicrobial properties" [10]. This will help to improve on the storage stability of the complimentary food formulations.

### 1.1 Statement of the Problem

Most infants suffer from protein energy malnutrition not because of the economic status of the country but because of inability to utilize the available raw materials to meet their daily requirements. Locally accessible ingredients such as sweet potatoes, soybean and date palm fruit are still highly underutilized in formulating complementary foods despite the fact that they are highly available, nutritionally rich, low in anti-nutritional with good functional properties, and forms a low viscous porridge.

## 2. METHODOLOGY

The research used a combination of quantitative approach to determine the vitamins and mineral composition (AOAC 2015) and a qualitative approach to evaluate the functional properties of food formulations.

**Procurement of materials:** Sweet potato was obtained from the National Roots and Crops Research Institute (NRCRI), UMUDIKE. Soybeans and dates palm fruits were obtained from the Wurukum Market, Makurdi. Nestle Cerelac was obtained from a local supermarket in Makurdi. Procured materials were taken to the CEFTER Food Processing Laboratory, Benue state university for processing.

**Equipment:** Standard equipment from the CEFTER food processing laboratory was used for the processing of the products. This equipment included but not limited to knives, bows, trays, driers, blender, mixer, oven, sieves (0.5 mm), measuring cylinder and weighing scale. Standard analytical equipment from the Chemistry laboratory were used for lab analysis.

**Preparation of raw materials:** Standard processing procedures and chemicals were used in the preparation of the flours and complementary food formulations.

**Sweet potato flour:** Mature orange-flesh sweet potato tubers (*Ipomoea batatas*), was cleaned and trimmed manually using knives. They were washed, peeled and sliced using a manual food slicer in to smaller sizes of approximately 10 mm thickness to facilitate the drying process. The sliced sweet potatoes were blanched for 5mins at 70 °C and dried in air circulating oven (Gallenkamp S/No 90/02/190, UK) at 60 °C for 48 h according to the method of Marcel et al. [11] with slight modification. The dried samples were milled to pass a 0.5 mm sieve for flour blends [12].

**Soy bean flour:** The soy bean seeds were sorted, washed and soaked in water for 24 hours. The soaked beans were placed in a sieve and the water drained off. The beans were then boiled for 30 minutes to inactivate enzyme activity and also to make decortication easy. The soybeans were dehulled by robbing between palms and the peels drain off using potable water. It was then oven dried at 60 °C for 48 h in a hot-air oven, after which they were roasted at 120 °C for 30 mins. The roasted beans were milled with a laboratory mill (Laboratory Blender of model KM 901D; Kenwood Electronic, Hertfordshire, UK) into flour. The flour were sieved through a 0.5 mm mesh sieve [7].

**Date palm flour:** The date flour was prepared by sorting the date palm nuts to remove all foreign particles and washing. It was followed by de-seeding through the manually cutting the nut using knife and removing the seed. The inside was cleaned and size reduction of the nut for efficient drying and milling processes. The pulp with the pericarp was then oven dried at 60 °C for 72 h [13]. The dried pulp was milled into fine flour using laboratory blender. The flour was then sieved through a 0.5 mm mesh sieve (British Standard) to obtain fine homogenous particle size flour and packaged.

**Food product and diet formulation:** Flours from orange-fleshed sweet potatoes, soy beans, and Date palm fruits were blended in different proportions as presented in Table 1, to develop varieties of complementary flour of composite mixtures obtained from different levels/percentages of flour substitution. These ratios

were arrived at, based on their protein content through material balancing to give 16 g protein/100g food as recommended by WFP, [14] and UNICEF 2016 for infant diets. Therefore, from the treatment combinations, five samples (CFA, CFB, CFC, CFD & CFE) were generated with the aid of the material balancing and Nestle Cerelac as standards.

**Analysis:**

**Functional properties of complementary food formulations:** The functional properties (Bulk density, Water Absorption Capacities (WAC), Oil Absorption Capacities (OAC), Gelatinization temperature, Foaming capacity, Swelling Capacity of complementary food formulations were determined by the method described in Okoronkwo et al. [1].

**Determination of vitamins content of complementary food formulations:** Vitamin A was determined by the calorimetric method described by AOAC 2012. Vitamins B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub>, B<sub>6</sub>, B<sub>9</sub> and Vitamin C were determined through the spectrophotometric method, described by Abuengmoh et al., [15]. Exactly 10 g of the sample was extracted with 50 ml EDTA/TCA (50 g in 50 ml of water) extracting Solution for 1 hour and filtered through a What man filter paper into a 50ml volumetric flask and made up to the mark with the extracting solution. Twenty (20 ml) of the extract was pipetted into a 250 ml conical flask and 10 ml of 30% KI was added and also 50 ml of distilled water added. This was followed by 2 ml of 1% starch indicator. This was titrated against 0.01 ml CuSO<sub>4</sub> solution to a dark end point.

**Determination of mineral elements of the flours and their complementary blends formulations:**

“Calcium (Ca), Zinc (Zn), Iron (Fe) was determined using Atomic Absorption Spectrophotometer (AAS) as described by AOAC, (2012). One gram (1 g) of the sample was first digested with 30 ml of aqua regia, which was a mixture of concentrated HNO<sub>3</sub> and HCl, in a ratio of 1:3. The digested sample was filtered and increased up to 50 ml with deionized water. The aliquots of the digested filtrate were used for AAS using filters that match the different elements” (AOAC, 2012). “The sodium (Na) content was determined using flame emission photometer (Sherwood Flame Photometer 410, Sherwood Scientific Ltd. Cambridge, UK) with NaCl as the standards” (AOAC 2012). “The potassium (K) content was determined using flame emission photometer (Sherwood Flame Photometer 410, Sherwood Scientific Ltd. Cambridge, UK) with KCl as the standards” (AOAC 2012). Phosphorous concentration in the sample was measured colourimetrically using the molybdovanadate method used by Abuengmoh et al. [15].

**Statistical analysis:** Experiments were conducted in triplicates and analysis of variance (one way ANOVA) were performed and results were separated using the Multiple Ranges Duncan’s test (p < 0.05) on a statistical software of Statistical Package for the Social Sciences (SPSS), Version 28. The mean values and standard deviations were calculated where appropriate, reported with differences considered at 95% significant level [16].

**Table 1. Sample formulation of complimentary food**

Samples	Orange-fleshed Sweet potato (%)	Soybean Flour (%)		Date (%)
		Boiled/Oven-dried	Sun-dried/Toasted	
CON	/	/	/	/
CFA	57.5	42.5	/	/
CFB	48	42	/	10
CFC	48	21	21	10
CFD	58	/	42	/
CFE	47.5	/	41.5	10

**KEY:**

CON: Nestle Cerelac

CFA: Boiled and Oven-dried Soy Flour/Orange Fleshed Sweet Potato

CFB: Date palm, Boiled and Oven-dried Soy Flour/Orange Fleshed Sweet Potato

CFC: Date palm, Boiled and Oven-dried/Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

CFD: Non-Sweetened Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

CFE: Sweetened Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

### 3. RESULTS AND DISCUSSION

**Functional properties of complementary food formulations:** The functional properties of flours analysed were as shown in Table 2. The functional properties determined were bulk density, water absorption capacity, oil absorption capacity, gelation temperature, foam capacity, and swelling capacity.

**Bulk density of complementary food formulations:** The bulk density ranged from 0.66 (g/ml) in sample to 0.91 (g/ml) in sample CON. It was observed that the bulk density increased significantly ( $p < 0.05$ ) based on the treatment method of the soy bean flour. The samples with sun-dried and toasted soy flour showed the least bulk density compared to the samples with boiled and oven-dried soy flour. The sun-dried and toasted soy bean was well dried with the least moisture content thus making the seeds to be milled into smaller particle size than the boiled and oven-dried soy bean grain and hence the reduction in bulk density. This result was similar to the findings of Bello et al., [17] where the values of bulk density in malted food formulations were significantly ( $p < 0.05$ ) lower than the values in the non-malted food formulations. The significance of this is that the less bulky flour has higher nutrient density, since more flour can be packaged in the same given volume. There was significant ( $p < 0.05$ ) increase in bulk density of Nestle Cerelac food formulation.

**Water absorption capacity (WAC) of complementary food formulations:** The water absorption capacity (WAC) ranged from 1.82 (g/ml) in sample to 2.63 (g/ml) in sample CON. It was observed that the water absorption capacity increased significantly ( $p < 0.05$ ) based on the treatment method of the soy bean flour. The samples with boiled and oven-dried soy flour showed the least water absorption capacity compared to the samples with sun-dried and toasted soy bean flour. The sun-dried and toasted soy bean was well dried with the least moisture content than the later thus making the samples to absorb more water than the samples with boiled and oven-dried soy bean flour and hence the increase in water absorption capacity. This result was similar to the findings of Bello et al., [17] where the values of water absorption capacity in malted food formulations were significantly ( $p < 0.05$ ) higher than the values in the non-malted food formulations. There was significant ( $p < 0.05$ ) increase in water absorption capacity of Nestle Cerelac food formulation. Water absorption capacity or characteristics

represent the ability of a product to associate with water under conditions where water is limited.

**Oil absorption capacity (OAC) of complementary food formulations:** The oil absorption capacity (OAC) ranged from 1.39 (g/ml) in sample CON to 1.72 (g/ml) in sample CFE. There was a significant difference ( $p < 0.05$ ) between samples CON every other sample. It was also observed that the oil absorption capacity increased significantly ( $p < 0.05$ ) based on the treatment method of the soy bean flour. The samples with boiled and oven-dried soy flour showed moderate oil absorption capacity compared to the samples with sun-dried and toasted soy bean flour that recorded the highest value of oil absorption capacity. This result was similar to the findings of Bello et al., [17] where the values of oil absorption capacity in malted food formulations were significantly ( $p < 0.05$ ) higher than the values in the non-malted food formulations. The water and oil binding capacity of food protein depend upon the intrinsic factors like amino acid composition, protein conformation and surface polarity or hydrophobicity. The ability of the proteins of these flours to bind with oil makes it useful in food system where optimum oil absorption is desired. This makes flour to have potential functional uses in foods such as sausage production. The OAC also makes the flour suitable in facilitating enhancement in flavour and mouth feel when used in food preparation.

**Gelatinization temperature (GT) of complementary food formulations:** The gelatinization temperature (GT) of the blended food formulations ranged from 69.98 °C in sample CON to 80.04 °C in sample CFE. The samples with boiled and oven-dried soy flour were significantly ( $p < 0.05$ ) different from the samples sun-dried and toasted soy flour. This implies that the processing method of soy bean flour greatly influences the. It was also observed that the gelatinization temperature increased significantly ( $p < 0.05$ ) based on the processing method of the soybean flour. The samples with boiled and oven-dried soy flour showed lower gelatinization temperature compared to the samples with sun-dried and toasted soy bean flour that recorded the highest value of gelatinization temperature. This result was similar to the findings of Bello et al., [17] where the values of oil absorption capacity in malted food formulations were significantly ( $p < 0.05$ ) higher than the values in the non-malted food formulations.

**Table 2. Functional Properties of complementary food formulations**

Samples	Bulk Density (g/ml)	WAC (g/ml)	OAC (g/ml)	GT (°C)	FC (%)	SC (g/ml)
CON	0.91 <sup>a</sup> ±0.01	2.63 <sup>a</sup> ±0.01	1.39 <sup>c</sup> ±0.01	69.98 <sup>c</sup> ±0.01	64.31 <sup>a</sup> ±0.01	2.75 <sup>c</sup> ±0.01
CFA	0.81 <sup>b</sup> ±0.01	1.82 <sup>c</sup> ±0.01	1.43 <sup>b</sup> ±0.01	71.81 <sup>b</sup> ±0.02	29.61 <sup>e</sup> ±0.02	3.45 <sup>ab</sup> ±0.01
CFB	0.73 <sup>bc</sup> ±0.01	1.92 <sup>c</sup> ±0.01	1.50 <sup>b</sup> ±0.01	73.13 <sup>b</sup> ±0.02	35.17 <sup>d</sup> ±0.02	3.18 <sup>b</sup> ±0.02
CFC	0.69 <sup>c</sup> ±0.01	2.06 <sup>b</sup> ±0.01	1.57 <sup>ab</sup> ±0.02	76.36 <sup>ab</sup> ±0.02	40.61 <sup>c</sup> ±0.02	3.22 <sup>b</sup> ±0.02
CFD	0.67 <sup>c</sup> ±0.01	2.18 <sup>b</sup> ±0.01	1.63 <sup>a</sup> ±0.02	78.14 <sup>a</sup> ±0.02	46.36 <sup>bc</sup> ±0.01	3.60 <sup>a</sup> ±0.01
CFE	0.66 <sup>c</sup> ±0.01	2.31 <sup>ab</sup> ±0.01	1.72 <sup>a</sup> ±0.02	80.04 <sup>a</sup> ±0.02	52.41 <sup>b</sup> ±0.01	3.43 <sup>ab</sup> ±0.01

Values are means ± standard deviation of triplicate determinations.

Mean scores within column with same letters are not significantly different ( $p < 0.05$ ).

Key

CON: Nestle Cerelac

CFA: Non-Sweeten Boiled and Oven-dried Soy Flour/Orange Fleshed Sweet Potato

CFB: Sweeten-Boiled and Oven-dried Soy Flour/Orange Fleshed Sweet Potato

CFC: Sweetened-Boiled and Oven-dried/Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

CFD: Non-Sweetened Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

CFE: Sweetened Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

**Foam capacity of complementary food formulations:** The foam capacity ranged from 29.61 % in sample CFA to 64.31 % in sample CON. It was observed that the foam capacity increased significantly ( $p < 0.05$ ) based on the treatment method of the soy bean flour. The lowest foam capacity was observed for sample CFA (29.61%) and the samples with boiled and oven-dried soy flour were significantly ( $p < 0.05$ ) different from the samples sun-dried and toasted soy flour. The samples sun-dried and toasted soy flour recorded higher values of foam capacity. The highest value was observed for sample CON (64.31%) and it was also significantly different from other samples. Protein in the dispersion may cause a lowering of the surface tension at the water air interface, thus always been due to protein which forms a continuous cohesive film around the air bubbles in the foam.

**Swelling capacity of complementary food formulations:** The swelling capacity ranged from 2.75 g/ml in sample CON to 3.60 g/ml in sample CFD. It was observed that the swelling capacity increased significantly ( $p < 0.05$ ) in samples with higher ratios of orange-fleshed sweet potato. The lowest swelling capacity was observed for sample CON (2.75 g/ml). The samples with soy bean flour and orange-fleshed sweet potato flour only were significantly ( $p < 0.05$ ) different/higher than the samples that had date palm pulp flour. The swelling capacity of flours depends on size of particles, types of variety and types of processing methods or unit operations. As per results, the flour of orange-fleshed sweet potato has more swelling capacity as compared to date palm pulp flour. This result agrees with those of Bello et al., [17] who worked on food formulations from malted, non-malted maize and *Moringa oleifera*. They observed that swelling capacity

increase with addition of *Moringa oleifera* which could be as a result of *Moringa oleifera* containing more fibre than sorghum and soybean. The malted flour whose starch was already dextrinised could not swell as much as the non-malted food formulations. The higher swelling index of the non-malted samples and also with addition of *Moringa oleifera* seed flour was as a result of starch and fibre.

**Vitamin Composition of complementary food formulations:** The study evaluated the vitamin composition of complementary food formulations as shown in Table 3. The vitamins that were determined were vitamin A, vitamin B<sub>1</sub>, vitamin B<sub>2</sub>, vitamin B<sub>3</sub>, vitamin B<sub>6</sub>, vitamin B<sub>9</sub>, vitamin C.

**Vitamin A content in the complementary food formulations:** The vitamin A content of the complementary food blend ranged from 110.32 (µg/100 g) in sample CON to 685.63 (µg/100 g) in Sample CFA. It was observed that the vitamin A content increased significantly ( $p < 0.05$ ) in samples with higher ratios of orange-fleshed sweet potato. The lowest vitamin A content was observed for sample CON 110.32 µg/100 g. The samples with soy bean flour and orange-fleshed sweet potato flour only were significantly ( $p < 0.05$ ) different/higher than the samples that had date palm pulp flour. The vitamin A content of samples depends on the quantity of the orange-fleshed sweet potato flour in the sample since orange-fleshed sweet potato is rich in β-carotenes which is pro-vitamin A. As per results, the flour of orange-fleshed sweet potato has more vitamin A as compared to date palm pulp flour. This result agrees with those of Okoronkwo et al., [1] who worked on complementary food formulations from Solid-State Fermentation of Fonio, Soybean and Orange-Fleshed Sweet Potato Blends. One of the easiest ways to

introduce more vitamin A into an infant's diet is by the addition of carotene-rich plant-based foods. Vitamin A helps in the prevention of xerophthalmia and keratomalacia of the eyes [18]. Vitamin A is an essential nutrient that helps to build up the immune system of infants against a number of infections and sustains the integrity of the epithelial linings [1].

**Vitamin B<sub>1</sub> (Thiamine) content in the complementary food formulations:** The vitamin B<sub>1</sub> (Thiamine) content of the complementary food blend ranged from 0.42 mg/100g in sample CON to 0.85 mg/100g in sample. It was observed that the vitamin B<sub>1</sub> content increased significantly ( $p < 0.05$ ) in samples with higher ratios of orange-fleshed sweet potato. The lowest vitamin B<sub>1</sub> content was observed for sample CON 0.42 mg/100g and it was significantly ( $p < 0.05$ ) different from all the other formulations. The samples with soy bean flour and orange-fleshed sweet potato flour only were significantly ( $p < 0.05$ ) different/higher than the samples that had date palm pulp flour. Thiamine helps in the treatment of beriberi and in the maintenance of healthy attitude in humans [18]. The vitamin B<sub>1</sub> of the complementary food formulations depends largely on the quantity of the orange-fleshed sweet potato flour in the sample since orange-fleshed sweet potato is a vegetable and is very rich in vitamins. There were significant ( $p < 0.05$ ) differences between the complementary food formulations not containing date palm pulp flour and those containing date palm pulp flour [1]. Vitamins are very important for fighting against diseases in the body.

**Vitamin B<sub>2</sub> (Riboflavin) content in the complementary food formulations:** The vitamin B<sub>2</sub> (Riboflavin) content of the complementary food blend ranged from 0.41 mg/100g in sample CON to 0.49 mg/100g in sample CFA. It was observed that the vitamin B<sub>2</sub> content increased significantly ( $p < 0.05$ ) in samples with higher ratios of orange-fleshed sweet potato. The lowest vitamin B<sub>2</sub> content was observed for sample CON 0.41 mg/100g and it was significantly ( $p < 0.05$ ) different from all the other formulations. Riboflavin is essential for growth and development in infants and young children [1]. The samples with soy bean flour and orange-fleshed sweet potato flour only were significantly ( $p < 0.05$ ) different/higher than the samples that had date palm pulp flour. The vitamin B<sub>2</sub> of the complementary food formulations depends largely on the quantity of the orange-fleshed sweet potato flour in the

sample since orange-fleshed sweet potato is a vegetable and is very rich in vitamins. Vitamins are very important for fighting against diseases in the body. There were significant ( $p < 0.05$ ) differences between the complementary food formulations not containing date palm pulp flour and those containing date palm pulp flour [1].

**Vitamin B<sub>3</sub> (Niacin) content in the complementary food formulations:** The vitamin B<sub>3</sub> (Niacin) content of the complementary food blend ranged from 2.55 mg/100g in sample CFC to 4.21 mg/100g in sample CON. It was observed that the vitamin B<sub>3</sub> content increased insignificantly ( $p < 0.05$ ) in samples with higher ratios of orange-fleshed sweet potato. The lowest vitamin B<sub>3</sub> content was observed for sample CFC comprising of dates, boiled and oven-dried/sun-dried/toasted soy flour and orange-fleshed sweet potato (2.55 mg/100g) and it was not significantly ( $p < 0.05$ ) different from all the other formulations except sample CON. Niacin is a component of the respiratory co-enzyme (NAD) that is responsible for tissue oxidation in the body. This result agreed with those observed by Okoye et al., [18]. The vitamin B<sub>3</sub> content of the complementary food formulations depends slightly on the quantity of the orange-fleshed sweet potato flour in the sample since orange-fleshed sweet potato is a vegetable and is very rich in vitamins. There was a significant ( $p < 0.05$ ) differences between sample CON and all other complementary food formulation.

**Vitamin B<sub>6</sub> (Pyridoxine) content in the complementary food formulations:** The vitamin B<sub>6</sub> (Pyridoxine) content of the complementary food blend ranged from 0.30 mg/100g in sample CON to 1.03 mg/100g in sample CFA. It was observed that the vitamin B<sub>6</sub> content increased significantly ( $p < 0.05$ ) in samples with sun-dried and toasted soy flour. The lowest vitamin B<sub>6</sub> content was observed for sample CON 0.30 mg/100g and it was significantly ( $p < 0.05$ ) different from all the other formulations. The samples with sun-dried and toasted soy flour were significantly ( $p < 0.05$ ) different/higher than the samples that were made of boiled and oven-dried soy flour. The vitamin B<sub>6</sub> content of the complementary food formulations depends slightly on the quantity of soy bean flour and its processing method since boiling will cause some of the vitamins to leach out. There were significant ( $p < 0.05$ ) differences between the complementary food formulations made of boiled and oven-dried soy flour and sun-dried and toasted soy flour [1].

**Table 3. Vitamin content in the complementary food formulations**

Samples	Vit. A (µg /100g)	Vit. B <sub>1</sub> (mg/100g)	Vit. B <sub>2</sub> (mg/100g)	Vit. B <sub>3</sub> (mg/100g)	Vit. B <sub>6</sub> (mg/100g)	Vit. B <sub>9</sub> (µg/100g)	Vit. C (mg/100g)
CON	110.32 <sup>c</sup> ±0.03	0.42 <sup>c</sup> ±0.21	0.41 <sup>c</sup> ±0.01	4.21 <sup>a</sup> ±0.01	0.30 <sup>b</sup> ±0.01	27.32 <sup>c</sup> ±0.01	65.15 <sup>a</sup> ±0.01
CFA	685.63 <sup>a</sup> ±0.04	0.83 <sup>a</sup> ±0.01	0.47 <sup>a</sup> ±0.02	2.96 <sup>b</sup> ±0.02	0.63 <sup>b</sup> ±0.01	175.86 <sup>ab</sup> ±0.11	11.03 <sup>d</sup> ±0.01
CFB	584.11 <sup>b</sup> ±0.35	0.61 <sup>b</sup> ±0.01	0.44 <sup>b</sup> ±0.00	2.73 <sup>b</sup> ±0.02	0.63 <sup>b</sup> ±0.02	147.63 <sup>b</sup> ±0.06	16.93 <sup>b</sup> ±0.02
CFC	585.63 <sup>b</sup> ±0.04	0.71 <sup>ab</sup> ±0.01	0.45 <sup>b</sup> ±0.01	2.55 <sup>b</sup> ±0.02	0.74 <sup>ab</sup> ±0.02	165.78 <sup>ab</sup> ±0.18	16.74 <sup>b</sup> ±0.02
CFD	679.28 <sup>a</sup> ±0.09	0.85 <sup>a</sup> ±0.01	0.49 <sup>a</sup> ±0.01	2.86 <sup>b</sup> ±0.02	1.03 <sup>a</sup> ±0.01	185.83 <sup>a</sup> ±0.11	15.58 <sup>c</sup> ±0.02
CFE	588.24 <sup>b</sup> ±0.08	0.71 <sup>ab</sup> ±0.01	0.46 <sup>ab</sup> ±0.01	2.66 <sup>b</sup> ±0.02	0.97 <sup>a</sup> ±0.02	193.85 <sup>a</sup> ±0.04	17.42 <sup>b</sup> ±0.01

Values are means ± standard deviation of triplicate determinations. Mean scores within the column with same letters are not significantly different (p<0.05).

Key

CON: Nestle Cerelac

CFA: Non-Sweeten Boiled and Oven-dried Soy Flour/Orange Fleshed Sweet Potato

CFB: Sweeten-Boiled and Oven-dried Soy Flour/Orange Fleshed Sweet Potato

CFC: Sweetened-Boiled and Oven-dried/Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

CFD: Non-Sweetened Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

CFE: Sweetened Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

**Table 4. Mineral composition of complementary food formulations**

Samples	Ca (mg/100g)	K (mg/100g)	Na(mg/100g)	P (mg/100g)	Fe (mg/100g)	Zn (mg/100g)	I <sub>2</sub> (µg/100g)
CON	351.32 <sup>a</sup> ±0.03	632.85 <sup>c</sup> ±0.01	190.11 <sup>a</sup> ±0.01	292.87 <sup>b</sup> ±0.21	10.73 <sup>c</sup> ±0.01	25.34 <sup>a</sup> ± 0.01	47.32 <sup>b</sup> ± 0.01
CFA	261.84 <sup>c</sup> ±0.09	696.68 <sup>b</sup> ±0.06	35.85 <sup>b</sup> ± 0.04	539.14 <sup>a</sup> ±0.03	16.52 <sup>b</sup> ±0.02	5.33 <sup>c</sup> ± 0.02	90.07 <sup>a</sup> ± 0.12
CFB	264.24 <sup>b</sup> ±0.11	711.23 <sup>a</sup> ±0.06	37.35 <sup>b</sup> ± 0.02	544.73 <sup>a</sup> ±1.02	18.12 <sup>ab</sup> ±0.02	6.86 <sup>bc</sup> ± 0.02	90.10 <sup>a</sup> ± 0.01
CFC	264.82 <sup>b</sup> ±0.08	715.37 <sup>a</sup> ±0.06	38.66 <sup>b</sup> ± 0.18	545.47 <sup>a</sup> ±0.58	18.72 <sup>ab</sup> ±0.02	6.42 <sup>bc</sup> ± 0.02	90.17 <sup>a</sup> ± 0.01
CFD	262.45 <sup>c</sup> ±0.11	706.68 <sup>ab</sup> ±0.06	37.85 <sup>b</sup> ± 0.13	542.92 <sup>a</sup> ±0.05	17.82 <sup>b</sup> ±0.02	6.67 <sup>bc</sup> ± 0.02	90.15 <sup>a</sup> ± 0.11
CFE	265.35 <sup>b</sup> ±0.05	718.23 <sup>a</sup> ±0.06	39.72 <sup>b</sup> ± 0.02	546.72 <sup>a</sup> ±0.15	19.91 <sup>a</sup> ±0.02	7.43 <sup>b</sup> ± 0.02	90.19 <sup>a</sup> ± 0.01

Values are means ± standard deviation of triplicate determinations.

Mean scores within the column with same letters are not significantly different (p>0.05).

Key

CON: Nestle Cerelac

CFA: Non-Sweeten Boiled and Oven-dried Soy Flour/Orange Fleshed Sweet Potato

CFB: Sweeten-Boiled and Oven-dried Soy Flour/Orange Fleshed Sweet Potato

CFC: Sweetened-Boiled and Oven-dried/Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

CFD: Non-Sweetened Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato

CFE: Sweetened Sun-dried and toasted Soy Flour/Orange Fleshed Sweet Potato



**Vitamin B<sub>9</sub> (Folic acid) content in the complementary food formulations:**

The vitamin B<sub>9</sub> (Folic acid) content of the complementary food blend ranged from 27.32 mg/100g in sample CON to 193.85 mg/100g in sample CFD. It was observed that the vitamin B<sub>6</sub> content increased significantly ( $p < 0.05$ ) in samples with sun-dried and toasted soy flour. The lowest vitamin B<sub>6</sub> content was observed for sample CON 0.30 mg/100g and it was significantly ( $p < 0.05$ ) different from all the other formulations. This result agreed with those observed by Okoye et al., [18]. The samples with sun-dried and toasted soy flour were significantly ( $p < 0.05$ ) different/higher than the samples that were made of boiled and oven-dried soy flour. The vitamin B<sub>6</sub> content of the complementary food formulations depends slightly on the quantity of soy bean flour and its processing method since boiling caused some of the vitamins to leach out. Folic acid functions as a co-enzyme in the body. There were significant ( $p < 0.05$ ) differences between the complementary food formulations made of boiled and oven-dried soy flour and sun-dried and toasted soy flour [1].

**Vitamin C (Ascorbic acid) content in the complementary food formulations:**

The vitamin C (Ascorbic acid) content of the complementary food blend ranged from 11.03 mg/100g in sample CFA to 65.15 mg/100g in sample CON. It was observed that the vitamin C content increased significantly ( $p < 0.05$ ) in samples with addition of date palm pulp flour. The lowest vitamin C content was observed for sample CFA (11.03 mg/100g) and it was significantly ( $p < 0.05$ ) different from all the other formulations. The vitamin C content of the complementary food formulations was generally low compared to that of sample CON. This result agreed with those observed by Okoye et al., [18]. Sample CON (Nestle Cerelac) contained more than the RDI; this could be because it is a fortified food [1]. The RDI of vitamin C for infants of 7–12 months is 50 mg/day. Vitamin C helps to form and repair red blood cells, bones and tissues; it also helps cuts and wounds to heal, boosts the immune system and keep infections away. Ascorbic acid is important in the prevention of scurvy. There were significant ( $p < 0.05$ ) differences between the complementary food formulations not containing date palm pulp flour and those containing date palm pulp flour [18].

**Mineral Composition of complementary food formulations:**

The research study also

evaluated the mineral composition of the bread samples as shown in Table 4. The minerals that were determined included potassium, calcium, sodium, magnesium, phosphorus, and Iron.

**Calcium (Ca) content in the complementary food formulations:**

The Calcium, content of the complementary food blend ranged from 261.84 mg/100g in sample CFA to 351.32 mg/100g in sample CON. It was observed that the Calcium content increased significantly ( $p < 0.05$ ) in samples with addition of date palm pulp flour. The lowest Calcium content was observed for sample CFA (261.84 mg/100g) and it was significantly ( $p < 0.05$ ) different from all the other formulations. All the samples contained more than the minimum recommended daily intake [14]. The RDI of Calcium for infants of 7–12 months is 260 mg/day. Calcium helps to build strong and teeth. Calcium is also necessary for the optimal growth and development of infants and young children. There were significant ( $p < 0.05$ ) differences between the complementary food formulations not containing date palm pulp flour and those containing date palm pulp flour. All these minerals are needed for the osmotic regulation of the body processes.

**Potassium (K) content in the complementary food formulations:**

The Potassium content of the complementary food blend ranged from 632.85 mg/100g in sample CON to 718.23 mg/100g in sample CFE. It was observed that the Potassium content increased significantly ( $p < 0.05$ ) in samples with sun-dried and toasted soy flour. The lowest Potassium content was observed for sample CON (632.85 mg/100g) and it was significantly ( $p < 0.05$ ) different from all the other formulations. The samples with sun-dried and toasted soy flour were significantly ( $p < 0.05$ ) different/higher than the samples that were made of boiled and oven-dried soy flour. The Potassium content of the complementary food formulations depends slightly on the quantity of date palm pulp flour. There were significant ( $p < 0.05$ ) differences between the complementary food formulations containing date palm pulp flour and those without date palm pulp flour. Potassium is an indispensable micronutrient that is necessary for cell metabolism by maintaining the sodium potassium pump and electrolytes.

**Sodium (Na) content in the complementary food formulations:**

The Sodium content of the complementary food blend ranged from 35.85 mg/100g in sample CFA to 190.11 mg/100g in sample CON. It was observed that the Sodium

content increased significantly ( $p < 0.05$ ) in samples with addition of date palm pulp flour. The lowest Sodium content was observed for sample CFA (35.85 mg/100g) and it was significantly ( $p < 0.05$ ) different from all the other formulations. All the samples contained more than the minimum recommended daily intake [14]. Sodium is an indispensable micronutrient that is necessary for cell metabolism by maintaining the sodium potassium pump and electrolytes.

**Phosphorus (P) content in the complementary food formulations:** The Phosphorus content of the complementary food blend ranged from 292.87 mg/100g in sample CON to 546.72 mg/100g in sample CFE. It was observed that the phosphorus content increased significantly ( $p < 0.05$ ) in samples with addition of date palm pulp flour. The lowest phosphorus content was observed for sample CON (292.87 mg/100g) and it was significantly ( $p < 0.05$ ) different from all the other formulations. The samples with sun-dried and toasted soy flour were not significantly ( $p < 0.05$ ) different/higher than the samples that were made of boiled and oven-dried soy flour. The phosphorus content of the complementary food formulations depends slightly on the quantity of date palm pulp flour.

**Iron (Fe) content in the complementary food formulations:** The Iron content of the complementary food blend ranged from 10.73 mg/100g in sample CON to 19.91 mg/100g in sample CFE. It was observed that the iron content increased significantly ( $p < 0.05$ ) in samples with addition of date palm pulp flour. The lowest iron content was observed for sample CON (10.73 mg/100g) and it was significantly ( $p < 0.05$ ) different from all the other formulations. The samples with sun-dried and toasted soy flour were not significantly ( $p < 0.05$ ) different/higher than the samples that were made of boiled and oven-dried soy flour. The iron content of the complementary food formulations depends slightly on the quantity of date palm pulp flour.

**Zinc (Zn) content in the complementary food formulations:** The Zinc content of the complementary food blend ranged from 5.33 mg/100g in sample CFA to 25.34 mg/100g in sample CON. It was observed that the Zinc content increased significantly ( $P < 0.05$ ) in samples with addition of date palm pulp flour. The lowest Zinc content was observed for sample CFA (5.33 mg/100g) and it was significantly ( $p < 0.05$ ) different from all the other

formulations. All the samples contained more than the minimum recommended daily intake [14]. "Zinc is an indispensable micronutrient that is necessary for protein synthesis, cell growth and differentiation, immune function and intestinal transport of water and electrolytes. Symptoms of zinc deficiency include dermatitis, retarded growth, diarrhea, etc. All age groups of the population are at risk of zinc deficiency, but infants and young children are the most vulnerable. Zinc supplementation trials conducted over the last few decades in children from developing countries have clearly demonstrated the positive benefits of improved zinc status, including improved growth rates and reductions in the incidence of various infectious diseases" [19].

**Iodine (I<sub>2</sub>) content in the complementary food formulations:** The Iodine content of the complementary food blend ranged from 47.32 ( $\mu\text{g}/100\text{g}$ ) in sample CON to 90.19 ( $\mu\text{g}/100\text{g}$ ) in sample CFE. It was observed that the iodine content increased significantly ( $P < 0.05$ ) in samples with addition of date palm pulp flour. The lowest iodine content was observed for sample CON (47.32  $\mu\text{g}/100\text{g}$ ) and it was significantly ( $p < 0.05$ ) different from all the other formulations. The samples with sun-dried and toasted soy flour were not significantly ( $p < 0.05$ ) different/higher than the samples that were made of boiled and oven-dried soy flour. The iodine content of the complementary food formulations depends slightly on the quantity of date palm pulp flour. "Iodine is a mineral contained in some foods that the body need in order to produce thyroid hormones. This hormone regulates the body's metabolism as well as many other vital activities. Thyroid hormones are also required by the body for appropriate bone and brain development during pregnancy and infancy. The dietary reference value (DRV) is 150 $\mu\text{g}$  of iodine per day for adults over the age of 18. During pregnancy and lactation, requirements increase to 200 $\mu\text{g}$  per day" (food code Service, 2013).

#### 4. CONCLUSION

Acceptable complimentary food can be produced from sweet potatoes, soybean flour and date fruit using an optimum level of 48% OFSP, 42 % boiled and oven-dried soy flour and 10% date flour (sample CFB). The formulated samples improved the physicochemical properties of complementary foods and thus improved overall nutritional profile of food. The experimental samples were very rich in vitamin A which is an

essential nutrient that helps to build up the immune system of infants against a number of infections and sustains the integrity of the epithelial linings. The content of essential minerals in the samples included calcium, phosphorus, sodium, iron, zinc, iodine, and potassium. These minerals play crucial roles in osmotic regulation, various physiological functions in the body and are necessary for maintaining overall health and well-being.

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Okoronkwo NC, Okoyeuzu CF, Eze CR, Mbaeyi-nwaoha IE, Agbata CP. Quality evaluation of complementary food produced by solid-state fermentation of fonio, soybean and orange-fleshed sweet potato blends. *Fermentation MDPI Journal*. 2023;9(10.3390):250–270.
2. Amagloh Francis, Kweku JC. Sweet potato-based complementary food for infants in Ghana. *International Journal of Food Sciences and Nutrition*. 2012;33(1): 3–10.
3. Damak AMA, Akubor PI, Ariahu CC, Okereke GO. Functional and nutritional properties of various flour blends of arrowroot starch and wheat flour. *AFSJ*. 2022, Apr 28;21(6):12-25. Available:<https://journalafsj.com/index.php/AFSJ/article/view/545>
4. Chiedu UC, Abiodun MS, Steve IO, Eguono EE, Joy AT. Physicochemical, nutritional and functional properties of composite flour blends from whole wheat, sweet potato, defatted peanut and rice bran. *Eur. J. Nutr. Food. Saf.* 2023, Dec 14;15(12):41-55. [cited 2024 Jun. 6] Available:<https://journalenfs.com/index.php/EJNFS/article/view/1366>
5. Alemayehu Eleni KA, TA. Complementary food product development from sorghum enriched with chickpea and orange-fleshed sweet potato: The case of dawuro zone, South Western Ethiopia complementary food product development from sorghum enriched with chickpea and orange-fleshed swe. *Jimma University College of Agriculture and Veterinary Medicine Journal*. 2014;7(2):1–119.
6. Owhero JO, Akpogheli PO, Edo GI, Ojulari AE, Agbo JJ. Proximate, mineral, sensorial and microbiological properties of chin-chin produced from okra seed and wheat flour blends. *Food Chemistry Advances*. 2023, Oct 1;2:100298.
7. Olatunde RO, Oyewole SJ, Abioye OD, BV, AG. Quality evaluation of sweet potato-based complementary food. *Agrosearch International Journal*. 2020; 20(1):94–105.
8. Ocheme OB, Adedeji OE, Chinma CE, Yakubu CM, Ajibo UH. Proximate composition, functional, and pasting properties of wheat and groundnut protein concentrate flour blends. *Food Science & Nutrition*. 2018, Jul;6(5):1173-8.
9. Mmari Mercy Wilfred, John N, Kinyuru JO, HL. Nutrient composition and consumer acceptability of soybean- sweet potato based complementary food fortified with longhorn grasshopper (*ruspolia differens*). *Jomo Kenyatta University of Agriculture and Technology Journal*. 2017;2(3):322–3902.
10. Abdul Z, Wen K, Atan H, Mat K, Khalid HM, Dini N, Anis S, Sukri M, Che H, Seong L, Hakim M, Mohd B, Hanafiah A, Mijanur M, Khairul M, Abdul A, Wee W, Shahman N, Ahmad N, Dawood MAO. Palm date meal as a non-traditional ingredient for feeding aquatic animals: A review. *Aquaculture Reports*. 2022;25(June): 101233. Available:<https://doi.org/10.1016/j.aqrep.2022.101233>
11. Marcel R Mary, JSC, CEO. Nutritional evaluation of complementary porridge formulated from orange- - fleshed sweet potato, amaranth grain, pumpkin seed, and soybean flours. *Food Science & Nutrition, Researchgate*. 2021;1–18. Available:<https://doi.org/10.1002/fsn3.2675>
12. Truong VD, Avula RY, Pecota KV, Yencho GC, Lam IL. Sweetpotato Production, Processing, and Nutritional Quality: Vol. II; 2018.
13. Ahaotu I, Eze O, Maduka N. Quality assessment of corn-breadfruit-date flour and sensory evaluation of chin-chin

- prepared using the composite flours. *Advances in Biotechnology & Microbiology*. 2021;16(3). Available: <https://doi.org/10.19080/AIBM.2021.16.555940>
14. WFP. Nutritional guidance for complementary food. World Food Programme Cereal, Super Publication. 2018;68/70(February):1–4.
  15. Abuengmoh P, Ahure D, Igoli NN. Proximate, vitamin and mineral composition of bread produced from wheat, banana and mango flour blends. *International Journal of Food Science and Nutrition*. 2022;7(3):92–99. ISSN: 2455-4898 DOI: [www.foodsciencejournal.com](http://www.foodsciencejournal.com)
  16. Yakum KN, Ariaahu CC, Ariaahu EC, Igoli JO. Physicochemical properties of fortified coconut milk based chocolate - like drinks as influenced by cocoa powder and sugar levels. *American Journal of Food Science and Technology*. 2024;3(1): 19–29. Available: <https://doi.org/10.54536/ajfst.v3i1.2573>
  17. Bello AA, Gernah DI, Ariaahu CC, Ikya JK. Physico-chemical and sensory properties of complementary foods from blends of malted and non-malted sorghum, soybean and moringa oleifera seed flours. *American Journal of Food Science and Technology*. 2020;8(1):1–13. Available: <https://doi.org/10.12691/ajfst-8-1-1> food code Service, U. S. P. H. (2013). *Food Code*.
  18. Okoye JI, Egbujie AE, Ene GI. Evaluation of complementary foods produced from sorghum, soybean and irish potato composite flours. *Science World Journal*. 2021;16(3):206–211.
  19. Assohoun MCN, Djeni TN, Koussémon-camara M, Brou K. Effect of fermentation process on nutritional composition and aflatoxins concentration of doklu, a fermented maize based food. *Food and Nutrition Sciences*. 2013, November 4; 1120–1127.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/118742>