



Hydrogen Bond Energies in Formation of Water Molecule Clusters

Dimitar Mehandjiev ^a, Ignat Ignatov ^{b*}, Nikolai Neshev ^c, Georgi Gluhchev ^d
and Christos Drossiankis ^e

^a Institute of General and Inorganic Chemistry, Bulgarian Academy of Sciences, Acad. G. Bonchev Str., Bl. 11, Sofia 1113, Bulgaria.

^b Scientific Research Center of Medical Biophysics, 32 N. Kopernik Str., Sofia 1111, Bulgaria.

^c Sofia University "St. Kliment Ohridski", Faculty of Physics, Sofia, Bulgaria.

^d Institute of Information and Communication Technologies, Bulgarian Academy of Sciences, Acad., G. Bonchev Str., Bl. 2, Sofia 1113, Bulgaria.

^e IAWG- INTERNATIONALE Akademie für Wissenschaftliche Geistheilung, Hauptplatz 44, A-2293 Marchegg, Austria, Frankfurt, Germany.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/PSIJ/2021/v25i1230298

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83944>

Original Research Article

Received 24 October 2021
Accepted 27 December 2021
Published 29 December 2021

ABSTRACT

During the last decades, the interest in water structure has increased as an important theoretical and practical problem, connected with different industrial and biological applications. Theoretically, the mechanisms of water molecules cluster formation have been of special interest. It has been assumed that aggregation of water molecules depends on the energy of hydrogen bonds. The character of this process was investigated in the present study. The approach was based on measurements of the wetting angle θ of water droplets at different energy levels during their evaporation. Taking into account previous findings that θ depends on the average energy of hydrogen bonds between water molecules, it was assumed that the size of water clusters is related to the value of θ , measured at different energy levels. This assumption was confirmed by the obtained experimental results.

Keywords: Water; hydrogen bonds; water clusters; energy.

*Corresponding author: E-mail: mbioph@abv.bg;

1. INTRODUCTION

The interest in elucidating the structure of water has especially increased in connection with studies of hydrophobic boundary surfaces and in processes with practical application such as electrolysis, structuring of organic molecular crystals, non-polar media and theoretical analyses. New effects of water are also being described.

The wetting angle, also known as contact angle, quantitatively describes how liquids wet solid surfaces. In terms of geometry, it is the angle formed at the intersection point of the three phase boundaries between a liquid, solid and air. The wetting angle is a common predictor of the qualities and behavior of solid surfaces after physical or chemical treatment [1,2].

In 1983, Antonov showed that, during evaporation of water droplets on different surfaces, the wetting angle is related with the average energy of the hydrogen bonds. For this purpose, he and co-authors designed specialized equipment for real time measurements of this angle with sufficient accuracy. So, the relation between the wetting angle and the average energy of the hydrogen bonds during droplet evaporation can be expressed as follows:

$$\theta = \arccos(-1-bE) \quad (1)$$

where θ is the wetting angle, E is the average energy of the hydrogen bonds, b is a temperature-dependent parameter [3, 4, 5].

It has also been found that the wetting angle changes discretely as water droplets evaporate. The shape of these curves depends on the composition of investigated water and other liquids, and this has opened up new possibilities for studying biochemical and biological effects.

Studies have shown that clusters with different numbers of water molecules are formed on the basis of hydrogen bonds in bulk water.

The following spectral methods have been generally used to study water clusters - H-NMR [6, 7], far-infrared [8], vibration-rotation-tunneling (VRT) [9], neutron diffraction [10], SCC-DFTB Method [11,12], NES [13,14]. An earlier authors' study has estimated that, at ($E = -0.1387$ eV) ($\lambda = 8.95$ μm) ($\tilde{\nu} = 1117$ cm^{-1}), water clusters of approx. 21 water molecules are formed in catholyte water [15]. Based on this, a cluster

model was proposed with 20 water molecules in a dodecahedral structure with diameter of the circumscribed sphere equal to 0.822 nm.

This result corresponds to our result by applying the Antonov effect to "discrete" evaporation of water drops. Analyses for clusters of $n = 6-20$ water molecules are available in a number of publications [16, 17, 18]. Cases of $n = 1-6$ have been described in [19].

It has also been found that at $n = 20$ about 25% of water molecules are structured with this number [20].

The question arises to what extent the discrete change in the wetting angle during evaporation of water droplets depends on the formation of clusters with different numbers of water molecules. This is the purpose of the present study.

2. MATHERIALS AND METHODS

Electrochemically activated catholyte water [21,22] with the following parameters was used for the study: pH = 9.30, oxidation reduction potential (ORP) = - 450 mV.

The wetting angle θ was measured with a specially designed apparatus which is described in detail in [13,14, 20]. Measured wetting angles of 10 water droplets during their evaporation from a solid surface varied from 74 to 0 deg. The measurements were sequential and N denotes the number of measurements taken from the initial value of θ_0 to 0. The time interval for each reading was 10minutes. From the obtained values of θ , the average energy of the hydrogen bonds E in eV was calculated according to the formula:

$$E = \gamma^0 (1+\cos\theta)/\tau(1+\cos\theta_0) \quad (2)$$

where θ is the wetting angle, E is the energy of the hydrogen bonds.

The expression $\tau (1+\cos\theta_0) / \gamma^0$ is equal to 14.33 eV at 20°C.

Evaporation of water drops was performed in a hermetic chamber with stable temperature of 20°C [21]. The drops were placed on BoPET (biaxially-oriented polyethylene terephthalate) foil with 350 μm thicknesses.

3. RESULTS AND DISCUSSION

The average weight reduction values of the investigated water drops during their evaporation and to the corresponding values of the wetting angle are shown in Table 1. The average initial height of the droplets was $h_i=2.30$ mm and their average final height was $h_f=0.26$ mm. The layering structure of water has a periodicity of 0.30 ± 0.03 nm [23].

Fig. 1 shows the dependence of the wetting angle on the sequential number of the corresponding measurement. This angle decreased from 74 deg to 10 deg and the dependence had a step character.

Table 1. Average weight reduction values of water drops during their evaporation and the corresponding values of the wetting angle

Wetting angle (°) θ	Weight mg
74	48.0
66	37.9
58	29.0
50	21.3
42	14.8
34	9.4
26	5.3
18	2.4
10	0.6




Fig. 1. Dependence of the wetting angle θ on the sequential number of the corresponding measurement

Fig. 2 shows the change in the average energy of the hydrogen bonds calculated with formula (2).

As it can be seen from the above figure, the average energy ($-E$) of the hydrogen bonds increases from 0.0912 to 0.1387 eV. Its dependence on the wetting angle is shown in Fig. 3.




Fig. 2. Dependence of the average energy of hydrogen bonds E on the sequential number of the corresponding measurement




Fig. 3. Dependence of the average energy of hydrogen bonds on the wetting angle θ




Fig. 4. Dependence of ΔE on N

It illustrates that the smaller the wetting angle, the higher the average hydrogen bonds energy is. From the point of view of the physical process, when the water evaporates from the drop, the energy of the hydrogen bonds increases. The dependence of the wetting angle is not linear, as can be seen from Fig. 1. The wetting angle decreases with greater values as evaporation of water from the drop advances. However, the energy of the hydrogen bonds E increases linearly, as shown in Fig. 2. Therefore, the energy of the hydrogen bonds increases stepwise, but with a constant value. Fig. 4 shows the difference at each step of the change of E .

As can be seen from Fig. 4, ΔE is a constant value during the whole process of evaporation of water drops. Then the question arises about what determines this constancy in the magnitude of ΔE . As it was pointed above, clusters of different numbers of water molecules form in bulk water. They also determine the amount of average energy of the hydrogen bonds. If the evaporation of water from water drops causes disintegration of these clusters, each step will correspond to the loss of a single water molecule. Then the average energy of the hydrogen bonds in the system will decrease with the value of the energy of the bond that had been associated with this water molecule in the cluster.

If this release energy is equal to 0.0025 eV, as can be seen from Fig. 4, then this will also determine the step in the change of E , which is calculated from the wetting angle θ . This fact makes it possible to calculate the number of water molecules involved in the formation of clusters for each energy presented in Fig. 2. If

the corresponding values are divided by 0.0025 and then by 2, because each hydrogen bond is associated with two water molecules, numbers are obtained that correspond in order of magnitude to the numbers of molecules that form particular clusters in bulk water.

Therefore, it can be assumed that the stepwise nature of the change in the energy of the hydrogen bonds is due to the disintegration of the existing clusters in bulk water. This determines the nature of the change and the wetting angle of water drops during evaporation.

4. CONCLUSION

Clusters of different numbers of water molecules are formed in water on the basis of hydrogen bonds. This results in different values of the average energy of the hydrogen bonds, proved by measurements of the wetting angle during water droplets evaporation from surfaces of different materials. When droplets evaporate, disintegration of water clusters takes place through separation of water molecules from them.

This changes the average energy of the hydrogen bonds and causes its stepwise change. On this basis, the abrupt changes in the wetting angle during evaporation of water droplets can be explained. Clusters can be formed in water, composed of more than 20 water molecules and with a high average hydrogen bonds energy.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Gao Y, Fang H, Ni K. A hierarchical clustering method of hydrogen bond networks in liquid water undergoing shear flow. *Scientific Reports*. 2021;11: 9542.
- Jelev J, Antonov A, Galabova T. Method and Device for Evaluation of Bio-Psychophysical Influence of Radio, Television and Media Products Upon Humans. US20070276192A1; 2003.
- Antonov A, Yuskesselieva L, Method for determination of structural changes in liquids, Author's certificate of invention. 1983;43821.
- Antonov A. An optical method version for determination of the wetting angle of liquids. *Comptes Rendus de l'Académie Bulgare des Sciences*. 1984;37:1199.
- Antonov A, Yuskesselieva L, Teodossieva I. Influence of Ions on the Structure of Water Under Conditions Far away from Equilibrium. *Physiologie*. 1989;26(4):255-260.
- D' Angelo P, Zitolo A, Aquilanti G, Migliorati V. Using a combined theoretical and experimental approach to understand the structure and dynamics of imidazolium-based ionic liquids/water mixtures. 2. EXAFS Spectroscopy, *The Journal of Physical Chemistry B*. 2013;117(41): 12516-12524.
Available: <https://doi.org/10.1021/jp404868a>
- Turov V, Krupskaya T, Barvinchenko V, Lipkovska N, Kartel M, Suvorova L. Peculiarities of water cluster formation on the surface of dispersed KCl: The influence of hydrophobic silica and organic media. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*. 2016;499:97-102.
Available: <https://doi.org/10.1016/j.colsurfa.2016.03.069>
- Liu K, Fellers R, Viant M, McLaughlin R, Brown M, Saykally R. A long path length pulsed slit valve appropriate for high temperature operation: Infrared spectroscopy of jet-cooled large water clusters and nucleotide bases. *Review of Scientific Instruments*. 1998;67(2).
Available: <https://doi.org/10.1063/1.1146605>
- Liu K, Cruzan J, Saykally R. Water clusters. *Science*. 1996;271(5251):929-933.
DOI: 10.1126/science.271.5251.929
- Yoshida K, Ishuda S, Yamaguchi T. Molecular Physics. Hydrogen bonding and clusters in supercritical methanol-water mixture by neutron diffraction with H/D substitution combined with empirical potential structure refinement modeling. 2019;117(22):3227-3310.
Available: <https://doi.org/10.1080/00268976.2019.1633481>
- Choi T, Jordan K.) Application of the SCC-DFTB Method to $H^+(H_2O)_6$, $H^+(H_2O)_{21}$, and $H^+(H_2O)_{22}$, *The Journal of Physical Chemistry B*. 2010;114(20):6932-6936.
Available: <https://doi.org/10.1021/jp912289e>
- Goyal P, Elstner M, Coi Q. J. *Phys. Chem. B*. Application of the SCC-DFTB method to neutral and protonated water clusters and bulk water. 2011;115(20):6790-6805.
Available: <https://doi.org/10.1021/jp202259c>
- Todorov S, Damianova A, Sivriev I, Antonov A, Galabova T. Water energy spectrum method and investigation of the variations of the H-bond structure of natural waters. *Comptes Rendus de l'Académie Bulgare des Sciences*. 2008; 61(5251):857-862.
- Gramatikov P, Antonov A, Gramatikova A. A study of the properties and structure variations of water systems under the stimulus of outside influences. *Fresenius' Journal of Analytical Chemistry*. 1992; 343(1):134-135.
- Ignatov I, Gluhchev G, Neshev N, Mehandjiev D. Structuring of water clusters depending on the energy of hydrogen bonds in electrochemically activated waters Anolyte and Catholyte. *Bulgarian Chemical Communications*. 2021;53(20): 234-239.
DOI: 10.34049/bcc.53.2.5376
- Masella M, Flament JP. *J. Chem. Phys.* A pairwise and two many-body models for water: Influence of nonpairwise effects upon the stability and geometry of $(H_2O)_n$ cyclic ($n=3-6$) and cagelike ($n=6-20$) clusters. 1997;107:9105.
Available: <https://doi.org/10.1063/1.475202>
- Koyasu K, Atobe J, Furuse S, Nakajima A. Anion photoelectron spectroscopy of transition metal- and lanthanide metal-silicon clusters: MSi_n^- ($n=6-20$). *J. Chem. Phys.* 2008;129(21):214301.
Available: <https://doi.org/10.1063/1.3023080>

18. Neela YI, Mahadvi AS, Sasty GN. Hydrogen bonding in water clusters and their ionized counterparts. J. Phys. Chem. B. 2010;114(51):17162–17171. Available:<https://doi.org/10.1021/jp108634z>
19. Kawai Y, et al. Reactions of protonated water clusters $H^+(H_2O)_n$ ($n=1-6$) with dimethylsulfoxide in a guided ion beam apparatus. Chemical Physics Letters. 2003; 377(1-2):69-73. Available:[https://doi.org/10.1016/S0009-2614\(03\)01095-9](https://doi.org/10.1016/S0009-2614(03)01095-9)
20. Antonov A. Dissertation thesis for degree Doctor of physical sciences. Blagoevgrad. 1995;1-254.
21. Kimbrough et al. The effect of electrolysis and oxidation–reduction potential on microbial survival, growth, and disinfection. International Journal of Environment and Pollution. 2006;211-221.
22. Lapin AA, Kalayda AA, Filippov SD, Zelenkov VN. Biochemical effects of molecular hydrogen in aqueous systems. IOP Conference Series: Earth and Environmental Science. 2019;288: 012054.
23. Uhling M, et al. Atomically resolved interfacial water structures on crystalline hydrophilic and hydrophobic surfaces. Nanoscale. 2021;13:5275-5282. DOI: 10.1039/D1NR00351H

© 2021 Mehandjiev et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83944>