

Impact of Nitrogen and Silicon Levels on Yield and Relative Economics of Transplanted Rice under Temperate Conditions

Rukhsana Jan ^{a*}, Shamshir Ul Hussan ^b, Kawsar Rasool ^c, Habibullah ^d
and Mehvish Mansoor ^e

^a Division of Agronomy, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir (SKUAST-K), Shalimar-190025, Srinagar, UT-J and K, India.

^b Division of Genetics and Plant Breeding, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir (SKUAST-K), Shalimar-190025, Srinagar, UT-J and K, India.

^c Division of Entomology, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu (SKUAST-K), Shalimar-190025, Srinagar, UT-J and K, India.

^d CSIR-Indian Institute of Integrative Medicine, Field station, Bonera-Pulwama, India.

^e Division of Soil Science, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu (SKUAST-K), Shalimar-190025, Srinagar, UT-J and K, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2022/v41i383979

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93293>

Original Research Article

Received 23 August 2022
Accepted 29 October 2022
Published 01 November 2022

ABSTRACT

The field experiment entitled "Impact of Nitrogen and Silicon application on yield and relative economics of transplanted rice under temperate conditions" was conducted at Agronomy Research Farm of Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir during Kharif seasons 2014-2015. The soil of the experiment was silty clay loam in texture, neutral in reaction with medium available nitrogen (442.88kg ha^{-1}), phosphorus (9.3kg ha^{-1}), potassium (221.60kg ha^{-1}) and low available silicon (280.23kg ha^{-1}). The experiment was laidout in Factorial Randomized Completely Block Design assigning combinations of three nitrogen levels viz. 120, 150 and 180N kg ha^{-1} and four silicon levels viz. Control, 5%, 10% and 15% Si with four replications.

*Corresponding author: E-mail: rukhsanamalik2323@gmail.com;

The results revealed that higher grain yield (76.00q ha^{-1}), straw yield (93.97q ha^{-1}) and harvest index (44.84%) were recorded with 120Nkg ha^{-1} . So far as silicon levels are concerned, 15%Si observed higher grain yield (76 q ha^{-1}), straw yield (93.97 q ha^{-1}) and harvest index (44.84%). Relative economics revealed that 120Nkg ha^{-1} with 15%Si realized higher benefit cost ratio (Rs. 2.12).

Keywords: Benefit cost ratio; nitrogen; rice; silicon and yield.

1. INTRODUCTION

“Rice is a staple food for more than half of the world’s population. Globally, it is grown on acreage of 164.19mha with total production of 509.87 million tonnes and productivity 3.10 ton per hectare” [1]. “In Jammu and Kashmir rice is grown on an area of 0.3million ha with a production of 0.6 million tonnes and productivity of 2 tonnes per hectare” [1]. “It is essential to the rice plant, with about 75% of leaf nitrogen associated with chloroplasts, which are physiologically important in dry matter production” [2]. “There are many factors affecting rice yield and quality, such as variety, accumulation temperature during the growth period, and relatively minor factors such as cultivation techniques (fertilizer rates, fertilization period, water management, etc.)” Laenoi et al., [3]; Zhu et al. [4]; Cao et al., 2017 and Zhou et al., [5]. Due to the effect of increased yield brought by the application of nitrogen fertilizer, an increasing number of nitrogen fertilizers have been applied in rice production.

“Silicon is the second most abundant element in soil after oxygen. Rice is considered to be a silicon accumulator plant and tends to actively accumulate Si to tissue concentrations of 5% or higher” [6]. “Reduced amount of silicon in plant produces necrosis, disturbance in leaf photosynthetic efficiency, growth retardation and reduces grain yield in cereals especially rice” [7]. “The positive effects of silicon on rice growth and production were manifested when it was specifically supplied during the reproductive growth stage (panicle initiation to heading) as compared to vegetative and ripening stages. Panicle initiation stage application exerted a feed-forward effect on photosynthesis coupled with increase in both stomatal conductance and biochemical capacity to fix CO_2 as reported” by Lavinsky et al., [8], who also surmised that “proper levels of Si in reproductive structures played an unidentified role in increasing the yields of rice. Silicon also has a major role in increasing yield attributing characters like

number of spikelets, filled spikelet percentage, test weight and total grain yield” [9,10,11] in rice. Keeping in view the above facts, the study was designed with the objective to study the impact of Nitrogen and Silicon levels on yield and relative economics of transplanted rice under temperate conditions.

2. MATERIALS AND METHODS

The field experiment was carried out at Research Farm of Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar during 2014-2015. The mean maximum (27.50°C and 27.30°C), mean minimum temperatures (12.60°C and 13.20°C), mean maximum relative humidities (82.50% and 82.80%) and mean minimum relative humidities (55.30% and 56.20%) were recorded for the entire cropping seasons during 2014 and 2015 respectively. The total rainfall received during the experimentation period was 33.3mm during 2014 and 29.5mm during 2015. The experiment comprised two factors (three nitrogen levels and four silicon levels) was laid out in a randomized completely block design with four replications. The factors included three nitrogen levels: N_1 : 120kg/ha, N_2 :150kg/ha and N_3 :180kg/ha and four silicon levels: Si_0 : Control, Si_1 :5%, Si_2 :10% and Si_3 :15%. The following observations were recorded during the experiment are:

Grain yield: Grain yield of each plot was recorded separately as kg plot^{-1} and then converted into q ha^{-1} .

Straw yield: Straw yield of each plot was computed by deducting the grain yield from biological yield and expressed as q ha^{-1} .

Harvest index: It was calculated by the following formula:

$$\text{H.I} = \text{Economic yield} / \text{Biological yield} \times 100$$

2.1 Relative Economics

Economics of different treatments was worked out on the basis of grain and straw yield per

hectare. The cost of input and output was estimated as per prevailing market rates. The benefit : cost ratio (returns per rupee invested) was determined as:

$$\text{Benefit cost ratio} = \frac{\text{Net returns}}{\text{Total cost of cultivation}}$$

2.2 Statistical Analysis

The data obtained in respect of various observations were statistically analysed by the method described by Cochran and Cox [12] using Ne-software. The significance of 'F' and 'T' was tested at 5% level of significance. The critical difference value was determined wherever 'F' test was found to be significant.

3. RESULTS AND DISCUSSION

From the Table 1 and Fig. 1 indicated that the grain yield, straw yield and harvest index showed the significant variation among the nitrogen and silicon levels. Significantly higher grain yield (76.00q ha⁻¹), straw yield (93.97q ha⁻¹) and harvest index (44.84%) were observed with 120kg N ha⁻¹ followed by 150kg N ha⁻¹ (grain yield: 74.03q ha⁻¹, straw yield: 91.66q ha⁻¹ and harvest index: 44.51%). It is because of higher yield attributing characters which ultimately caused the increase of rice grain yield. Similar results were reported by Huang et al., [13]. "Also, the increment of grain yield by the application of nitrogen upto a certain level was reported" by Zhaowen et al., [14]. Similar trend of the effect of nitrogen levels on straw yield were also reported by Pramanik and Bera [15]. However, harvest

index were significantly higher with 120kgN ha⁻¹. Rahman et al., [16], also reported similar trend. Significantly lower grain yield (68.83q ha⁻¹) straw yield (88.19q ha⁻¹) and harvest index (43.96%) were recorded with 180kg N ha⁻¹. Among different silicon levels, 15% Si recorded the higher grain yield (76.76q ha⁻¹), straw yield (94.72q ha⁻¹) and harvest index (44.80%) followed by 10%Si (grain yield: 74.58q ha⁻¹, straw yield: 92.89q ha⁻¹, and harvest index: 44.40%) being at par with 5%Si (grain yield:73.21q ha⁻¹, straw yield:91.96q ha⁻¹, and harvest index:44.24%). It is because of silicon is responsible to control stomatal activity, photosynthesis, water use efficiency which ultimately results in better vegetative growth results in higher yield. And it may also be attributed to leaf erectness which facilitated better penetration of sunlight leading to higher photosynthetic activity of plant and higher production of carbohydrates. Similar results were also noticed by Singh et al., [17] and Pati et al., [18]. Significantly lower grain yield (67.27q ha⁻¹) straw yield (85.52q ha⁻¹) and harvest index (43.84%) were observed with control [19].

The data relative economics presented in Table 2 revealed that benefit cost ratio remained higher (Rs.2.13) when 120 kg N ha⁻¹ and 15% Si were applied. The same treatment also recorded the higher net returns of Rs.106107. The efficiency of a treatment is finally decided in terms of economics (benefit cost ratio) of the treatment and it might be due to the higher grain and straw yield and minimum cost of nitrogen fertilizers as cost of silicon does not have a greater effect.

Table 1. Impact of nitrogen and silicon levels on grain yield, straw yield and harvest index of transplanted rice (Pooled data of two years)

Treatments	Grain yield (q ha ⁻¹)	Straw yield (q ha ⁻¹)	Harvest index (%)
Nitrogen levels (kg ha⁻¹)			
N ₁ :120	76.00	93.97	44.84
N ₂ :150	74.03	91.66	44.51
N ₃ :180	68.83	88.19	43.96
SEm±	0.51	0.54	0.09
CD(p≤0.05)	1.52	1.59	0.28
Silicon levels			
Si ₀ :Control	67.27	85.52	43.84
Si ₁ : 5%	73.21	91.96	44.24
Si ₂ :10%	74.58	92.89	44.40
Si ₃ :15%	76.76	94.72	44.80
SEm±	0.57	0.63	0.12
CD(p≤0.05)	1.71	1.84	0.37




Fig. 1(a)




Fig. 1(b)

Fig. 1. (a and b). Impact of nitrogen and silicon levels on grain yield, straw yield and harvest index of transplanted rice (Pooled data of two years)

Table 2. Relative economics of transplanted rice as influenced by nitrogen and silicon levels (pooled data of two years)

Treatments	Cost of Cultivation (Rs. ha ⁻¹)	Gross returns (Rs. ha ⁻¹)	Net returns (Rs. ha ⁻¹)	B:C
N ₁ Si ₀	48522	138929	90407	1.86
N ₁ Si ₁	49146	148280	99134	2.01
N ₁ Si ₂	49490	150174	100684	2.03
N ₁ Si ₃	49834	155941	106107	2.13
N ₂ Si ₀	48687	132364	83677	1.71
N ₂ Si ₁	49311	145581	96270	1.95
N ₂ Si ₂	49655	147330	97675	1.96
N ₂ Si ₃	49999	153206	103207	2.06
N ₃ Si ₀	48852	128447	79595	1.62
N ₃ Si ₁	49476	137549	88073	1.78
N ₃ Si ₂	49820	139223	89403	1.79
N ₃ Si ₃	50164	140566	90402	1.80

4. CONCLUSION

From this study we concluded that silicon applications can significantly improve rice yield if applied at proper time with feasible concentration. Optimum nitrogen fertilizer is an important practice for increasing rice yield. Considering the influence of nitrogen and silicon levels, it can be suggested that for realizing economically higher grain yield of rice with 120Nkg ha⁻¹ and 15% Si based nutrient management, as it helped to achieve the higher net returns (Rs. 106107) and the higher B:C ratio (Rs. 2.13).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- DES. Directorate of economics and statistics. Econ Surv. 2020;123:129-30.
- Dalling MJ. The physiological basis of nitrogen redistributing during filling in cereals. In: Exploitation of physiological and genetic variability to enhance crop productivity. Rockville, MD: American Society of Plant Physiologists. 1995;55-71.
- Laenoi S, Rerkasem B, Lordkaew S, Prom-u-thai C. Seasonal variation in grain yield and quality in different rice varieties. Field Crops Res. 2018;221:350-7. DOI: 10.1016/j.fcr.2017.06.006
- Zhu DW, Zhang HC, Guo BW, Xu K, Dai QG, Wei HY, et al. Effects of nitrogen level on yield and quality of japonica soft super rice. J Integr Agric. 2017;16(5): 1018-27. DOI: 10.1016/S2095-3119(16)61577-0
- Zhou CC, Huang YC, Jia BY, Wang Y, Wang Y, Xu Q, et al. Effects of cultivar, nitrogen rate, and planting density on rice-grain quality. Agronomy. 2018;8(11): 246. DOI: 10.3390/agronomy8110246
- Epstein E. Silicon in plants: facts vs. concepts. In: Datnoff LE, Synder GH, Korndorfer, editors. Silicon in agriculture; Elsevier. Amsterdam. 2002;1-15.
- Shashidhar HE, Chandrashekhar N, Narayanaswamy C, Mehendra AC, Prakash NB. Calcium silicate as silicon source and its interaction with nitrogen in aerobic rice. Silicon in. Agriculture: 4th International Conference 26-31 October, South Africa pp. 2008;93.
- Lavinsky AO, Detmann KC, Reis JV, Ávila RT, Sanglard ML, Pereira LF, et al. Silicon improves rice grain yield and photosynthesis specifically when supplied during the reproductive growth stage. J Plant Physiol. 2016;206:125-32. DOI: 10.1016/j.jplph.2016.09.010, PMID 27744227.
- Rani YA, Narayanan A, Devi V, Subbaramamma P. The effect of silicon application on growth and yield of rice plants. Annals Plant Physiol. 1997;11:125-8.
- Ahmad A, Afzal M, Ahmad AUH, Tahir M. Effect of foliar application of silicon on yield and quality of rice (*Oryza sativa* L.). Cercet Agronomic Moldova. 2013;XLVI(3):155.
- Jawahar S, Vijayakumar D, Bommera R, Jain N, Jeevanandham. effect of silixol granules on growth and yield of rice. Int J Curr Res Acad Rev. 2015;3(1):168-74.
- Cochran WG, Cox GM. Experimental designs. 2. Ufl. 1964;1964:XIV + 617S.,Preis 68 s.
- Huang M, Zou YB, Jiang P, Bing Xia I, Ao HJ. Relationship between grain yield and yield components in super hybrid rice. Agric Sci China. 2011;1537-1544;10.
- Zhaowen MO, Shenggang P, Bingshun L, Hua T, Meiyang D, Qingshan L et al. Growth and Yield response of five super rice (*Oryza sativa* L.) varieties to nitrogen fertilizer rates. J Food Agric Environ. 2013;11(3-4):1257-62.
- Pramanik K, Bera AK. Effect of seedling age and nitrogen fertilizer on growth, chlorophyll content, yield and economics of hybrid rice (*Oryza sativa* L.). Int J Agron Plant Prod. 2013;4:3489-99.
- Singh R, Ahuja U, Ahuja SC. Basmati for prosperity. Indian Farming. 2006;56: 33-6.
- Pati S, Pal B, Badole S. Effect of silicon fertilization on growth, yield and nitrogen uptake of rice. Commun Soil Sci Plant Anal. 2015;47(3):1081-6.
- Rahman MH, Khatun MM, Mamun MAA, Islam MZ, Islam MR. Effect of number of seedling hill-1 and nitrogen levels on growth and yield of BRRI dhan 32. J Soil Nat. 2007;1(2):01-7.

19. Cao XM, Sun HY, Wang CG, Ren XJ, Liu HF, Zhang ZJ. Effects of late-stage nitrogen fertilizer application on the starch structure and cooking quality of rice. J Sci Food Agric. 2018;98(6):2332-40. DOI: 10.1002/jsfa.8723, PMID 28991369.

© 2022 Jan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93293>