



Genetic Insights for Nutritional Traits in Elite Rice (*Oryza sativa* L.) Crosses Using Generation Mean Analysis

N. Lingaiah^{a+++*}, Ch. Surender Raju^{b#},
N. Sarla^{c†}, K. Radhika^{d‡}, V. Venkanna^{e^}
and D. Vishnu Vardhan Reddy^{f###}

^a Agricultural Research Station, Kampasagar, Nalgonda (Dist), Telangana State-508 207, India.

^b Institute of Rice Research, ARI, Rajendranagar, PJTSAU, Hyderabad-30, India.

^c Crop Improvement Section, ICAR-IIRR, Rajendranagar, Hyderabad-30. (GPBR), India.

^d Advanced Post Graduate Centre Lam Farm Opp. Regional Agricultural Research Station, Guntur, Andhra Pradesh-522034, India.

^e Agricultural College, Aswaraopet, Bhadradi Kothagudem (Dist) 507301, India.

^f PJTSAU, Rajendranagar, Hyderabad-30, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i121476

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93967>

Original Research Article

Received: 17/09/2022

Accepted: 23/11/2022

Published: 03/12/2022

ABSTRACT

The present investigation was carried out to understand the genetics and inheritance of yield components, physical, chemical and nutritional quality traits using generation mean analysis. In the present experiment, the fitting of digenic interaction model indicates the prevalence of interaction

⁺⁺ Senior Scientist (Pl.Br.) & Head;

[#] Principal Scientist (Rice), (Retired);

[†] National Professor;

[‡] Professor (GPBR);

[^] Associate Professor (GPBR);

^{###} Dean of Student Affairs (Retired);

*Corresponding author: E-mail: nlrashi80@gmail.com;

effects especially additive x additive [i] type in addition to the main additive and dominance gene effects in case of amylose content. The overall genetic analysis in the three crosses suggests a simple pedigree method to develop desirable lines from only cross viz., RP-Bio-5478-185 x NH-686 would be feasible. The generation mean analysis clearly indicated that apart from the additive and dominance gene effects, significant magnitudes [i] type were observed to play greater role in expression of iron content in rice. In comparison to the estimates of [d], the component due to dominance genetic effects [h] is very high in all the three crosses uniformly. However, due to prevalence of [h] component significantly in all the crosses, inter mating in early segregating generations, would further increase chances to develop iron enriched homozygous lines. Studies for zinc content revealed that among the interactions, [i] and [j] types were found to be significant, whereas the [l] component was totally negligible. From the study, all the three crosses were considered as superior on one or other way and among these three crosses, the last one (RP-Bio-5478-185 x NH-686) was considered as superior one. Simple pedigree method to develop desirable lines for protein and zinc, inter mating in early segregating generations for iron would be more feasible methods to develop nutrient rich desirable lines.

Keywords: Rice; generation mean analysis; digenic interaction model; χ^2 test; epistasis.

1. INTRODUCTION

Research efforts focused on development of high-yielding, disease & pest resistant and quality varieties and adoption of modern production technologies resulted in enhanced production leading to self-sufficiency in the country. Application of biometrical procedures in plant breeding in recent years led to a greater understanding of genetics of quantitative traits such as yield and proved to be useful to plant breeders for genetic analysis and planning for sound breeding programme. Along with yield, grain and nutritional quality has also become a primary consideration in rice breeding programs not only in India but also in various rice growing countries across the world. Grain quality characteristics are very important in rice breeding as it is predominantly consumed as a whole grain. Like yield, grain quality is not easily amenable to selection due to its complex nature. Lack of clear cut perception regarding the component traits of good quality rice is one of the important reasons for the tardy progress in breeding for rice varieties with fine grain quality. The quality characters of rice include physical attributes like grain appearance, kernel length/breadth ratio, milling parameters like hulling, milling percentages, head rice recovery etc. cooking quality and eating qualities like elongation ratio, gelatinization temperature. Now, nutritional elements like protein, iron and zinc contents were included as important quality attributes in view of their role in human health maintenance. Genetic variation for micronutrients in rice was studied and reported for iron and zinc [1] (Zhang et al., 2004). Iron and zinc contents in brown and milled rice of national and

international germplasm need to be estimated for the identification of donors for future deployment in the nutritional breeding programs. Incorporation of these characters into high yielding semi dwarf varieties are essential in this era of quality breeding.

The ability of parents to combine well depends upon complex interaction among genes, which cannot be predicted from yield performance and adaptability of the parents [2]. Most of the reports for gene action in rice are based on the diallel mating which does not provide information regarding non-allelic gene actions. The non-allelic gene actions could inflate the measures of additive and dominance components. Estimation of gene effects based on generation mean analysis provides requisite information to formulate the breeding strategy. Thus, generation mean analysis is a useful technique in plant breeding for estimating main gene effects (additive and dominance) and their digenic (additive x additive, additive x dominance, and dominance x dominance) interactions responsible for inheritance of quantitative traits. It helps us in understanding the performance of the parents used in crosses and potential of crosses to be used either for heterosis exploitation or pedigree selection. The nature of inheritance and type of gene action governing quality traits are many and complex and when genetically manipulated to some extent paved the way for the success in quality improvement through conventional breeding methods. Hence a better understanding of the factors that contribute to the overall grain quality of rice will lay the foundation for developing new breeding and selection strategies for combining high quality with high

yield. Genetic enhancement of key food crops with enhanced nutrients is advocated as the most promising approach to address the problem of malnutrition [3,4] which can be possible with understanding the genetic analysis among quality and nutritional traits in rice.

Considering the fact that nutrient traits are the most important complex traits and that their improvement is the most frequent goal of rice breeding programs. Keeping this aspect in view, the present investigation was undertaken to study the gene action governing yield, physico-chemical and nutritional traits in rice.

2. MATERIALS AND METHODS

The present investigation was carried out at the Regional Agricultural Research Station, Warangal, Telangana State, India which is located at an altitude of 304 M above MSL, 17.97° N latitude and 79.60° E longitude during two years 2014-15 and 2015-16 utilizing both *Kharif* and *Rabi* crop seasons in each year. The main crop seasons in Telangana State, India can be called as rainy (June-Dec) and post rainy (Nov-April) seasons.

2.1 Development of Material for Generation Mean Analysis

Five parents viz. MTU 1010, WGL-32100, RP-Bio-5478-185, NH-686 and RP-Bio-5478-166 were selected based on contrasting characters and developed material (F_1 , F_2 , BC_1 and BC_2) for 3 independent crosses (MTU 1010 X NH – 686, WGL-32100 X RP-Bio-5478-166 and RP-Bio-5478 -185 X NH-686) to study the presence of non allelic interactions through generation mean analysis for quality and nutritional characteristics. Variation for kernel dimensions (long slender x short bold, medium slender x short bold and short bold x short bold) and flowering duration was considered as criteria for selection of parents for generation mean analysis.

2.2 Generation of F_1 Hybrids

2.2.1 Hybridization & crop management techniques

During *Kharif* 2014, five parents were transplanted each in four rows in a crossing block at spacing of 20 x 15 cm and 4 sets were maintained. Crosses were effected to produce three F_1 s. The selected parents for study of generation mean analysis were sown at

staggered intervals of seven days to facilitate continuous availability of pollen during crossing. Twenty eight days old seedlings were transplanted at spacing of 30 x 15 cm. Hybridization was done by clipping method of emasculation as suggested by Jennings et al. (1981). Spikelets were emasculated in the afternoon and pollination was done in morning of next day between 11 – 1.00 P.M. Adequate care was taken to produce sufficient seed required for studying different generations.

2.3 Study of Generations

Three crosses (MTU 1010 X NH-686, WGL-32100 X RP-Bio-5478-166 and RP-Bio-7458-185 X NH-686) were selected wherein the parents had contrasting features for different traits under study. The F_1 's generated from five parents during *Kharif*, 2014, and corresponding F_2 's (seed obtained from F_1 generation raised during successive *Rabi*, 2014-15 and crosses were selfed and backcrossed with their respective parents to obtain backcrosses (BC_1 and BC_2) generations respectively. Parents were also selfed to ensure 100% genetic purity. Thus, seed of six basic generations, P_1 , P_2 , F_1 , F_2 , BC_1 and BC_2 was in hand for these three crosses at the end of the season *Rabi* 2014-15.

The experimental material (P_1 , P_2 , F_1 , F_2 , BC_1 , BC_2) was planted separately side by side for this study and data generated. The number of rows for P_1 , P_2 , F_1 , F_2 , BC_1 and BC_2 were 1, 1, 1, 8, 5, and 5 respectively. The mean data was used for final statistical analysis.

2.4 Estimation of Quality and Nutritional Traits

2.4.1 Amylose content (%)

Amylose (%) content in the rice sample was estimated as per the procedure given by Jennings et al. [5].

2.4.2 Crude protein content (%)

Protein content (%) = N content x 5.65

Nitrogen content of the rice grain was estimated by following micro-kjeldahl method [6].

2.4.3 Grain iron concentration zinc concentration (ppm)

Grain iron and zinc concentrations were determined by X-Ray fluorescence Spectrometry (XRF) (EDXRF, model-X-supreme8000).

2.5 Statistical Analysis

2.5.1 Generation means and gene effects

The concept of Generation Mean Analysis (GMA) was developed by Hayman and Mather [7] and Jinks and Jones [8] for the estimation of genetic components of variation (m, d, h, i, j, l). This technique involves six different generations viz., parents (P_1 and P_2), their F_1, F_2 and back crosses (BC_1 and BC_2). Accordingly, the means were computed for each generation of P_1, P_2, F_1, F_2, BC_1 and BC_2 for each cross over three replications in the present study. The variance and corresponding standard errors of the means were computed from the deviations of the individual values obtained from individual plants for each of the generation in each cross.

2.5.2 Joint Scaling test

Data were subjected to joint scaling test [9] in view of the advantages associated with this test. The Cavalli's joint scale test has two distinct advantages over scaling tests of Mather, 1949. Firstly, this test combines very effectively several scaling tests into one utilizing the entire available 6 generation mean and offers a more genetical approach. Secondly, this test provides best possible estimates of all the parameters needed to describe differences among the means of families, in addition to the test of adequacy of additive-dominance model. In the present study, the characters which fail to show significant variation among generations in any cross was not subjected to further genetic analysis of generation means. As only 6 generations are available in the event of significance of 6 parameters (m, d, h, i, j, l) and significance of χ^2 test also further analysis cannot be carried out and the result of χ^2 test cannot be interpreted due to non availability of degrees of freedom. Fortunately, in the present study, there were certain components which were non-significant, these were deleted and further analysis was carried out and χ^2 test was done using left over degrees of freedom.

Initially we tried a model with just 'm' first, if this adequately explained the variations in the trait, there was no need to proceed further to estimate any other genetical parameters, otherwise next higher parameters like d, h etc., were introduced until χ^2 test value become non significant [10].

First it was fitted into 3 parameter model, when χ^2 value was significant, it indicated that data

does not fit into additive - dominance model, hence, sequential model was followed. The data was first subjected to 6 parameter model, and when all the components were significant in six parameter model, no further analysis was carried out. In the event of non significance of any one of the components, the same was deleted and data fitted into 5 parameter model. This procedure was followed until the χ^2 value is non significant and all the components were significant in the model. Fitness in 5 or 4 parameter model indicated the presence of digenic non allelic interaction.

The following procedure was followed to estimate the components in the joint scaling test. The parameters $m, [d]$ and $[h]$ estimated from the observed mean of the available types of generations were compared with expected values derived from the estimates of these three parameters. The six equations which are obtained by equating the observed family means to their expectations in terms of $m, [d], [h], [i], [j]$ and $[l]$ were used for estimating these parameters. Since, the number of equations is higher than the number of parameters to be estimated, least square technique was followed.

2.6 Statistical Analyses

Means of three experimental values were used. Entire analyses were carried out by using Indostat available at Professor Jayashankar Telangana State Agricultural University, Hyderabad.

3. RESULTS AND DISCUSSION

The mean and variances of mean values of different generations utilized for generation mean analysis with respect to grain quality and nutritional traits of three crosses are presented in tabular form (Tables 1, 2 & 3).

Generation mean analysis was carried out using 6 generations means (P_1, P_2, F_1, F_2, B_1 & B_2). The adequacy of additive – dominance model was tested by using A, B, C, D scales of Mather [11] (Tables 4, 5 & 6).and also further confirmed by the Joint scaling test of Cavalli [9] (Table 7). As the Joint scaling test is a comprehensive test of simple additive – dominance model replacing A, B, C, D scaling test and involves weighted regression analysis, the components would be estimated and tested with more precision. Hence, a sequential method was followed to identify the best fit model in which the all the

possible components would be significant and at the same time the χ^2 test value becomes non significant. Major advantage of this test is that we can delete a non significant component in the sequential process and estimate the remaining significant components with maximum likelihood precisions.

Initially we tried a model with just 'm' first, if this adequately explained the variations in the trait, there was no need to proceed further to estimate any other genetical parameters, otherwise next higher parameters like d, h etc., were introduced until χ^2 test value become non significant [10].

In the present study, fortunately at least one component out of six was non significant, having at least one degrees of freedom to facilitate the χ^2 test. This procedure was adopted for three crosses uniformly and estimated the possible parameters by weighted least square method and discussed thoroughly (Table 7).

For amylase content, the fitting of digenic interaction model indicates the prevalence of interaction effects especially additive x additive [i] type in addition to the main additive and dominance gene effects. These findings are in agreement with the earlier reports of Mahalingam and Natarajan [12] and Subbalaxmi et al. [13]. In the present study, both additive and dominance genetic effects were found to be significant. In addition, the [i] type of epistasis was also found to play a greater role in expression of amylase content in desirable direction. A duplicate type of epistasis was registered in the second cross (WGL-32100 x RP-Bio-5478-166) as the signs were in opposite sides. Keeping in view the highly significant estimates of fixable, additive x additive type of genetic variation coupled with existence of higher magnitudes of additive gene effects, a better scope is expected for isolation of promising genotypes through simple selection procedures. However, attempting few crosses among the selected segregating genotypes and postponing the selections for some time may further improve the breeding efficiency [14].

For protein content, significant values of 'm' were estimated in the crosses studied. In two crosses (MTU 1010 x NH-686 and WGL-32100 x RP-Bio-5478-166) the interaction effect was observed which was of duplicate type as the [h] and [I] components showed opposite signs. The genetic variation was largely controlled by dominance gene effects of negative type in these two crosses, thus a little scope was indicated for

improvement of protein content for these two crosses. In view of the dominance gene effects on negative side and lowering of desirable interaction effects due to mutual cancellation, heterosis breeding would not be profitable. The overall genetic analysis in the three crosses suggests a simple pedigree method to develop desirable lines from only cross viz., RP-Bio-5478-185 x NH-686.

Iron deficiency is probably the most wide spread micro nutrient deficiency in rural areas, where rice is the staple food. Therefore, apart from breeding, the approaches like genetic engineering, biochemical and physical have been frequently used as prospective methods to regulate iron content and bio availability in rice grains. This present study was envisaged to understand the genetics of iron content as little efforts were made so far in these lines. The generation mean analysis clearly indicated that apart from the additive and dominance gene effects, significant magnitudes of digenic interactions especially [i] type were observed to play greater role in expression of iron content in rice. In one cross, RP-Bio-5478-185 x NH-686, all the types of digenic interactions were present with duplicate dominant epistasis. In comparison to the estimates of [d], the component due to dominance genetic effects [h] is very high in all the three crosses uniformly. It is interesting to note that [i] type of epistasis which is fixable in nature is highly significant giving a way for direct selection. However, due to prevalence of [h] component significantly in all the crosses, inter mating in early segregating generations, would be further useful to develop iron enriched homozygous lines.

Apart from iron, zinc is another important element to be consumed at optimum level as per the dietary requirements. Therefore, in most of the programmes both iron and zinc would be considered for simultaneous improvement. To understand the genetic architecture of zinc content in rice, the generations mean analysis was carried out and fitted in various models. The first cross combination was fitted in digenic interaction 4 parameter model, whereas, other two crosses were fitted at 5 parameter level. Among the interactions, [i] and [j] types were found to be significant, whereas the [I] component was totally negligible. When fixable components are taken into consideration, the [i] type (additive x additive) was observed to be more predominant in desirable side in all the three crosses, whereas [d] type was significant

Table 1. Estimates of generation mean and variances of mean for nutrient characters for the cross MTU 1010 x NH-686

Character	MTU 1010 x NH-686											
	P ₁		P ₂		F ₁		F ₂		BC ₁		BC ₂	
	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean
Amylose content (%)	23.02 ±0.23	0.05	20.94 ±0.33	0.10	24.89 ±0.22	0.05	20.76 ±0.18	0.03	22.68 ±0.44	0.19	22.06 ±0.86	0.74
Protein content (%)	7.40 ±0.22	0.04	7.54 ±0.16	0.02	9.64 ±0.21	0.04	7.66 ±0.47	0.22	7.51 ±0.17	0.03	7.76 ±0.23	0.05
Iron content (ppm)	9.80 ±0.02	0.007	10.03 ±0.12	0.01	10.02 ±0.24	0.05	9.17 ±0.07	0.005	9.46 ±0.07	0.006	9.79 ±0.08	0.007
Zinc content (ppm)	11.11 ±0.06	0.004	13.76 ±0.18	0.03	13.00 ±0.16	0.02	10.78 ±0.69	0.48	11.26 ±0.42	0.17	13.00 ±0.29	0.08

Table 2. Estimates of generation mean and variances of mean for nutrient characters for the cross WGL-32100 X RP-Bio-5478-166

Character	WGL-32100 X RP-Bio-5478-166											
	P ₁		P ₂		F ₁		F ₂		BC ₁		BC ₂	
	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean
Amylose content (%)	22.68 ±0.27	0.07	19.68 ±0.66	0.43	22.77 ±0.16	0.02	21.19 ±1.41	1.98	21.51 ±0.36	0.13	19.53 ±0.61	0.37
Protein content (%)	7.11 ±0.06	0.004	8.53 ±0.26	0.07	8.88 ±0.06	0.004	7.98 ±0.54	0.29	10.82 ±0.37	0.01	8.12 ±0.27	0.07
Iron content (ppm)	10.75 ±0.41	0.17	12.89 ±0.20	0.04	11.99 ±0.21	0.04	9.91 ±0.17	0.03	14.30 ±0.39	0.14	10.78 ±0.37	0.14
Zinc content (ppm)	14.44 ±0.11	0.01	19.69 ±0.16	0.02	22.69 ±0.22	0.05	14.15 ±0.28	0.08	14.30 ±0.39	0.15	20.81 ±0.47	0.22

Table 3. Estimates of generation mean and variances of mean for nutrient characters for the cross RP-Bio-5478 -185 x NH-686

Character	RP-Bio-5478 -185 x NH-686											
	P ₁		P ₂		F ₁		F ₂		BC ₁		BC ₂	
	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean	Mean	Variance of mean
Amylose content (%)	19.36 ±0.40	0.16	21.36 ±0.25	0.06	21.01 ±0.45	0.20	19.29 ±0.30	0.09	19.66 ±0.26	0.07	21.13 ±0.44	0.19
Protein content (%)	7.84 ±0.16	0.02	7.56 ±0.15	0.02	8.36 ±0.20	0.04	7.80 ±0.42	0.18	7.97 ±0.07	0.005	7.79 ±0.13	0.01
Iron content (ppm)	11.64 ±0.23	0.05	9.85 ±0.20	0.04	15.32 ±0.35	0.12	10.15 ±0.52	0.27	10.70 ±0.28	0.07	14.01 ±0.24	0.06
Zinc content (ppm)	21.31 ±0.34	0.12	13.78 ±0.22	0.05	19.15 ±0.28	0.08	14.44 ±0.25	0.06	14.44 ±0.26	0.07	18.34 ±0.37	0.14

only in one cross i.e RP-Bio-5478-185 x NH-686, hence all the three crosses were considered as superior on one or other way. This findings is in agreement with the earlier report of Maddeppa Mallimar et al., [15]. The dominance effects were prevalent in desirable direction in all the crosses where as the additive component was significantly negative in first and second

crosses. In view of the predominant role of dominance effects and also the fixable interaction effects, the breeding methods like recurrent selection, biparental mating would be highly advantageous and among these three crosses, the last one (RP-Bio-5478-185 x NH-686) was considered as superior one [16].

Table 4. Caling tests of generation means for the cross MTU-1010 x NH-686 for nutrient traits

S. No	Character	MTU-1010 x NH-686			
		Scales			
		A	B	C	D
1	Amylose content (%)	-2.55* ± 0.94	-1.70 NS ± 1.76	-10.72** ± 0.94	-3.23** ± 1.03
2	Protein content (%)	-2.01** ± 0.46	-1.65 **± 0.54	-3.57 NS ± 1.98	0.04 NS ± 1.00
3	Iron content (ppm)	-0.90** ± 0.29	-0.47NS ± 0.32	-3.17** ± 0.57	-0.89** ± 0.18
4	Zinc content (ppm)	-1.51 NS ± 0.86	-0.75 NS ± 0.63	-7.75** ± 2.81	-2.74 NS ± 1.48

*Significant at 5 % level, ** Significant at 1 % level

Table 5. Scaling tests of generation means for the cross WGL-32100 X RP-Bio-5478-166 for nutrient traits

S. No	Character	WGL-32100 X RP-Bio-5478-166			
		Scales			
		A	B	C	D
1	Amylose content (%)	-2.42** ± 0.79	-3.38** ± 1.40	-3.14 NS ± 5.69	1.33 NS ± 2.90
2	Protein content (%)	-1.80** ± 0.27	-1.17 NS ± 0.62	-1.50 NS ± 2.19	0.73 NS ± 1.12
3	Iron content (ppm)	-1.10 NS ± 0.88	-3.32** ± 0.81	-7.97** ± 0.93	-1.77* ± 0.63
4	Zinc content (ppm)	-8.53** ± 0.82	-0.75 NS ± 0.98	-22.89** ± 1.25	-6.80** ± 0.84

*Significant at 5 % level, ** Significant at 1 % level

Table 6. Scaling tests of generation means for the cross RP-Bio-5478 -185 x NH-686 for nutrient traits

S. No	Character	RP-Bio-5478 -185 x NH-686			
		Scales			
		A	B	C	D
1	Amylose content (%)	-1.04NS ± 0.81	-0.10NS ± 1.02	-5.58** ± 1.60	-2.22** ± 0.80
2	Protein content (%)	-0.25NS ± 0.30	-0.34NS ± 0.37	-0.91NS ± 1.76	-0.15NS ± 0.86
3	Iron content (ppm)	-5.56** ± 0.70	2.85** ± 0.63	-11.52** ± 2.24	-4.40** ± 1.11
4	Zinc content (ppm)	-11.57** ± 0.70	3.75** ± 0.83	-15.61** ± 1.22	-3.89** ± 0.68

*Significant at 5 % level, ** Significant at 1 % level

Table 7. Genetic components of nutrient traits estimated through Joint scale test

Cross	Genetic components	Character			
		Amylose content (%)	Protein content (%)	Iron content (ppm)	Zinc content (ppm)
MTU 1010 x NH-686	m	16.64** ± 0.42	7.49** ± 0.13	8.29** ± 0.26	9.77** ± 0.84
	[d]	1.03** ± 0.19	-	-0.17** ± 0.05	-1.35** ± 0.09
	[h]	8.26** ± 0.57	-1.68* ± 0.69	1.79** ± 0.43	3.24** ± 0.91
	[i]	5.35** ± 0.47	-	1.67** ± 0.26	2.68** ± 0.86
	[j]	-	-	-	-
	[l]	-	3.84** ± 0.73	-	-
	χ^2 value/probability		0.327(0.849) NS	1.026(0.795) NS	2.763(0.251) NS
Model fitted		4 parameter	3 parameter duplicate	4 parameter	4 parameter
WGL-32100 x RP-Bio-5478-166	m	21.12** ± 0.34	7.87** ± 0.12	7.80** ± 0.40	5.94** ± 0.58
	[d]	1.59** ± 0.32	-0.76** ± 0.12	-0.90** ± 0.21	-2.62** ± 0.09
	[h]	-3.54* ± 1.49	-2.31** ± 0.53	4.17** ± 0.54	16.80** ± 0.71
	[i]	-	-	4.10** ± 0.46	11.13** ± 0.59
	[j]	-	-	-	-7.45 ** ± 1.23
	[l]	5.19** ± 1.36	3.32** ± 0.50	-	-
	χ^2 value/probability		0.519/ (0.771) NS	1.542/ (0.463) NS	3.849/ (0.146) NS

Cross	Genetic components	Character			
MTU 1010 x NH-686		Amylose content (%)	Protein content (%)	Iron content (ppm)	Zinc content (ppm)
Model fitted		4 parameter duplicate	4 parameter duplicate	4 parameter	5 parameter
Type of epistasis		-	-	-	-
RP-Bio-5478 -185 x NH-686	m	17.17** ± 0.75	7.66** ± 0.10	-	9.73** ± 0.55
	[d]	-1.02** ± 0.21	0.57** ± 0.20	0.89** ± 0.15	3.76 ** ± 0.20
	[h]	3.48** ± 1.09	-	23.58** ± 0.80	9.41** ± 0.75
	[i]	2.67** ± 0.79	-	10.73** ± 0.15	7.81** ± 0.36
	[j]	-	-	-8.38* ± 0.80	-15.33** ± 0.98
	[l]	-	-	-8.26** ± 1.00	-
χ ² value/probability		1.815/(0.404)	3.548/(0.471) NS	0.744/ (0.388)	0.000/ (0.989) NS
Model fitted		4 parameter	2 parameter	5 parameter duplicate	5 parameter duplicate
Type of epistasis		-	-	-	-




Fig. 1. Mean values of 6 Generations of 3 crosses for Iron content

C1: MTU 1010 x NH-686, C2: WGL – 3200 x RP-Bio-5478-166, C3: RP-Bio-5478 -185 x NH – 686




Fig. 2. Mean values of 6 generations of 3 crosses for zinc content

C1: MTU 1010 x NH-686, C2: WGL – 3200 x RP-Bio-5478-166, C3: RP-Bio-5478 -185 x NH – 686

4. CONCLUSION

The fitting of digenic interaction model indicates the prevalence of interaction effects especially additive x additive [i] type in addition to the main additive and dominance gene effects in case of amylase content. A duplicate type of epistasis was registered in the second cross (WGL-32100 x RP-Bio-5478-166) as the signs were in opposite sides. Simple pedigree method to develop desirable lines for protein and zinc, inter mating in early segregating generations for iron would be more feasible methods to develop nutrient rich desirable lines.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Gregorio GB, Senadhira D, Htut H, Graham RD. Breeding for trace mineral density in rice. Food Nutr. Bull. 2000; 21:382-386.
- Allard RW. Principles of plant breeding. John Wiley and Sons Inc., New York, USA. 1960;485.
- Graham RD, Welch RM, Bouis HF. Addressing micronutrient malnutrition through enhancing the nutritional quality of staple foods: principles, perspectives and knowledge gaps. Adv. Agron. 2001;70:77-142.
- Bouis HE. Plant breeding: A new tool for fighting micronutrient malnutrition. J Nutrition. 2002;132:491-494.
- Jennings PR, Coffman WR, Kaufman MHE, Grain quality: Rice improvement. International Rice Research institute, Philippines. 1979;Chapter 6:101-120.
- A.O.A.C. Official methods of analysis, 13th ed. Association of official analytical chemists. Washington, D.C. 1980;376-384.
- Hayman B.I, Mather K. The description of genic interactions in continuous variation. Biometrics. 1955;11:69-82.
- Jinks JL, Jones RM.. Estimation of the components of heterosis. Genetics. 1958;43:223-234.
- Cavalli LL. .An analysis of linkage in quantitative inheritance. Quantitative Inheritance. H.M.S.O, London. 1952;135–144.
- Kearsey MJ, Pooni HS. The genetical analysis of quantitative traits. Chapman and Hall,. London; 1996.
- Mather K . Biometrical Genetics. Methuen and Co. Ltd., London; 1949.
- Mahalingam L, Nadarajan., N. Genetic analysis of grain quality characteristics of two line rice hybrids. Electronic Journal of Plant Breeding. 2010;1:983-988.
- Subbalakshmi K., Shunmugavalli N, Muthuswamy A. Generation mean analysis for yield and quality traits in F₂ and F₃ generation of rice (*Oryza sativa* L.). Electronic Journal of Plant Breeding. 2016; 7:491-495.
- Ramli, AB, Rafii MY, Latif MA, Saleh GB. Omar, G. B. Generation mean analysis of grainquality traits in selected rice populations derived from different amylose characteristics. Journal of the Science of Food and Agriculture. 2016;96:1593-1600.
- Maddeppa Mallimar P, Surendra Babagouda Patil, Satish TN, Mahantesh Jogi, Ramaling Hundekar. Study The Inheritance of Iron and Zinc in Segregating Population of Rice (*Oryza sativa* L.). Int. J. Pure App. Biosci. 2017;5:888-892.
- Zhang MW, Guo BJ, Peng ZM. Genetic effects of Fe, Zn, Mn and P content in indica black pericarp rice and their genetic correlations with grain characteristics. Euphytica. 2014;135:315-323.

© 2022 Lingaiah et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93967>