



# Integrated Management of Root-Knot Nematode (*Meloidogyne* Spp.) Using Fresh Organic Manure and *Crotalaria brevidens* Intercrop for Improved Growth and Yield of Tomato

James Ngelenzi Munywoki <sup>a\*</sup>, Joshua Otieno Ogweno <sup>a</sup>  
and Mwanarusi Saidi <sup>a</sup>

<sup>a</sup> Department of Crops, Horticulture and Soils, Egerton University, P.O.Box 536-20115, Egerton, Kenya.

## Authors' contributions

This work was carried out in collaboration among all authors. Author JNM was responsible for performing the experiment, data collection, statistical data analysis, interpretation of the results and manuscript preparation. Authors JOO and MS conceived the idea, designed the experiment and edited the manuscript. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2022/v34i430859

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/78865>

Original Research Article

Received 20 October 2021  
Accepted 25 December 2021  
Published 21 February 2022

## ABSTRACT

Tomato (*Solanum lycopersicum* L.) is one of the most important vegetables worldwide. Tomato productivity is limited by, among other factors, biotic constraints including root-knot nematodes (*Meloidogyne* species) which lower both quality and quantity of the crop. Two trials were conducted at the Horticulture Research and Teaching Field, Egerton University, Kenya, to evaluate effects of fresh organic manure and slender leaf (*Crotalaria brevidens* var. *brevidens* Benth.) intercrop on management of root-knot nematode during tomato production as well as establish their effect on tomato growth and yield. Variables measured were nematode infestation, plant height, stem collar diameter, number of internodes and branches, number of fruits, and fresh fruit weight. Use of fresh organic manure and slender leaf intercrop suppressed the nematode population by 27.8% to 53.5% reduction and significantly increased tomato fruit numbers by 22.2% to 49.7% and fresh fruit weight by 24.5% to 80.4% when used alone or in combinations compared to the control treatment. The combination of fresh goat dung and

\*Corresponding author: E-mail: [jamezngelenzi@gmail.com](mailto:jamezngelenzi@gmail.com);

slender leaf intercrop resulted in the highest reduction in nematode population of 53.5% and highest increase in total number of fruits of 49.7% and fresh fruit weight of 80.4%.

**Keywords:** *Solanum lycopersicum*; galls; goat manure; cow manure; slender leaf.

## 1. INTRODUCTION

Tomato (*Solanum lycopersicum* L.) is one of the most important vegetables worldwide. Tomato productivity is limited by, among other factors, biotic constraints including root-knot nematodes (*Meloidogyne* spp.) which reduce both the quality and quantity of tomato. Root-knot nematodes can cause up to 80% yield losses especially in heavily infested tomato growing fields [1]. The short life cycle of six to eight weeks enables root-knot nematode populations to survive and proliferate in the presence of a suitable host and their numbers will build up to a maximum as the crop reaches maturity [2,3]. Control of root-knot nematodes is difficult since the pathogen is soil-borne with a wide host range and usually attacks the underground parts of the plant [4].

Currently, the primary control measures employed against root-knot nematodes are crop rotation and chemical nematicides as a pre-plant fumigation of soil. Although crop rotation can be an effective method of managing root-knot nematode, it is not feasible among smallholder crop production systems due to limited land resource. Chemical nematicides, on the other hand, are effective and fast-acting but are currently being reappraised with respect to their effects on the environment and human health as well as the high cost of purchase [5]. There is, therefore, need for development of alternative sustainable management strategies for the pest that are ecofriendly and relatively affordable to smallholder tomato growers.

Alternative strategies for management of root-knot nematodes such as heat treatment, soil solarization, crop rotation with non-host plants, intercropping, and application of organic soil amendments of crop and/or animal origin [6] that do not pollute the environment have been emphasized to researchers, farmers, and scientists [7]. Studies have shown that host plants can be protected against root-knot nematode infection through intercropping practices especially with plants with nematicidal or nematostatic properties [3]. *Crotalaria* has been documented as a poor host to many plant-parasitic nematodes including *Meloidogyne* species. *Crotalaria* increases bacterivorous

nematode population densities and nematode-trapping fungal propagules, thus enhancing microbial activities against *Meloidogyne* species [8]. Besides pest suppression or control, other associated benefits of intercropping other crops with *Crotalaria* are greater yield stability, greater land-use efficiency, increased competitive ability toward weeds, and improvement of soil fertility [9].

Use of organic manure reduces negative effects of root-knot nematode on host crop with a resultant increase in growth and yield of the crop [10]. An organic manure amendment improves activities of non-pathogenic organisms and directly suppresses soilborne pathogens during decomposition [11]. Predatory nematodes numbers tend to be higher in organic amended soil thus suppressing parasitic nematodes in the soil [12]. Well-decomposed composts are stable and mineralize slowly releasing low concentrations of toxic nematicidal compounds [13]. Nahar et al. [14] proposed that raw organic manure may be more effective than composted manure as toxic compounds could build up quickly to reach toxicity threshold level for nematodes and other soilborne pathogens.

Integrated Crop Management (ICM) strategies have been developed worldwide for many crops and have resulted in reduced pesticide use, higher crop yields and economic value, as well as reduced economic risks for farm management owing to lower variation in the severity of pest problems [15]. Further improvement of such programs will promote their widespread applicability and improve on performance of crop production systems. This study sought to investigate the effects of integrating fresh organic manure (cow and goat dung) and slender leaf (*Crotalaria brevidens* var. *brevidens* Benth.) intercrop as an alternative management strategy for suppressing root-knot nematode during tomato production, as well as establish the effect of the strategy on growth and fruit yield of the crop.

## 2. MATERIALS AND METHODS

Two trials (July to October 2019 and January to April 2020) were conducted at the Horticulture


Research and Teaching Field of Egerton University, Njoro, Kenya. The field lies at a latitude of 0°23' S and longitude 35°35' E in the Lower Highland III Agro Ecological Zone (LH3), at an altitude of ≈2238 m above sea level. The soils are predominantly vitric mollic andosols with a pH of 6.0 to 6.5 [16]. The monthly mean temperature and monthly rainfall during the study period are presented in Fig. 1.

Tomato cultivar 'Rio Grande' and slender leaf (*Crotalaria brevidens* var. *brevidens* Benth.) seeds were obtained from Kenya Seed Company in Nakuru - Kenya. 'Rio Grande' is a determinate tomato cultivar with a high yielding potential and one of the most popular among open field tomato growers in Kenya. Slender leaf is an annual or short-lived perennial herb in the family Fabaceae. The genus is common in the tropics and subtropics, with the greatest number of species occurring in Africa.

The experimental design was a 2 x 3 factorial arranged in a randomized complete block design (RCBD) with three replications. The two factors were cropping regime at two levels (tomato intercropped with slender leaf and monocrop tomato) and fresh organic manure at three levels (cow dung, goat dung, and no organic manure),

for a total of six treatment combinations; (i) fresh cow dung with no slender leaf; (ii) fresh goat dung with no slender leaf; (iii) fresh cow dung and slender leaf; (iv) fresh goat dung and slender leaf; (v) slender leaf with no organic manure; (vi) control (monocrop tomato with no organic manure). Each experimental unit consisted of 5 troughs each 2 m long, 0.4 m wide and 0.2 m high in a 2 x 2 m plot separated from each other by 1 m path. Each block consisted of 6 such arrangements to accommodate the six treatment combinations. Individual blocks were 17 m long separated from each other by 1.5 m path. Soil for filling the troughs was sterilized by covering with airtight clear polyethylene sheets for a period of eight weeks. Troughs filled with sterilized soil were placed on a clear polyethylene sheet to prevent contaminating the field with nematodes.

Tomato seeds were sown in rows spaced 20 cm apart in a raised nursery bed and transplanted into troughs filled with soil in the field when the seedlings were 5 weeks old. Diammonium phosphate (DAP) fertilizer was applied at the rate of 240 kg·ha<sup>-1</sup> [17] and thoroughly mixed with the soil prior to transplanting. Tomato seedlings were transplanted at 60 cm apart within the trough giving a population density of 4 plants/trough and



**Fig. 1. Average monthly air temperature (°C) and precipitation (mm) during tomato production over the two trials (July to October 2019 and January to April 2020)**

Source: Egerton University Engineering Department (2020)

a total of 20 plants per experimental unit. In intercropped plots, slender leaf seedling rows were made across the trough at 60 cm between adjacent rows of slender leaf and 30 cm from individual tomato rows giving a total of three rows of slender leaf per trough. Slender leaf seeds were drilled at the same time with the sowing of tomato seeds in the nursery so as to have them well established by the time of transplanting tomato seedlings. Thinning of the slender leaf seedlings was done when plants were about 10 cm tall to a spacing of 10 cm between plants giving a total of four plants per row and twelve plants per trough. Tomato plants were top dressed with Calcium Ammonium Nitrate at the rate of 240 kg·ha<sup>-1</sup> [17] applied in two splits; first split at three weeks after transplanting and the second split three weeks later.

Two weeks after tomato transplanting, second instar juvenile stage of *Meloidogyne* species inoculum suspension of approximately 250 juveniles per plant was sprinkled on each experimental unit. Thereafter, fresh organic manure at a rate of 15 ton·ha<sup>-1</sup> was also applied in furrows made adjacent to tomato plants in individual troughs as per the designated treatments. Fresh cow and goat dung were obtained from the Tatton Agriculture Park (TAP) of Egerton University.

Nematode infestation was determined by evaluating nematode population in the soil and gall assessment on tomato roots at the end of the experiment. To determine the nematode population in the soil, second stage juveniles (J2) were extracted from 100 cm<sup>3</sup> soil from each experimental unit, using the method described by Coyne et al. [18]. The soil samples were placed in sieves lined with double layered tissue paper. The sieves were half immersed in plastic beakers containing 250 ml of distilled water for 24 hours to allow nematode migration into the water underneath. Quantification of juveniles was done under a light microscope with gridded petri dishes. Three 1 ml replicate samples were drawn from the well mixed suspension to establish the average number of juveniles per millimeter and later determine the number of nematodes per 100 cm<sup>3</sup> soil of each experimental unit, and subsequently the average number of nematodes per treatment combination. For gall assessment, plants were gently uprooted, and their roots washed with tap water to remove the adhering soil. Galling was determined by counting the number of galls and data obtained recorded as number of galls per plant (no./plant).

Four plants from the inner rows of each experimental unit were randomly selected and tagged for collection of data on plant height, stem collar diameter, number of internodes, number of branches and yield. Plant growth data were collected on a two-week interval beginning at the second week after application of treatments and ending at first harvest. Plant height of the tagged plants was measured in centimeters (cm) from the ground level to the point of growth using a meter ruler and the number of internodes and branches were counted and recorded as number of internodes or branches per plant (no./plant). At the same time, stem collar diameter of the four tagged tomato plants was measured at  $\approx$  2 cm from the ground level using a digital vernier caliper (Model 599-577-1/USA) and the data obtained was used to compute the average stem collar diameter of plants for the different treatments in millimeters (mm).

Tomato fruits from each experimental unit were harvested twice every week at breaker stage. At each harvest, tomato fruits from each tagged plant were physically counted and weighed in kilograms (kg/plant) using a weighing balance (ATZ; Shanghai Precision and Scientific Instrument Co., Shanghai, China). Data obtained were later used to compute the average number of fruits per plant (no./plant) and weight of fruits per plant (kg/plant) for the different treatments.

The Proc univariate procedure of SAS (ver. 9.1; SAS Institute, Cary, NC) was used to determine normality and equal variances assumptions of analysis of variance of the data before analysis. Where assumptions were not met, appropriate transformation was done. Data were subjected to ANOVA using the GLM procedure of SAS at  $p \leq 0.05$ . Initial analysis showed no significant interaction between treatments and trial for most variables measured. Data for the two trials were thus pooled together and analyzed using the statistical model RCB model:  $Y_{ijk} = \mu + \alpha_i + \beta_j + \tau_k + \beta\tau_{jk} + \epsilon_{ijk}$ , where  $Y_{ijk}$  is the tomato response,  $\mu$  is the overall mean,  $\alpha_i$  is the  $i^{\text{th}}$  blocking effect,  $\beta_j$  is the effect due to  $j^{\text{th}}$  organic manure,  $\tau_k$  is the effect due to  $k^{\text{th}}$  cropping regime,  $\beta\tau_{jk}$  is interaction effect of the  $j^{\text{th}}$  organic manure and  $k^{\text{th}}$  cropping regime,  $\epsilon_{ijk}$  is the random error term. Means for significant treatments, at the F test, were separated using Tukey's honestly significant difference test at  $p \leq 0.05$ .

### 3. RESULTS

#### 3.1 Effects of Slender Leaf Intercrop and Fresh Organic Manure on Nematode Infestation on Tomato Growth and Productivity

Use of fresh organic manure and slender leaf intercrop either alone or in combination reduced nematode population in the soil by 27.8–53.5% in comparison with the control treatment (Table 1). Plots with fresh goat dung and slender leaf intercrop recorded the lowest juvenile root-knot nematode population in the soil and had the highest reduction of 53.5% of juvenile root-knot nematode population in the soil compared to the control treatment. Plots where tomato was intercropped with slender leaf with no fresh manure applied had higher numbers of juvenile root-knot nematodes in the soil among intercropped and fresh manure treated plots but had lower numbers compared to the control with a 27.8% reduction in the population of juvenile root-knot nematodes in the soil. Among the other treatments, plots applied with fresh cow dung and slender leaf intercrop had slightly lower soil nematode population followed by those applied with fresh goat dung and fresh cow dung in that order. When comparing the main effects of growing tomato with or without slender leaf intercrop, intercropping with slender leaf significantly reduced the number of juvenile root-knot nematode populations by 24.4% compared to when tomato was grown as a monocrop (Table 1). Similarly, growing tomato with fresh cow or goat dung significantly reduced juvenile root-knot nematode population by 32.9% and 37.7%, respectively, compared to when no fresh organic manure was applied (Table 1).

Similar to the trends observed for juvenile root-knot nematode population in the soil, the number of galls on tomato roots was consistently lower on tomato plants grown with fresh organic manure and slender leaf intercrop. The use of fresh organic manure and slender leaf intercrop reduced number of galls in the tomato roots by 58.4–65.4% in comparison with the control treatment. Overall, the lowest number of root galls was recorded in plots with fresh goat dung and slender leaf intercrop while the highest number was with the control treatment (Table 1). Intermediate number of root galls was obtained for the other treatments but the differences in the number of galls among all intercrop and fresh manure treatments were not statistically significant. A comparison of the two cropping

regimes showed that intercropping tomato with slender leaf reduced the number of root-knot nematode galls by 36.8% compared to when tomato was grown as a monocrop with the reduction being statistically significant. Similarly, growing tomato with fresh organic manure significantly reduced the number of root-knot nematode galls on tomato roots by 44.8% and 48.7% for fresh cow and fresh goat manure, respectively, compared to when no fresh organic manure was applied (Table 1).

#### 3.2 Effects of Slender Leaf Intercrop and Fresh Organic Manure on Tomato Plant Growth

Growing tomato under combined application of fresh organic manure and slender leaf intercrop did not significantly influence tomato plant height but resulted in relatively better growth than when the two technologies were used separately or not used at all (Table 2). In most sampling dates, tomato plants tended to be tallest in plots with fresh goat dung treatment and shortest in the control treatment. During most sampling dates, plant height also tended to be higher in plots with slender leaf intercrop and fresh organic manure (goat or cow dung) compared to plots with either slender leaf intercrop or cow dung alone. By the final sampling date (at 70 days after transplanting or DAT), tomato plants were tallest in plots treated with goat dung and shortest in the control treatment. Among treatments, tomato plants grown with fresh cow dung and slender leaf intercrop tended to be taller followed by those in plots with goat dung and slender leaf intercrop, then those grown with fresh cow dung, and the shortest plants were those grown with slender leaf intercrop with no fresh organic manure. When comparing the two cropping regimes, tomato intercropped with slender leaf had taller plants compared to monocrop, although the difference in height was not statistically significant (Fig. 2A). Similarly, tomato plants grown with fresh organic manure were significantly taller than those grown without application of fresh organic manure during all sampling dates except at 28 DAT when the difference was not statistically significant (Fig. 2A).

Tomato collar diameter was also significantly affected by the use of fresh organic manure and slender leaf intercrop during production (Table 2). The lowest collar diameter amongst all treatments was observed in the control plants. Growing tomato plants under combined use of

fresh goat or cow manure with slender leaf intercrop resulted in thicker stems compared to the other treatments in most sampling dates. Tomato plants intercropped with slender leaf had slightly thicker stems compared to the monocrop tomato plants (Fig. 2B). Likewise, growing tomato with fresh organic manure generally resulted in significantly thicker plants compared to when no fresh organic manure was applied in all sampling dates (Fig. 2B).

The number of tomato stem internode was also influenced by the use of fresh organic manure and slender leaf intercrop during production (Table 2). Using the two technologies in combination resulted in higher number of internodes than where they were not used at all. Internode numbers also tended to be slightly higher for monocrop tomato grown with fresh cow or goat dung compared to the internode numbers recorded for control plants. Comparing the main effects, tomato plants intercropped with slender leaf had slightly higher number of internodes compared to tomato plants grown as a monocrop (Fig. 2C). Tomato grown with fresh organic manure also had higher number of

internodes compared to when no fresh organic manure was applied (Fig. 2C).

Similarly, the use of fresh organic manure and slender leaf intercrop significantly improved plant branching ability (Table 2). The number of branches per plant was higher for plants grown with fresh organic manure and slender leaf intercrop compared to plants grown where the two technologies were each used in isolation or were not used at all in most sampling dates. In most sampling dates, plants grown with cow dung and slender leaf intercrop had the highest number of branches while the least number of branches per plant was recorded on control treatment plants. Comparing the two cropping regime, tomato intercropped with slender leaf had more branches compared to tomato grown as a monocrop, although the difference was not statistically significant except during the final data collection date at 70 DAT (Fig. 2D). Similarly, tomato grown with fresh organic manure had significantly more branches compared to those grown without fresh organic manure at all data collection dates (Fig. 2D).

**Table 1. Effects of slender leaf intercrop and fresh organic manure on root-knot nematode juvenile population in the soil and on the number of galls of 'Rio Grande' tomato**

Manure	Cropping regime				Manure Means	Reduction (%)
	Intercrop	Reduction (%)	Monocrop	Reduction (%)		
	Number of juveniles/100 g soil					
Cow	31.5b*	48.7**	39.4b	35.8	35.5b	32.9
Goat	28.6b	53.5	37.3b	39.2	33.0b	37.7
No manure	44.3ab	27.8	61.4a		52.9a	
Cropping regime Means	34.8b		46.1a			
Reduction (%)	24.4					
	Number of root galls/plant					
Cow	120.1b	62.9	133.4b	58.8	126.8b	44.8
Goat	112.0b	65.4	123.3b	62.0	117.6b	48.7
No manure	134.9b	58.4	324.0a		229.5a	
Cropping regime Means	122.4b		193.6a			
Reduction (%)	36.8					

\*Means followed by the same letter within a letter series and within a variable are not significantly different according to Tukey's honestly significant difference test at  $p \leq 0.05$ .

\*\*Percent reduction computed in reference to nematode (juvenile) population or galls in the control treatment (No slender leaf (monocrop), no fresh manure).

**Table 2. Effects of slender leaf intercrop and fresh organic manure on plant growth during tomato production**

Treatment		Days after Transplanting			
Fresh manure		28	42	56	70
Cropping regime					
Plant height (cm)					
Intercrop	Cow	31.0*	47.4	64.3	73.5
	Goat	32.2	45.5	61.0	72.5
	No manure	32.4	44.6	57.7	66.8
Monocrop	Cow	29.2	46.1	63.6	71.5
	Goat	31.9	47.9	64.3	74.7
	No manure	30.5	41.7	55.5	64.6
Collar diameter (mm)					
Intercrop	Cow	6.8 ab**	9.1 ab	11.3 a	12.5 ab
	Goat	7.5 a	9.5 a	11.2 a	12.7 a
	No manure	6.5 b	9.0 ab	10.2 ab	11.5 ab
Monocrop	Cow	6.4 b	9.5 a	11.0 ab	12.3 ab
	Goat	6.6 b	9.2 ab	11.0 ab	12.4 ab
	No manure	6.2 b	7.7 b	9.3 b	10.6 b
Number of internodes (no./plant)					
Intercrop	Cow	8.0	11.0	12.9 a	13.7
	Goat	8.3	10.9	12.3 ab	13.5
	No manure	8.0	10.6	12.2 ab	13.7
Monocrop	Cow	8.0	10.4	12.6 a	13.4
	Goat	7.9	10.7	12.5 a	13.5
	No manure	7.8	9.8	11.2 b	12.5
Number of branches (no./plant)					
Intercrop	Cow	3.5 b	6.3	7.2 a	9.6 a
	Goat	4.7 a	6.0	7.3 a	9.6 a
	No manure	3.1 b	5.1	6.3 ab	8.7 ab
Monocrop	Cow	4.0 ab	6.3	7.2 a	9.4 a
	Goat	3.5 b	6.0	6.8 ab	8.9 ab
	No manure	3.2 b	4.7	5.9 b	7.9 b

\*Means in a column not followed by a letter within a variable are not significantly different according to F-test at  $p \leq 0.05$ .


\*\*Means in a column followed by the same letter within a variable are not significantly different according to Tukey's honestly significant difference test at  $p \leq 0.05$

### 3.3 Effects of Slender Leaf Intercrop and Fresh Organic Manure on Tomato Yield

Tomato plants grown with fresh organic manure and slender leaf intercrop had significantly higher fruit yield in terms of number of fruits per plant and fresh fruit weight per plant compared to the control treatment (Table 3). Growing tomato plants using fresh organic manure and slender leaf intercrop increased the number of tomato fruits per plant and fresh fruit weight per plant by 22.2–49.7% and 24.5–80.4%, respectively in comparison with the control treatment. Generally, the highest fruit number and total fruit weight per plant was obtained on plants grown with fresh goat dung and slender leaf intercrop while the lowest yields were obtained under the control treatment. In terms of fruit numbers among the other treatments, plants grown with fresh goat dung gave higher fruit numbers than those grown with fresh cow dung and slender leaf intercrop,

followed by those grown with fresh cow dung, and the least number of fruits per plant obtained on plants grown with a slender leaf intercrop alone. On the other hand, fresh fruit weight per plant was higher on plants grown with fresh cow dung and slender leaf intercrop followed by those grown with fresh goat dung, then those grown with fresh cow dung with the lowest fresh weight obtained from plants grown without fresh manure (Table 3).

Comparing the two cropping regimes, tomato intercropped with slender leaf had more fruits (5.9%) and higher fruit weight per plant (17.0%) compared to tomato grown as a monocrop (Table 3). Similarly, growing tomato with fresh organic manure significantly enhanced tomato fruit yield in terms of fruit number (fresh cow dung 22.1% and fresh goat dung 32.4%) and fresh fruit weight per plant (fresh cow dung 33.9% and goat dung 47.8%) compared to when the plants were grown without application of fresh organic manure (Table 3).



**Fig. 2. Main effects of slender leaf intercrop and fresh organic manure on growth variables; (A) Plant height, (B) Collar diameter, (C) Number of internodes and (D) Number of branches during tomato production. Means followed by the same letter within a variable are not significantly different according to Tukey's honestly significant difference test at  $p \leq 0.05$**

**Table 3. Effects of slender leaf intercrop and fresh organic manure on yield of tomato during tomato production**

Manure	Cropping regime		Monocrop		Manure Means	Increase (%)
	Intercrop	Increase (%)		Increase (%)		
	Number of fruits (no./plant)					
Cow	32.5ab*	33.1**	33.7ab	38.1	33.1h	22.1
Goat	36.5a	49.7	35.2ab	44.4	35.9h	32.4
No manure	29.8ab	22.2	24.4b		27.1i	
Cropping regime Means	32.9e		31.1e			
Increase (%)	5.9					
	Fresh fruit weight (kg/plant)					
Cow	1.6ab	58.8	1.5ab	43.1	1.5h	33.9
Goat	1.8a	80.4	1.6ab	52.9	1.7h	47.8
No manure	1.3ab	24.5	1.0b		1.2i	
Cropping regime Means	1.6e		1.4e			
Increase (%)	17.0					

\*Means followed by the same letter within a letter series and within a variable are not significantly different according to Tukey's honestly significant difference test at  $p \leq 0.05$ .

\*\*Percent increased calculated in reference to yield of the control treatment (No slender leaf (monocrop), no fresh manure).

#### 4. DISCUSSION

In the current study, use of fresh organic manure and slender leaf intercrop proved of potential benefit in tomato production. Root-knot nematodes both in the soil and on the roots of tomato plant were lower under fresh organic manure and slender leaf intercrop treatments than in the control treatment. Addition of amendments in soil enhances microbial activities and increases number of antagonists [19]. According to Mohammed and Alan [20], organic manure such as livestock waste stimulates the generation of predators and parasites of plant parasitic nematode. Suppression of *Meloidogyne* spp. populations by fresh organic manure in the current study could thus have been a result of toxic effects during decomposition and increased microbial activities in the soil leading to a poor host environment for the root-knot nematodes thus inhibiting their penetration into the roots. Suppression of *Meloidogyne* spp. by fresh organic manure could also have been due to increased number of root-knot nematode antagonists generated during decomposition of fresh organic manure in the soil. According to Nahar et al. [14], fresh organic manures can be more effective to control root-knot nematodes compared to composted manures since toxic compounds can quickly reach threshold level required to control nematodes with rapid increase in number of beneficial microorganisms

and microbial activities. Similar to the findings of our study, studies by Abolusoro and Abolusoro [10] reported a significant reduction in the soil population of root-knot nematode and root gall index by organic manure amendment compared with the control treatment.

Similar to the effects of fresh organic manure on the population of juvenile root-knot nematodes in the soil and root galls on tomato roots, intercropping tomato with slender leaf reduced juvenile root-knot nematode populations and number of nematode root-knot galls compared to when tomato was grown as a monocrop. These findings corroborate findings by [3] who reported significantly lower number of juvenile nematodes in the soil and root galls in tomato plants when *Crotalaria* plants were used as intercrop. According to Wang et al. [8], *Crotalaria* intercrop suppresses plant-parasitic nematodes by enhancing microbial activities against *Meloidogyne* species. Some compounds of microbial origin exuded by *Crotalaria* roots disrupt biological activities of root-knot nematodes that are essential for the successful plant-nematode interaction [21].

From the results of the current study, combined use of fresh organic manure and slender leaf intercrop registered higher reduction in root-knot nematode (juvenile) population in the soil and the number of galls on the roots of tomato plants.

According to Ravindra et al. [22], combination of treatments is more effective than individual treatment in management of root-knot nematodes. The combined diverse mechanisms could reduce egg hatching factors, enhancing plant growth, inducing systemic resistance in the plant, alter root exudates and inhibit penetration of nematodes into the roots as well as reduced galling [23]. In the current study, higher suppression of root-knot nematode achieved with combined use of fresh goat dung and slender leaf intercrop could possibly be as a result of production of more root-knot nematode predatory microorganism and other competing beneficial organisms by the fresh goat dung manure and slender leaf intercrop as well as increased plant tolerance or resistance to nematode infection through enhanced plant growth. As noted in the results, treatments with lower number of juvenile root-knot nematode populations registered fewer number of root galls in the current study. This implies that root-knot nematodes stimulate formation of root galls. Studies by Kankam and Adomako [24] reported a strong positive correlation of nematode juveniles and root galling, where the root galling on plant root increase with increase in nematode population in the rhizosphere of the plant.

Growing tomato plants with application of fresh organic manure and slender leaf intercrop resulted in significantly taller and thicker stems and plants with more internodes and branches. Better plant growth observed on tomato grown under these treatments in the current study could be as a result of lower number of root-knot nematodes and root galls registered under these treatments leading to the better crop performance. On the other hand, the reduced plant growth under the control treatment could partly have been as a result of high number of root galls which could have hindered plant water and nutrients supply by the roots. Root galls caused by root-knot nematode leads to vascular damage which disturbs water and mineral uptake resulting in severe reduction in plant growth [25]. Besides, improved plant growth in fresh manure and slender leaf intercrop treatments could also be associated with increased supply of plant nutrients following application of fresh organic manure [26] and use of *Crotalaria* plants [8]. Apart from improving soil fertility, *Crotalaria* plants competes with weeds without becoming a weed and grows vigorously to provide good ground coverage [8] which may have contributed to better performance of tomato plants intercropped with slender leaf.

Increased tomato yield recorded under fresh organic manure and slender leaf intercrop in the current study could be associated with the reduced root-knot nematode and enhanced tomato growth recorded under these treatments. Increased tomato yield following application of fresh organic manure in the current study could be attributed to better nutrient availability and growth promoting substances leading to higher fruit set percent. According to Soni et al. [27], application of organic nutrient sources increases activities of beneficial microorganisms due to increased organic pool in the soil, resulting in production of growth promoting substances and improved nutrient availability. Good ground coverage by slender leaf plant could also have led to reduced evapotranspiration rate resulting to improved soil moisture status as well as enhanced uptake of nutrients resulting to higher photosynthesis. Increased photosynthesis increases the assimilation rates leading to more food being manufactured and translocated to active sinks thus promoting vigorous growth and production of more fruits [28].

## 5. CONCLUSIONS

Based on the results, we conclude that the combined use of fresh organic manure and slender leaf intercrop is a potential strategy for use in suppressing root-knot nematode populations with a resultant increase in growth and yield of tomato. The present study forms a basis for advocating for the use of fresh organic manure and slender leaf as an effective method to control root-knot nematodes in field grown tomato. Among the treatments tested, combined use of fresh goat dung and slender leaf intercrop in tomato production stands to benefit tomato growers more in terms of higher suppression of root-knot nematode and higher tomato fruit yield. Based on the findings of our study, we recommend the use of fresh goat dung and slender leaf intercrop in management of root-knot nematode in tomato production within regions with similar climatic conditions to those of our study site. In addition, further studies on the use of these technologies using different tomato varieties and in different tomato growing climatic conditions to those of the current study are recommended to further validate the results.

## ACKNOWLEDGEMENTS

This is part of Doctor of Philosophy Degree in Horticulture project by James Ngelenzi Munywoki. The study was made possible by the

support of the African Development Bank (AfDB) through the Ministry of Agriculture - Kenya. We wish to acknowledge the Crop Molecular Biology and Biotechnology Laboratory of the Confucius Institute, Department of Crops, Horticulture and Soils, Egerton University, where part of the study was conducted. We also acknowledge the African Center of Excellence in Sustainable Agriculture and Agribusiness Management (CESAAM), Egerton University for facilitating the processing of research funds that made implementation of the study possible.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Kaşkavalci G. Effects of soil solarization and organic amendment treatments for controlling *Meloidogyne incognita* in tomato cultivars in Western Anatolia. Turk. J. Agric. For. 2007;31(3):159-167.
2. Shurtleff MC, Averre CW. Diagnosing plant diseases caused by nematodes. American Phytopathological Society (APS Press); 2000.
3. Bello TT, Fawole B, Claudius-Cole A. Management of root-knot nematodes (*Meloidogyne* spp) on tomato using antagonistic plants. J. Biol. agric. Healthcare. 2014;4(24):97-101.
4. Sikora RA, Fernandez E. Nematode parasites of vegetables. In: Luc M, Sikora RA, and J Bridge (Eds.). Plant parasitic nematodes in subtropical and tropical agriculture (2<sup>nd</sup> edition). 2005;319-392. CABI publishing.
5. Idorenyin AU, Ugwuoke KI. Pathogenicity of *Meloidogyne incognita* race 1 on turmeric (*Curcuma longa* L.) as influenced by inoculum density and poultry manure amendment. Plant Pathol. J. 2010;9(4): 162-168.
6. Singh SK, Khurma UR. Susceptibility of six tomato cultivars to the root-knot nematode, *Meloidogyne incognita*. S. Pac. J. Nat. Sci. 2007;25(1):73-77.
7. Mashela PW, Shimelis HA, Mudau FN. Comparison of the efficacy of ground wild cucumber fruits, aldcarb and fenamiphos on suppression of *Meloidogyne incognita* in tomato. J. Phytopathol. 2008;156(5): 264-267.
8. Wang KH, Sipes BS, Schmitt DP. Crotalaria as a cover crop for nematode management: A review. Nematropica. 2002;32:35-57.
9. Mousavi SR, Eskandari H. A general overview on intercropping and its advantages in sustainable agriculture. J. Appl. Environ. Biol. Sci. 2011;1(11):482-486.
10. Abolusoro SA, Abolusoro PF. Effects of organic manure types on the growth, yield as well as root and soil populations of root-knot nematodes (*Meloidogyne incognita*) of tomato. Sci. J. Agric. 2012;1(5):138-144.
11. Bonanomi G, Antignani V, Capodilupo M, Scala F. Identifying the characteristics of organic soil amendments that suppress soilborne plant diseases. Soil Biol. Biochem. 2010;42(2):136-144.
12. Summers H. Effects of organic manure on nematode control. J. Dis. Pest Control in Tropics. 2011;7(2):190-191.
13. Widmer TL, Abawi GS. Relationship between levels of cyanide in Sudan grass hybrids incorporated into soil and suppression of *Meloidogyne hapla*. J. Nematol. 2002;34(1):16-22.
14. Nahar MS, Grewal PS, Miller SA, Stinner D, Stinner BR, Kleinhenz MD, et al. Differential effects of raw and composted manure on nematode community, and its indicative value for soil microbial, physical and chemical properties. Appl. Soil Ecol. 2006;34(2-3):140-151.
15. Koul O, Dhaliwal GS, Cuperus GW. Integrated pest management potential, constraints and challenges. CABI, Wallingford, UK; 2004.
16. Jaetzold R, Schmidt H, Hornetz B, Shisanya CA. Farm management handbook of Kenya. Natural conditions and farm information, Vol.11/C, 2nd ed. Ministry of Agriculture/Deutsche Gesellschaft für Technische Zusammenarbeit GmbH, German Agency for Technical Cooperation, Nairobi, Kenya; 2006.
17. Horticultural Crops Development Authority, HCDA. Fruits and Vegetables. Agricultural Information Resource Centre. Nairobi, Kenya; 2006.
18. Coyne DL, Nicol JM, Claudius-Cole B. Practical plant nematology: a field and laboratory guide (3<sup>rd</sup> edition). International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria; 2018.

19. Wachira PM, Kimenju JW, Okoth SA, Mibey RK. Stimulation of nematode-destroying fungi by organic amendments applied in management of plant parasitic nematode. *Asian J. Plant Sci.* 2009; 8(2):153- 159.
20. Mohammed A, Alan MM. Utilization of waste material in nematode control. *Indian Phytopathol.* 2003;156:264-267.
21. Varkey S, Anith KN, Narayana R, Aswini S. A consortium of rhizobacteria and fungal endophyte suppress the root-knot nematode parasite in tomato. *Rhizosphere.* 2018;5:38-42.
22. Ravindra H, Sehgal M, Pawan AS, Archana BS, Shruti SA, Narasimhamurthy HB. Eco-friendly management of root-knot nematodes using acacia compost and bioagents in brinjal. *Pak. J. Nematol.* 2014; 32(1):33-38.
23. Khalil MS, Kenawy A, Gohrab MA, Mohammed EE. Impact of microbial agents on *Meloidogyne incognita* management and morphogenesis of tomato. *J. Biopestic.* 2012;5(1):28-35.
24. Kankam F, Adomako J. Influence of inoculum levels of root knot nematodes (*Meloidogyne* spp.) on tomato (*Solanum lycopersicum* L.). *Asian J. Agric. Food Sci.* 2014;2(2):171-178.
25. Abolusoro SA, Abolusoro PF, Mathew FO, Izuogu NB. Effects of organic and inorganic manures on the growth attributes of root-knot nematode (*Meloidogyne incognita*) infected Ethiopian egg plant (*Solanum aethiopicum*). *World J. Agric. Res.* 2013;1(6):104-107.
26. John LW, Jamer DB, Samuel LT, Warner LW. *Soil fertility and fertilizers: An introduction to nutrient management.* Pearson Education, Delhi, India. 2004;106-153.
27. Soni S, Kanawjia A, Chaurasiya R, Chauhan PS, Kumar R, Dubey S. Effect of organic manure and biofertilizers on growth, yield and quality of Strawberry (*Fragaria x ananassa* Duch) cv. Sweet Charlie. *J. Pharmacogn. Phytochem.* 2018; 2:128-132.
28. Dauda SN, Ajayi FA, Ndor E. Growth and yield of water melon (*Citrullus lanatus*) as affected by poultry manure application. *Elec. J. Env. Agric. Food Chem.* 2009;8 (4):305-311.

© 2022 Munywoki et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/78865>