



## **Application of Sulphur Levels and Different Micronutrients on Yield and its Attributing Traits of Garlic (*Allium sativum* L.)**

**Praveen Choyal<sup>a\*</sup>, O. P. Garhwal<sup>a</sup>, M. R. Choudhary<sup>a</sup>, Manju Netwal<sup>a</sup> and Kamal Mahala<sup>a</sup>**

<sup>a</sup> *Department of Horticulture, SKN College of Agriculture, Jobner, Jaipur, India.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJPSS/2022/v34i530861

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83163>

**Original Research Article**

**Received 02 December 2021**  
**Accepted 05 February 2022**  
**Published 21 February 2022**

### **ABSTRACT**

The field experiment was conducted at the Horticulture Farm, Sri Karan Narendra College of Agriculture, Jobner (Rajasthan) during 2019-20 and 2020-21. The experiment consisted of 20 treatment combinations with four soil application of sulphur (S<sub>0</sub>-control, S<sub>1</sub>-sulphur 20 kg/ha, S<sub>2</sub>-sulphur 40 kg/ha and S<sub>3</sub>-sulphur 60 kg/ha) and five levels of foliar application of micronutrients (M<sub>0</sub>-Control, M<sub>1</sub>-zinc sulphate @ 0.6%, M<sub>2</sub>-ferrous sulphate @ 0.2%, M<sub>3</sub>-borax @ 0.5 and M<sub>4</sub>-Ammonium molybdate @ 0.5%) in factorial randomized block design with three replications. The results of the study have clearly indicated that among sulphur levels, application of sulphur 60 kg/ha significantly improved growth parameters viz., neck thickness, number of cloves per plant, clove length, clove girth, polar diameter, equatorial diameter, enhanced significantly by application of sulphur 60 kg/ha (S<sub>3</sub>) in both the years as well as in pooled analysis. Similarly, among micronutrients foliar application of zinc sulphate @ 0.6% significantly influenced all the growth, yield attributes viz., neck thickness, number of cloves per plant, clove length, clove girth, polar diameter, equatorial diameter of garlic in both the years as well as in pooled analysis.

**Keywords:** *Garlic; sulphur; micronutrients; clove length; diameter of bulb.*

## 1. INTRODUCTION

Garlic is an important bulbous plant and used throughout India primarily as a spice or condiment. It is botanically known as *Allium sativum* L. is member of the genus *Allium*, which comprises approximately 750 species belonging to the family Alliaceae. It is one of the most important bulbous vegetable crops and is next to onion [1]. It is originated from Central Asia and later spread to Mediterranean region [2,3].

The bulb of garlic is compound in nature, consisting of numerous bulblets, so called as cloves, of different size, the whole surrounded by layers of white scale leaves. Allicin is the main biologically active component of freshly crushed garlic cloves, which is produced by the degradation of alliin, from results of alliinase activity [4,5]. It has many medicinal properties. It lowers blood cholesterol levels and antiplatelet aggregation, produces anti-inflammatory activity and inhibits cholesterol synthesis. Moreover, it has long been known to have antibacterial, antifungal, anticancer, antioxidant and antiviral activities [6].

Nutrients is a product of the magnitude of the effects of crop yields on each area. The plant needs nutrients that are essential for normal functioning and growth. The shortage of micronutrients over the past three decades has been a major obstacle to the production and production of vegetables in general and in particular. Therefore, there is an urgent need to correct individual deficiencies and to prevent further spread. The low production of Indian garlic is mainly due to the cultivation of low-yield / hybrid varieties, vulnerable to both biotic and abiotic factors. Therefore, a balanced diet is considered one of the major abiotic factors that negatively affect the growth and yield of garlic [7].

Application of boron can increase bulb size, number of cloves per bulb and yield of garlic [8]. Zinc is crucial for plant growth because it controls the synthesis of indole acetic acid, which noticeably regulates plant growth and also active many enzymatic reactions which are necessary for chlorophyll synthesis and carbohydrate formation [9]. Iron is an essential micronutrient for almost all living organisms because it plays critical role in metabolic processes such as DNA synthesis, respiration and photosynthesis. Further, many metabolic pathways are also activated by iron and it is a prosthetic group constituent of many enzymes. An imbalance

between the solubility of iron in soil and the demand for iron by the plant are the primary causes of iron chlorosis. Molybdenum is also an important micronutrient for plants, which plays a vital role in enzymes activity as nitrogenase, catalase and peroxidase [10,11].

Among the macronutrients, sulphur is one of the essential elements for building up pungency in garlic is attributed to presence of an alkaloid "Di allyl disulphide" in which sulphur is prime component. The application of sulphur in garlic is not only important from nutrient point of view but also it builds resistance in plants against pest and diseases. Its role in balanced fertilization and consequently in yield and quality improvement of garlic is being increasingly appreciated [12].

## 2. MATERIALS AND METHODS

The field experiment "Application of Sulphur Levels and Different Micronutrients on Yield and its Attributing traits of Garlic (*Allium sativum* L.)" was conducted at the Horticulture Farm, Sri Karan Narendra College of Agriculture, Jobner, Jaipur (Rajasthan) during *Rabi* seasons 2019-20 and 2020-21. Jobner is situated 45 km in West of Jaipur at 26°5' North latitude, 75°20' East longitude and at an altitude of 427 meters above mean sea level. This region falls under Agro-Climatic Zone-III A (Semi- Arid Eastern Plain Zone) of Rajasthan.

Jobner is belong to semi-arid region characterized by extremes of temperature both in summer and winter, low rainfall and moderate relative humidity. The annual average rainfall varies between 250 to 500 mm year<sup>-1</sup>, most of which is received in rainy season fall during July to early September, sporadic showers also received in winters. The maximum temperature ranges from 30 to 46°C during month of May and June, while in December and January, it falls down below -1°C and evaporation ranges from 1.2-6.9 mm per day. The experimental field is sand loam in texture, alkaline in reaction (pH 8.24) and poor in organic matter (0.11%) nitrogen (132.15 kg/ha), phosphorus (17.64 kg/ha) but medium in potassium content (161.80 kg/ha).

The experiment was laid out in factorial randomized block design with three replications consisting of twenty treatment combinations including five micronutrients viz; control (water spray), borax at 0.5%, zinc sulphate at 0.6%, ferrous sulphate at 0.2% and ammonium molybdate at 0.5% and four sulphur levels

(control, 20, 40 and 60 kg/ha). The sulphur was applied as soil application just before sowing and micronutrients (Zn, B, Fe and Mo) as foliar spray at 40 DAS. The recommended dose of NK for garlic was applied @ 120: 100 kg/ha, respectively. Full dose of potassium and half dose of nitrogen were applied as basal dose just before sowing and rest half dose of nitrogen was applied as top dressing in two split doses. The plot size was 1.5 × 1.2 m<sup>2</sup> (1.80 m<sup>2</sup>) and seed are sowing at spacing of 15 cm × 10 cm. The seeds of cv. G -282 procured from National Horticultural Research & Development Foundation in Karnal, (Haryana). The cloves of garlic were treated with Carbendazim @ 2 g/kg seed to control seed borne diseases. The seeds was sown on 16<sup>th</sup> October, 2019 and 22<sup>th</sup> October, 2020 manually with a seed rate of 500 kg /ha in row at 15 cm apart. All the cultural operations were followed continuously during crop growth. The observations for plant growth parameters and yield parameters like neck thickness, polar diameter, equatorial diameter, number of cloves per bulb, clove length and girth are taken after harvesting of crop.

### 3. RESULTS AND DISCUSSION

#### 3.1 Effect of Sulphur Levels

The data pertaining to the effect of various levels of sulphur on yield parameters of garlic revealed

that all the treatments significantly influenced the neck thickness, polar diameter, equatorial diameter, number of cloves per bulb, clove length and girth of garlic [Table 1, 2 and 3] during both the years and in pooled mean analysis. The maximum neck thickness (0.750, 0.795 and 0.773), polar diameter (4.96, 5.12 and 5.04), equatorial diameter (3.79, 3.94 and 3.87), number of cloves per bulb (17.88, 17.00 and 17.44), clove length (2.79, 2.91 and 2.85) and clove girth (1.04, 1.08 and 1.06) were found in treatment S<sub>3</sub> (Sulphur -60 kg/ha) in both the years and pooled mean respectively. However, the treatment S<sub>3</sub> (sulphur- 60 kg/ha) was found statistically at par to S<sub>2</sub> (Sulphur- 40 kg/ha). This might be due to magnificent role of sulphur is a key nutrient in garlic production; therefore, lack of its optimum supply in different plant parts limits the crop quality and also had poor utilization of macro and micronutrients [12]. These results are agreement with findings of [13] in garlic, [13] in onion and garlic, [14,15] in garlic, [16,17] in onion.

#### 3.2 Effect of Micronutrients

Data indicated that application of various micronutrients also had significant effect on the yield parameters of garlic during both the years and in pooled mean analysis [Tables 1, 2 and 3].

**Table 1. Effect of sulphur and micronutrients on number of cloves per bulb and neck thickness of garlic**

Treatments	Number of cloves per bulb			Neck thickness (cm)		
	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled
<b>A. Sulphur</b>						
S <sub>0</sub> (Control)	16.02	15.53	15.78	0.630	0.639	0.635
S <sub>1</sub> (Sulphur 20 kg/ha)	17.38	16.46	16.92	0.690	0.735	0.713
S <sub>2</sub> (Sulphur 40 kg/ha)	17.81	16.82	17.32	0.730	0.779	0.755
S <sub>3</sub> (Sulphur 60 kg/ha)	17.88	17.00	17.44	0.750	0.795	0.773
<b>SEm±</b>	<b>0.21</b>	<b>0.20</b>	<b>0.14</b>	<b>0.013</b>	<b>0.015</b>	<b>0.010</b>
<b>CD (P=0.05)</b>	<b>0.59</b>	<b>0.57</b>	<b>0.40</b>	<b>0.037</b>	<b>0.043</b>	<b>0.028</b>
<b>B. Micronutrients</b>						
M <sub>0</sub> (Control)	16.61	15.77	16.19	0.625	0.653	0.639
M <sub>1</sub> (Zinc sulphate @ 0.6%)	17.54	16.75	17.14	0.749	0.784	0.767
M <sub>2</sub> (Ferrous sulphate @ 0.2%)	17.39	16.56	16.97	0.712	0.752	0.732
M <sub>3</sub> (Borax @ 0.5%)	17.49	16.71	17.10	0.731	0.772	0.752
M <sub>4</sub> (Ammonium molybdate @ 0.5%)	17.33	16.48	16.91	0.683	0.723	0.703
<b>SEm±</b>	<b>0.23</b>	<b>0.22</b>	<b>0.16</b>	<b>0.015</b>	<b>0.017</b>	<b>0.011</b>
<b>CD (P=0.05)</b>	<b>0.66</b>	<b>0.63</b>	<b>0.45</b>	<b>0.042</b>	<b>0.048</b>	<b>0.031</b>

**Table 2. Effect of sulphur and micronutrients on diameter of bulb (polar and equatorial) of garlic**

Treatments	Polar diameter (cm)			Equatorial diameter (cm)		
	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled
<b>A. Sulphur</b>						
S <sub>0</sub> (Control)	4.23	4.28	4.26	3.19	3.33	3.26
S <sub>1</sub> (Sulphur 20 kg/ha)	4.55	4.68	4.62	3.45	3.60	3.53
S <sub>2</sub> (Sulphur 40 kg/ha)	4.86	5.02	4.94	3.69	3.86	3.78
S <sub>3</sub> (Sulphur 60 kg/ha)	4.96	5.12	5.04	3.79	3.94	3.87
<b>SEm±</b>	<b>0.10</b>	<b>0.12</b>	<b>0.08</b>	<b>0.07</b>	<b>0.08</b>	<b>0.05</b>
<b>CD (P=0.05)</b>	<b>0.28</b>	<b>0.33</b>	<b>0.21</b>	<b>0.20</b>	<b>0.23</b>	<b>0.15</b>
<b>B. Micronutrients</b>						
M <sub>0</sub> (Control)	4.28	4.37	4.32	3.26	3.37	3.32
M <sub>1</sub> (Zinc sulphate @ 0.6%)	4.89	4.99	4.94	3.71	3.87	3.79
M <sub>2</sub> (Ferrous sulphate @ 0.2%)	4.68	4.83	4.76	3.54	3.71	3.63
M <sub>3</sub> (Borax @ 0.5%)	4.79	4.89	4.84	3.62	3.81	3.71
M <sub>4</sub> (Ammonium molybdate @ 0.5%)	4.62	4.79	4.70	3.51	3.65	3.58
<b>SEm±</b>	<b>0.11</b>	<b>0.13</b>	<b>0.09</b>	<b>0.08</b>	<b>0.09</b>	<b>0.06</b>
<b>CD (P=0.05)</b>	<b>0.32</b>	<b>0.37</b>	<b>0.24</b>	<b>0.23</b>	<b>0.25</b>	<b>0.17</b>

**Table 3. Effect of sulphur and micronutrients on clove length and girth of garlic**

Treatments	Clove length (cm)			Clove girth (cm)		
	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled
<b>A. Sulphur</b>						
S <sub>0</sub> (Control)	2.39	2.45	2.42	0.88	0.90	0.89
S <sub>1</sub> (Sulphur 20 kg/ha)	2.57	2.64	2.61	0.96	0.98	0.97
S <sub>2</sub> (Sulphur 40 kg/ha)	2.71	2.80	2.76	1.01	1.04	1.03
S <sub>3</sub> (Sulphur 60 kg/ha)	2.79	2.91	2.85	1.04	1.08	1.06
<b>SEm±</b>	<b>0.05</b>	<b>0.05</b>	<b>0.03</b>	<b>0.02</b>	<b>0.02</b>	<b>0.01</b>
<b>CD (P=0.05)</b>	<b>0.13</b>	<b>0.15</b>	<b>0.10</b>	<b>0.05</b>	<b>0.06</b>	<b>0.04</b>
<b>B. Micronutrients</b>						
M <sub>0</sub> (Control)	2.39	2.45	2.42	0.89	0.91	0.90
M <sub>1</sub> (Zinc sulphate @ 0.6%)	2.78	2.88	2.83	1.04	1.07	1.05
M <sub>2</sub> (Ferrous sulphate @ 0.2%)	2.63	2.71	2.67	0.97	1.01	0.99
M <sub>3</sub> (Borax @ 0.5%)	2.73	2.83	2.78	1.01	1.04	1.02
M <sub>4</sub> (Ammonium molybdate @ 0.5%)	2.56	2.63	2.59	0.96	0.97	0.97
<b>SEm±</b>	<b>0.05</b>	<b>0.06</b>	<b>0.04</b>	<b>0.02</b>	<b>0.02</b>	<b>0.01</b>
<b>CD (P=0.05)</b>	<b>0.15</b>	<b>0.16</b>	<b>0.11</b>	<b>0.05</b>	<b>0.06</b>	<b>0.04</b>

The foliar spray of zinc sulphate-0.6 % (M<sub>1</sub>) registered maximum neck thickness (0.749, 0.784 and 0.767), polar diameter (4.89, 4.99 and 4.94), equatorial diameter (3.71, 3.87 and 3.79), number of cloves per bulb (17.54, 16.75 and 17.14), clove length (2.78, 2.88 and 2.83) and clove girth (1.04, 1.07 and 1.05) which were significantly higher over rest of the treatments except M<sub>2</sub> (ferrous sulphate @ 0.2%), M<sub>3</sub> (borax @ 0.5%) and M<sub>4</sub> (ammonium molybdate @ 0.5%) in both the years and pooled mean analysis which was found statistically at par to it.

Zinc is crucial for plant growth because it controls the synthesis of indole acetic acid, which noticeably regulates plant growth and also active many enzymatic reactions which are necessary for chlorophyll synthesis and carbohydrate formation [8]. Application of zinc also plays a role to increase the activity of nitrate reductase enzyme and enhanced synthesis of certain amino acids and protein. The results are also supported by [18] in tomato, [19] in okra, [20] in garlic, [21] in onion.

#### 4. CONCLUSION

Based on the results of two years experiments, it may be concluded that soil application of sulphur at 60 kg/ha combined with foliar spray of Zinc sulphate at 0.6% proved the most superior treatment combination in garlic fetching the significantly higher yield and its attributing traits. Although, sulphur application at 40 kg/ha along with foliar application of zinc sulphate at 0.6 % was found at par to it.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Hamma IL, Ibrahim U, Mohammed AB. Growth, yield and economic performance of garlic (*Allium sativum* L.) as influenced by farm yard manure and spacing in Zaria, Nigeria. *Journal of Agricultural Economics and Development*. 2013;2(1):001-005.
2. Simon W. The origin and distribution of garlic. *USDA Vegetable Crops Research Unit, USA*. 2001;1-3.
3. Kigori JM, Magaji MD, Yakudu AI. Productivity of two garlic (*Allium sativum* L.) cultivars as affected by different levels of nitrogenous and phosphorus fertilizers in Sokoto, Nigeria. *Proceeding of 41<sup>st</sup> Annual Conference on Bulletin of the Science Association of Nigeria, Usmanu Danfodiyo University, Sokoto*; 2005.
4. Bocchini P, Andalò C, Pozzi R, Galletti GC, Antonelli A. Determination of diallyl thiosulfinate (allicin) in garlic (*Allium sativum* L.) by high performance liquid chromatography with a post column photochemical reactor. *Analytica Chimica Acta*. 2001;441:37-43.
5. Rahman MM, Fazlic V, Saad NW. Antioxidant properties of raw garlic (*Allium sativum* L.) extract. *International Food Research Journal*. 2012;19:589-591.
6. Lawrence R, Lawrence K. Antioxidant activity of garlic essential oil (*Allium sativum* L.) grown in north Indian plains. *Asian Pacific Journal of Tropical Biomedicine*. 2011;1:51-54.
7. Ballabh K, Rana DK, Rawat SS. Effects of foliar application of micronutrients on growth, yield and quality of onion. *Indian Journal of Horticulture*. 2013;70(2):260-265.
8. Rani P, Panghal VPS, Rana MK, Duhan DS. Response of garlic to foliar application of urea and micronutrients. *International Journal of Tropical Agriculture*. 2015;33(4):2845-2849.
9. Vitosh ML, Silva GH. A rapid petiole sap nitrate test for potatoes. *Comm. In soil Science and plant Analysis*. 1994;25(3):183-190.
10. Marschner H. Mineral nutrition of higher plant. 2<sup>nd</sup> (ed.), Academic Press Limited. Text Book. 1995;864.
11. Tisdale SL, Nelson WL, Beaton JD. *Soil Fertility and Fertilizers*, 4th edition. Macmillan Publication Company, New York; 1985.
12. Campbell WH. Nitrate reductase structure, function and regulation. *Binding the gap between biochemistry and physiology. Annual Review of Plant Physiology and Plant Molecular Biology*. 1999;50:277-303.
13. Kumar A, Singh O. Role of sulphur in nutrient utilization and catalase activity in onion crop. *Indian Journal Agricultural Research*. 1994;28:15-19.
14. Srinidhi N. Studies on sulphur nutrition on onion and garlic in sulphur deficient, research report. *University of Agricultural Sciences, Dharwad*; 2000.
15. Jaggi RC. Sulphur levels and sources affecting yield and yield attributes in onion (*Allium cepa* L.). *Indian Journal of Agricultural Sciences*. 2005;75(3):154-156.
16. Verma D, Singh H. Response of varying levels of potassium and sulphur on yield and uptake of nutrients by onion. *Annals of Plant and Soil Research*. 2012;14(2):143-146.
17. Singh SK, Kumar M, Seema Singh PK, Yadav LM. Effect of sulphur sources and levels on growth, yield and quality of onion. *Current Journal of Applied Science and Technology*. 2019;33(2):1-4.
18. Raghavendra BH, Umamaheswarappa P, Srinivasa V, Salimath S, Hanumantappa M. Effect of different sources and levels of sulphur on yield and yield attributes in onion under central dry zone of Karnataka. *International Journal of Ecology and Environmental Sciences*. 2020;2(4):766-768.
19. Vitosh ML, Silva GH. A rapid petiole sap nitrate test for potatoes. *Comm. In soil Science and plant Analysis*. 1994;25(3): 183-190.

20. Patnaik MC, Raj GB, Reddy IP. Response of tomato (*Lycopersicon esculentum* L.) to zinc and iron. *Vegetable Science*. 2001; 28(1):78-79.
21. Kumar M, Sen NL. Effect of zinc, boron and GA<sub>3</sub> on yield of okra (*Abelmoschus esculentus* L.). *Indian Journal of Horticulture*. 2005;62(3):308-309.

© 2022 Choyal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/83163>